

06/14/99  
Jc573 U.S. PTO

**NON-PROVISIONAL  
UTILITY PATENT APPLICATION  
TRANSMITTAL - 37 CFR 1.53(b)**

☐ Duplicate  
(check, if applicable)

Jc490 U.S. PTO  
09/333159  
06/14/99

Assistant Commissioner for Patents  
**BOX PATENT APPLICATION**  
Washington, DC 20231

**Attorney Docket No. 10147-6**  
**First Named Inventor: Sean A. McCarthy**  
**Express Mail Label No. EL174022173US**  
**Total Pages of Transmittal Form: 2**

Transmitted herewith for filing is the **non-provisional** utility patent application entitled:

**NOVEL GENES ENCODING PROTEINS HAVING DIAGNOSTIC,  
PREVENTIVE, THERAPEUTIC, AND OTHER USES**

which is:

an ☒ Original; or

a ☐ Continuation, ☐ Divisional, or ☐ Continuation-in-part (CIP)  
of prior Application No. \_\_\_\_/\_\_\_\_, filed \_\_\_\_.

☐ This non-provisional patent application is based on Provisional Patent Application  
No. \_\_\_\_\_, filed \_\_\_\_\_.

Enclosed are:

- ☒ Specification (including Abstract) and claims: 148 pages.
- ☒ Unexecuted copy of Declaration.
- ☐ Copy of Declaration from prior application.
- ☐ Separate Power of Attorney (including 37 CFR 3.73(b) statement, if applicable).
- ☒ 53 sheets of drawings (formal) plus one copy.
- ☐ Microfiche computer program (Appendix).
- ☒ Nucleotide and/or Amino Acid Sequence Submission, including:
  - ☒ Computer readable copy ☒ Paper Copy ☒ Verified Statement.
- ☐ Under PTO-1595 cover sheet, an assignment of the invention.
- ☐ Certified copy of \_\_\_\_\_ Application No. \_\_\_\_\_, filed \_\_\_\_\_, is filed:
  - ☐ herewith or ☐ in prior application \_\_\_\_\_.
- ☐ Verified Statement Claiming Small Entity Status under 37 CFR 1.9 and 1.27.
  - ☐ was filed in the prior non-provisional application, and such status is still proper and desired (37 CFR 1.28(a));
  - ☐ is enclosed herewith; ☐ is no longer desired.
- ☐ Preliminary Amendment.
- ☐ Information Disclosure Statement, PTO-1449, and cited references.
- ☐ Other:

The filing fee has been calculated as shown below:

			SMALL ENTITY			LARGE ENTITY	
CLAIMS	NO. FILED	NO. EXTRA	BASIC FEE: \$380.			BASIC FEE: \$760	
Total	23-20 =	3	X9	\$	OR	X18	\$54.00
Independent	3-3=	0	X39	\$	OR	X78	\$-0-
Multiple Dependent Claims Present			\$130	\$	OR	\$260	\$
			<b>TOTAL</b>	<b>\$</b>	<b>OR</b>	<b>TOTAL</b>	<b>\$-0-*</b>

\*THE ABOVE CALCULATED FEE IS NOT BEING PAID AT THIS TIME

The Commissioner is hereby authorized to charge payment of the following fees or credit any overpayment to Deposit Account No. 16-0235. One additional copy of this sheet is enclosed.

- ☒ The above calculated filing fee \$-0-
- ☐ Any additional fees required under 37 C.F.R. § 1.16.
- ☒ Any additional fees required under 37 C.F.R. §1.17.
- ☒ If the filing of any paper during the prosecution of this application requires an extension of time in order for the paper to be timely filed, applicant(s) hereby petition(s) for the appropriate extension of time pursuant to 37 C.F.R. §1.136(a).

**CORRESPONDENCE ADDRESS:**

June 14, 1999  
(Date)

By:

GARY D. COLBY, Ph.D., J.D.

Registration No. 40,961

**PANITCH SCHWARZE JACOBS & NADEL, P.C.**

One Commerce Square

2005 Market Street - 22nd Floor

Philadelphia, PA 19103-7086

Telephone: 215-567-2020

**Direct Dial: (215) 965-1285**

Facsimile: 215-567-2991

E-Mail: gdc@psjn.com

☒ Customer Number or Bar Code Label: **000570**

KDL/GDC/moh  
Enclosures

06/14/99  
Jc573 U.S. PTO

5  
NOVEL GENES ENCODING PROTEINS HAVING  
DIAGNOSTIC, PREVENTIVE, THERAPEUTIC, AND OTHER USES

Background of the Invention

10 The molecular bases underlying many human and animal physiological  
states (e.g. diseased and homeostatic states of various tissues) remain unknown.  
Nonetheless, it is well understood that these states result from interactions among the  
proteins and nucleic acids present in the cells of the relevant tissues. In the past, the  
complexity of biological systems overwhelmed the ability of practitioners to understand  
the molecular interactions giving rise to normal and abnormal physiological states.  
15 More recently, though, the techniques of molecular biology, transgenic and null mutant  
animal production, computational biology, pharmacogenomics, and the like have  
enabled practitioners to discern the role and importance of individual genes and  
proteins in particular physiological states.

20 Knowledge of the sequences and other properties of genes (particularly  
including the portions of genes encoding proteins) and the proteins encoded thereby  
enables the practitioner to design and screen agents which will affect, prospectively or  
retrospectively, the physiological state of an animal tissue in a favorable way. Such  
knowledge also enables the practitioner, by detecting the levels of gene expression and  
protein production, to diagnose the current physiological state of a tissue or animal and  
25 to predict such physiological states in the future. This knowledge furthermore enables  
the practitioner to identify and design molecules which bind with the polynucleotides  
and proteins, *in vitro*, *in vivo*, or both.

30 The present invention provides sequence information for  
polynucleotides derived from human and murine genes and for proteins encoded  
thereby, and thus enables the practitioner to assess, predict, and affect the physiological  
state of various human and murine tissues.

### Summary of the Invention

The present invention is based, at least in part, on the discovery of a variety of human and murine cDNA molecules which encode proteins which are herein designated TANGO 202, TANGO 234, TANGO 265, TANGO 273, TANGO 286, TANGO 294, and INTERCEPT 296. These seven proteins, fragments thereof, derivatives thereof, and variants thereof are collectively referred to herein as the polypeptides of the invention or the proteins of the invention. Nucleic acid molecules encoding polypeptides of the invention are collectively referred to as nucleic acids of the invention.

The nucleic acids and polypeptides of the present invention are useful as modulating agents in regulating a variety of cellular processes. Accordingly, in one aspect, the present invention provides isolated nucleic acid molecules encoding a polypeptide of the invention or a biologically active portion thereof. The present invention also provides nucleic acid molecules which are suitable as primers or hybridization probes for the detection of nucleic acids encoding a polypeptide of the invention.

The invention also features nucleic acid molecules which are at least 40% (or 50%, 60%, 70%, 80%, 90%, 95%, or 98%) identical to the nucleotide sequence of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA clone deposited with ATCC as one of Accession numbers 207219, 207184, 207228, 207185, 207220, and 207221 ("a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221"), or a complement thereof.

The invention features nucleic acid molecules which include a fragment of at least 15 (25, 40, 60, 80, 100, 150, 200, 250, 300, 350, 400, 450, 550, 650, 700, 800, 900, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000, 3500, 4000, 4500, or 4928) consecutive nucleotide residues of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof.

The invention also features nucleic acid molecules which include a nucleotide sequence encoding a protein having an amino acid sequence that is at least 50% (or 60%, 70%, 80%, 90%, 95%, or 98%) identical to the amino acid sequence of any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof.

In preferred embodiments, the nucleic acid molecules have the nucleotide sequence of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221.

Also within the invention are nucleic acid molecules which encode a fragment of a polypeptide having the amino acid sequence of any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, the fragment including at least 8 (10, 15, 20, 25, 30, 40, 50, 75, 100, 125, 150, or 200) consecutive amino acids of any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221.

The invention includes nucleic acid molecules which encode a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, wherein the nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule having a nucleic acid sequence encoding any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof.

Also within the invention are isolated polypeptides or proteins having an amino acid sequence that is at least about 50%, preferably 60%, 75%, 90%, 95%, or

98% identical to the amino acid sequence of any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, and 74.

Also within the invention are isolated polypeptides or proteins which are encoded by a nucleic acid molecule having a nucleotide sequence that is at least about 40%, preferably 50%, 75%, 85%, or 95% identical the nucleic acid sequence encoding any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, and isolated polypeptides or proteins which are encoded by a nucleic acid molecule consisting of the nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule having the nucleotide sequence of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, and 73.

Also within the invention are polypeptides which are naturally occurring allelic variants of a polypeptide that includes the amino acid sequence of any of SEQ ID NOs: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes under stringent conditions to a nucleic acid molecule having the nucleotide sequence of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 317, 319, or a complement thereof.

The invention also features nucleic acid molecules that hybridize under stringent conditions to a nucleic acid molecule having the nucleotide sequence of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221 or a complement thereof. In other embodiments, the nucleic acid molecules are at least 15 (25, 40, 60, 80, 100, 150, 200, 250, 300, 350, 400, 450, 550, 650, 700, 800, 900, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000, 3500, 4000, 4500, or 4928) nucleotides in length and hybridize under stringent conditions to a nucleic acid molecule having the nucleotide sequence of any of SEQ ID NOs: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof. In some embodiments,

the isolated nucleic acid molecules encode a cytoplasmic, transmembrane, extracellular, or other domain of a polypeptide of the invention. In other embodiments, the invention provides an isolated nucleic acid molecule which is antisense to the coding strand of a nucleic acid of the invention.

Another aspect of the invention provides vectors, e.g., recombinant expression vectors, comprising a nucleic acid molecule of the invention. In another embodiment, the invention provides isolated host cells, e.g., mammalian and non-mammalian cells, containing such a vector or a nucleic acid of the invention. The invention also provides methods for producing a polypeptide of the invention by culturing, in a suitable medium, a host cell of the invention containing a recombinant expression vector encoding a polypeptide of the invention such that the polypeptide of the invention is produced.

Another aspect of this invention features isolated or recombinant proteins and polypeptides of the invention. Preferred proteins and polypeptides possess at least one biological activity possessed by the corresponding naturally-occurring human polypeptide. An activity, a biological activity, and a functional activity of a polypeptide of the invention refers to an activity exerted by a protein or polypeptide of the invention on a responsive cell as determined *in vivo*, or *in vitro*, according to standard techniques. Such activities can be a direct activity, such as an association with or an enzymatic activity on a second protein or an indirect activity, such as a cellular processes mediated by interaction of the protein with a second protein.

By way of example, TANGO 202 exhibits the ability to affect growth, proliferation, survival, differentiation, and activity of human hematopoietic cells (e.g. bone marrow stromal cells) and fetal cells. TANGO 202 modulates cellular binding to one or more mediators, modulates proteolytic activity *in vivo*, modulates developmental processes, and modulates cell growth, proliferation, survival, differentiation, and activity. Thus, TANGO 202 can be used to prevent, diagnose, or treat disorders relating to aberrant cellular protease activity, inappropriate interaction (or non-interaction) of cells with mediators, inappropriate development, and blood and hematopoietic cell-related disorders. Exemplary disorders for which TANGO 202 is

useful include immune disorders, infectious diseases, auto-immune disorders, vascular and cardiovascular disorders, disorders related to mal-expression of growth factors, cancers, hematological disorders, various cancers, birth defects, developmental defects, and the like.

Further by way of example, TANGO 234 exhibits the ability to affect growth, proliferation, survival, differentiation, and activity of human lung, hematopoietic, and fetal cells and of (e.g. bacterial or fungal) cells and viruses which infect humans. TANGO 234 modulates growth, proliferation, survival, differentiation, and activity of gamma delta T cells, for example. Furthermore, TANGO 234 modulates cholesterol deposition on human arterial walls, and is involved in uptake and metabolism of low density lipoprotein and regulation of serum cholesterol levels. Thus, TANGO 234 can be used to affect development and persistence of atherogenesis and arteriosclerosis, as well as other vascular and cardiovascular disorders. Other exemplary disorders for which TANGO 234 is useful include immune development disorders and disorders involving generation and persistence of an immune response to bacterial, fungal, and viral infections.

Still further by way of example, TANGO 265 modulates growth and regeneration of neuronal and epithelial tissues, and guides neuronal axon development. TANGO 265 is a transmembrane protein which mediates cellular interaction with cells, molecules and structures (e.g. extracellular matrix) in the extracellular environment. TANGO 265 is therefore involved in growth, organization, and adhesion of tissues and the cells which constitute those tissues. Furthermore, TANGO 265 modulates growth, proliferation, survival, differentiation, and activity of neuronal cells and immune system cells. Thus, TANGO 265 can be used, for example, to prevent, diagnose, or treat disorders characterized by aberrant organization or development of a tissue or organ, for guiding neural axon development, for modulating differentiation of cells of the immune system, for modulating cytokine production by cells of the immune system, for modulating reactivity of cells of the immune system toward cytokines, for

modulating initiation and persistence of an inflammatory response, and for modulating proliferation of epithelial cells.

Yet further by way of example, TANGO 273 protein mediates one or more physiological responses of cells to bacterial infection, e.g., by mediating one or more of detection of bacteria in a tissue in which it is expressed, movement of cells with relation to sites of bacterial infection, production of biological molecules which inhibit bacterial infection, and production of biological molecules which alleviate cellular or other physiological damage wrought by bacterial infection. TANGO 273, a transmembrane protein, is also involved in transmembrane signal transduction, and therefore mediates transmission of signals between the extracellular and intracellular environments of cells. TANGO 273 mediates regulation of cell growth and proliferation, endocytosis, activation of respiratory burst, and other physiological processes triggered by transmission of a signal via a protein with which TANGO 273 interacts. The compositions and methods of the invention can therefore be used to prevent, diagnose, and treat disorders involving one or more physiological activities mediated by TANGO 273 protein. Such disorders include, for example, various bone-related disorders such as metabolic, homeostatic, and developmental bone disorders (e.g. osteoporosis, various cancers, skeletal development disorders, bone fragility and the like), disorders caused by or related to bacterial infection, and disorders characterized by aberrant transmembrane signal transduction by TANGO 273.

As an additional example, TANGO 286 protein is involved in lipid-binding physiological processes such as lipid transport, metabolism, serum lipid particle regulation, host antimicrobial defensive mechanisms, and the like. Thus, the compositions and methods of the invention can therefore be used to prevent, diagnose, and treat disorders involving one or more physiological activities mediated by TANGO 286 protein. Such disorders include, for example, lipid transport disorders, lipid metabolism disorders, obesity, disorders of serum lipid particle regulation, disorders involving insufficient or inappropriate host antimicrobial defensive mechanisms, vasculitis, bronchiectasis, LPS-related disorders such as shock, disseminated

intravascular coagulation, anemia, thrombocytopenia, adult respiratory distress syndrome, renal failure, liver disease, and disorders associated with Gram negative bacterial infections, such as bacteremia, endotoxemia, sepsis, and the like.

Further by way of example, TANGO 294 protein is involved in facilitating absorption and metabolism of fat. Thus, the compositions and methods of the invention can therefore be used to prevent, diagnose, and treat disorders involving one or more physiological activities mediated by TANGO 294 protein. Such disorders include, for example, inadequate expression of gastric/pancreatic lipase, cystic fibrosis, exocrine pancreatic insufficiency, medical treatments which alter fat absorption, obesity, and the like.

As another example, INTERCEPT 296 protein is involved in physiological processes related to disorders of the human lung and esophagus. Thus, the compositions and methods of the invention can be used to prevent, diagnose, and treat these disorders. Such disorders include, for example, various cancers, bronchitis, cystic fibrosis, respiratory infections (e.g. influenza, bronchiolitis, pneumonia, and tuberculosis), asthma, emphysema, chronic bronchitis, bronchiectasis, pulmonary edema, pleural effusion, pulmonary embolus, adult and infant respiratory distress syndromes, heartburn, and gastric reflux esophageal disease.

In one embodiment, a polypeptide of the invention has an amino acid sequence sufficiently identical to an identified domain of a polypeptide of the invention. As used herein, the term "sufficiently identical" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (e.g., with a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences have a common domain and/or common functional activity. For example, amino acid or nucleotide sequences which contain a common domain having about 65% identity, preferably 75% identity, more preferably 85%, 95%, or 98% identity are defined herein as sufficiently identical.

In one embodiment, the isolated polypeptide of the invention lacks both a transmembrane and a cytoplasmic domain. In another embodiment, the polypeptide lacks both a transmembrane domain and a cytoplasmic domain and is soluble under physiological conditions.

The polypeptides of the present invention, or biologically active portions thereof, can be operably linked to a heterologous amino acid sequence to form fusion proteins. The invention further features antibody substances that specifically bind a polypeptide of the invention such as monoclonal or polyclonal antibodies, antibody fragments, single-chain antibodies, and the like. In addition, the polypeptides of the invention or biologically active portions thereof can be incorporated into pharmaceutical compositions, which optionally include pharmaceutically acceptable carriers. These antibody substances can be made, for example, by providing the polypeptide of the invention to an immunocompetent vertebrate and thereafter harvesting blood or serum from the vertebrate.

In another aspect, the present invention provides methods for detecting the presence of the activity or expression of a polypeptide of the invention in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of activity such that the presence of activity is detected in the biological sample.

In another aspect, the invention provides methods for modulating activity of a polypeptide of the invention comprising contacting a cell with an agent that modulates (inhibits or enhances) the activity or expression of a polypeptide of the invention such that activity or expression in the cell is modulated. In one embodiment, the agent is an antibody that specifically binds to a polypeptide of the invention.

In another embodiment, the agent modulates expression of a polypeptide of the invention by modulating transcription, splicing, or translation of an mRNA encoding a polypeptide of the invention. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense with respect to the coding strand of an mRNA encoding a polypeptide of the invention.

5 The present invention also provides methods to treat a subject having a disorder characterized by aberrant activity of a polypeptide of the invention or aberrant expression of a nucleic acid of the invention by administering an agent which is a modulator of the activity of a polypeptide of the invention or a modulator of the expression of a nucleic acid of the invention to the subject. In one embodiment, the modulator is a protein of the invention. In another embodiment, the modulator is a nucleic acid of the invention. In other embodiments, the modulator is a peptide, peptidomimetic, or other small molecule.

10 The present invention also provides diagnostic assays for identifying the presence or absence of a genetic lesion or mutation characterized by at least one of: (i) aberrant modification or mutation of a gene encoding a polypeptide of the invention, (ii) mis-regulation of a gene encoding a polypeptide of the invention, and (iii) aberrant post-translational modification of a polypeptide of the invention wherein a wild-type form of the gene encodes a polypeptide having the activity of the polypeptide of the invention.

15 In another aspect, the invention provides a method for identifying a compound that binds to or modulates the activity of a polypeptide of the invention. In general, such methods entail measuring a biological activity of the polypeptide in the presence and absence of a test compound and identifying those compounds which alter the activity of the polypeptide.

20 The invention also features methods for identifying a compound which modulates the expression of a polypeptide or nucleic acid of the invention by measuring the expression of the polypeptide or nucleic acid in the presence and absence of the compound.

25 Other features and advantages of the invention will be apparent from the following detailed description and claims.

#### Brief Description of the Drawings

30 *Figure 1* comprises Figures 1A through 1H. The nucleotide sequence (SEQ ID NO: 1) of a cDNA encoding the human TANGO 202 protein described herein

is listed in Figures 1A and 1B. The open reading frame (ORF; residues 34 to 1458; SEQ ID NO: 2) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 3) of human TANGO 202 is listed. The nucleotide sequence (SEQ ID NO: 67) of a cDNA encoding the murine TANGO 202 protein described herein is listed in Figures 1C, 1D, and 1E. The ORF (residues 81 to 1490; SEQ ID NO: 68) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 69) of murine TANGO 202 is listed. An alignment of the amino acid sequences of human ("Hum."; SEQ ID NO: 3) and murine ("Mur."; SEQ ID NO: 69) TANGO 202 protein is shown in Figure 1F, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".". Figure 1G is a hydrophilicity plot of human TANGO 202 protein, in which the locations of cysteine residues ("Cys") and potential N-glycosylation sites ("Ngly") are indicated by vertical bars and the predicted extracellular ("out"), intracellular ("ins"), or transmembrane ("TM") locations of the protein backbone is indicated by a horizontal bar. Figure 1H is a hydrophilicity plot of murine TANGO 202 protein.

*Figure 2* comprises Figures 2A through 2Q. The nucleotide sequence (SEQ ID NO: 9) of a cDNA encoding the human TANGO 234 protein described herein is listed in Figures 2A through 2E. The ORF (residues 28 to 4386; SEQ ID NO: 10) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 11) of human TANGO 234 is listed. Figure 2F is a hydrophilicity plot of human TANGO 234 protein. An alignment of the amino acid sequences of human TANGO 234 ("Hum"; SEQ ID NO: 11) and bovine WC1 ("WC1"; SEQ ID NO: 78) proteins is shown in Figures 2G through 2I, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".". An alignment of the nucleotide sequences of an ORF encoding human TANGO 234 ("Hum"; SEQ ID NO: 10) and an ORF encoding bovine WC1 ("WC1"; SEQ ID NO: 79) proteins is shown in Figures 2J through 2Q, wherein identical nucleotide residues are indicated by ":".

*Figure 3* comprises Figures 3A through 3I. The nucleotide sequence (SEQ ID NO: 17) of a cDNA encoding the human TANGO 265 protein described

herein is listed in Figures 3A, 3B, and 3C. The ORF (residues 32 to 2314; SEQ ID NO: 18) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 19) of human TANGO 265 is listed. An alignment of the amino acid sequences of human TANGO 265 protein ("Hum."; SEQ ID NO: 19) and murine semaphorin B protein ("Mur."; SEQ ID NO: 70; GenBank Accession No. X85991) is shown in Figure 3D, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".". In Figures 3E, 3F, 3G, and 3H, an alignment of the nucleotide sequences of the cDNA encoding human TANGO 265 protein ("Hum."; SEQ ID NO: 17) and the nucleotide sequences of the cDNA encoding murine semaphorin B protein ("Mur."; SEQ ID NO: 71; GenBank Accession No. X85991) is shown. Figure 3I is a hydrophilicity plot of TANGO 265 protein.

*Figure 4* comprises Figures 4A through 4G. The nucleotide sequence (SEQ ID NO: 25) of a cDNA encoding the human TANGO 273 protein described herein is listed in Figures 4A and 4B. The ORF (residues 135 to 650; SEQ ID NO: 26) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 27) of human TANGO 273 is listed. The nucleotide sequence (SEQ ID NO: 72) of a cDNA encoding the murine TANGO 273 protein described herein is listed in Figures 4C, and 4D. The ORF (residues 137 to 652; SEQ ID NO: 73) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 74) of murine TANGO 273 is listed. An alignment of the amino acid sequences of human ("Hum."; SEQ ID NO: 27) and murine ("Mur."; SEQ ID NO: 74) TANGO 273 protein is shown in Figure 4E, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".". Figure 4F is a hydrophilicity plot of human TANGO 273 protein, and Figure 4G is a hydrophilicity plot of murine TANGO 273 protein.

*Figure 5* comprises Figures 5A through 5E. The nucleotide sequence (SEQ ID NO: 33) of a cDNA encoding the human TANGO 286 protein described herein is listed in Figures 5A and 5B. The ORF (residues 133 to 1497; SEQ ID NO: 34) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 35) of human TANGO 286 is listed. Figure 5C is a

hydrophilicity plot of TANGO 286 protein. An alignment of the amino acid sequences of human TANGO 286 ("286"; SEQ ID NO: 35) and BPI protein ("BPI"; SEQ ID NO: 38) protein is shown in Figure 5D, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".". An alignment of the amino acid sequences of human TANGO 286 ("286"; SEQ ID NO: 35) and RENP protein ("RENP"; SEQ ID NO: 39) is shown in Figure 5E, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".".

*Figure 6* comprises Figures 6A through 6E. The nucleotide sequence (SEQ ID NO: 45) of a cDNA encoding the human TANGO 294 protein described herein is listed in Figures 6A and 6B. The ORF (residues 126 to 1394; SEQ ID NO: 46) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 47) of human TANGO 294 is listed. An alignment of the amino acid sequences of human TANGO 294 protein ("294"; SEQ ID NO: 47) and a known human lipase protein ("HLP"; SEQ ID NO: 75; GenBank Accession No. NP\_004181) is shown in Figure 6C, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".". Figure 6D is a hydrophilicity plot of TANGO 294 protein. An alignment of the amino acid sequences of human TANGO 294 protein ("294"; SEQ ID NO: 47) and a known human lysosomal acid lipase protein ("LAL"; SEQ ID NO: 41) is shown in Figure 6E, wherein identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".".

*Figure 7* comprises Figures 7A through 7D. The nucleotide sequence (SEQ ID NO: 53) of a cDNA encoding the human INTERCEPT 296 protein described herein is listed in Figures 7A and 7B. The ORF (residues 70 to 1098; SEQ ID NO: 54) of the cDNA is indicated by nucleotide triplets, above which the amino acid sequence (SEQ ID NO: 55) of human INTERCEPT 296 protein is listed. Figure 7C is a hydrophilicity plot of INTERCEPT 296 protein. An alignment of the amino acid sequences of human INTERCEPT 296 protein ("296"; SEQ ID NO: 55) and *C. elegans* C06E1.3 related protein ("CRP"; SEQ ID NO: 40) is shown in Figure 7D, wherein

identical amino acid residues are indicated by ":" and similar amino acid residues are indicated by ".".

### Detailed Description of the Invention

The present invention is based, at least in part, on the discovery of a variety of human and murine cDNA molecules which encode proteins which are herein designated TANGO 202, TANGO 234, TANGO 265, TANGO 273, TANGO 286, TANGO 294, and INTERCEPT 296. These proteins exhibit a variety of physiological activities, and are included in a single application for the sake of convenience. It is understood that the allowability or non-allowability of claims directed to one of these proteins has no bearing on the allowability of claims directed to the others. The characteristics of each of these proteins and the cDNAs encoding them are now described separately.

#### TANGO 202

A cDNA encoding at least a portion of human TANGO 202 protein was isolated from a human fetal skin cDNA library. The corresponding murine cDNA was isolated from a bone marrow stromal cell cDNA library. The human TANGO 202 protein is predicted by structural analysis to be a type I membrane protein, although it can exist in a secreted form as well. The murine TANGO 202 protein is predicted by structural analysis to be a secreted protein.

The full length of the cDNA encoding human TANGO 202 protein (Figure 1; SEQ ID NO: 1) is 1656 nucleotide residues. The open reading frame (ORF) of this cDNA, nucleotide residues 34 to 1458 of SEQ ID NO: 1 (i.e. SEQ ID NO: 2), encodes a 475-amino acid transmembrane protein (Figure 1; SEQ ID NO: 3).

The invention thus includes purified human TANGO 202 protein, both in the form of the immature 475 amino acid residue protein (SEQ ID NO: 3) and in the form of the mature 456 amino acid residue protein (SEQ ID NO: 5). The invention also includes purified murine TANGO 202 protein, both in the form of the immature 470 amino acid residue protein (SEQ ID NO: 67) and in the form of the mature 451 amino acid residue protein (SEQ ID NO: 43). Mature human or murine TANGO 202

proteins can be synthesized without the signal sequence polypeptide at the amino terminus thereof, or they can be synthesized by generating immature TANGO 202 protein and cleaving the signal sequence therefrom.

In addition to full length mature and immature human and murine TANGO 202 proteins, the invention includes fragments, derivatives, and variants of these TANGO 202 proteins, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA molecule having the nucleotide sequence listed in SEQ ID NO: 1 or some portion thereof or SEQ ID NO: 67 or some portion thereof, such as the portion which encodes mature human or murine TANGO 202 protein, immature human or murine TANGO 202 protein, or a domain of human or murine TANGO 202 protein. These nucleic acids are collectively referred to as nucleic acids of the invention.

TANGO 202 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features. As used herein, the term "family" is intended to mean two or more proteins or nucleic acid molecules having a common or similar domain structure and having sufficient amino acid or nucleotide sequence identity as defined herein. Family members can be from either the same or different species (e.g. human and mouse, as described herein). For example, a family can comprise two or more proteins of human origin, or can comprise one or more proteins of human origin and one or more of non-human origin.

A common domain present in TANGO 202 proteins is a signal sequence. As used herein, a signal sequence includes a peptide of at least about 10 amino acid residues in length which occurs at the amino terminus of membrane-bound and secreted proteins and which contains at least about 45% hydrophobic amino acid residues such as alanine, leucine, isoleucine, phenylalanine, proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least

about 10 to 35 amino acid residues, preferably about 10 to 20 amino acid residues, and has at least about 35-60%, more preferably 40-50%, and more preferably at least about 45% hydrophobic residues. A signal sequence serves to direct a protein containing such a sequence to a lipid bilayer. Thus, in one embodiment, a TANGO 202 protein contains a signal sequence corresponding to amino acid residues 1 to 19 of SEQ ID NO: 3 (SEQ ID NO: 4) or to amino acid residues 1 to 19 of SEQ ID NO: 69 (SEQ ID NO: 42). The signal sequence is cleaved during processing of the mature protein.

TANGO 202 proteins can also include an extracellular domain. As used herein, an "extracellular domain" refers to a portion of a protein which is localized to the non-cytoplasmic side of a lipid bilayer of a cell when a nucleic acid encoding the protein is expressed in the cell. The human TANGO 202 protein extracellular domain is located from about amino acid residue 20 to about amino acid residue 392 of SEQ ID NO: 3 in the non-secreted form, and from about amino acid residue 20 to amino acid residue 475 of SEQ ID NO: 3 (i.e. the entire mature human protein). The murine TANGO 202 protein extracellular domain is located from about amino acid residue 20 to amino acid residue 470 of SEQ ID NO: 69 (i.e. the entire mature murine protein).

TANGO 202 proteins of the invention can also include a transmembrane domain. As used herein, a "transmembrane domain" refers to an amino acid sequence having at least about 20 to 25 amino acid residues in length and which contains at least about 65-70% hydrophobic amino acid residues such as alanine, leucine, phenylalanine, protein, tyrosine, tryptophan, or valine. In a preferred embodiment, a transmembrane domain contains at least about 15 to 30 amino acid residues, preferably about 20-25 amino acid residues, and has at least about 60-80%, more preferably 65-75%, and more preferably at least about 70% hydrophobic residues. Thus, in one embodiment, a TANGO 202 protein of the invention contains a transmembrane domain corresponding to about amino acid residues 393 to 415 of SEQ ID NO: 3 (SEQ ID NO: 7).

In addition, TANGO 202 proteins of the invention can include a cytoplasmic domain, particularly including a carboxyl-terminal cytoplasmic domain. As used herein, a "cytoplasmic domain" refers to a portion of a protein which is localized to the cytoplasmic side of a lipid bilayer of a cell when a nucleic acid

encoding the protein is expressed in the cell. The cytoplasmic domain is located from about amino acid residue 416 to amino acid residue 475 of SEQ ID NO: 3 (SEQ ID NO: 8) in the non-secreted form of human TANGO 202 protein.

TANGO 202 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Tables I (for human TANGO 202) and II (for murine TANGO 202), as predicted by computerized sequence analysis of TANGO 202 proteins using amino acid sequence comparison software (comparing the amino acid sequence of TANGO 202 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}).

Table I

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 3	Amino Acid Sequence
N-glycosylation site	47 to 50 61 to 64 219 to 222 295 to 298 335 to 338 347 to 350	NWTA NETF NYS NVSL NQTV NLSV
Protein kinase C phosphorylation site	70 to 72 137 to 139 141 to 143 155 to 157 238 to 240 245 to 247 277 to 279 307 to 309 355 to 357 387 to 389 418 to 420 421 to 423	TLK TSK SNK SQR TGR TIR THR SDR SSK SHR TFK SHR
Casein kinase II phosphorylation site	337 to 340 438 to 441 464 to 467	TVAE TSGE SQQD

Table I (Cont'd)

N-myristoylation site	53 to 58 120 to 125 136 to 141 162 to 167 214 to 219	GGKPCL GNLGCY GTSKTS GMESGY GACGGN
Kringle domain signature	85 to 90	YCRNPD
Kringle Domain	34 to 116	See Fig. 1
CUB domain	216 to 320	See Fig. 1

Table II

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 69	Amino Acid Sequence
N-glycosylation site	59 to 62 217 to 220 255 to 258 293 to 296 333 to 336 345 to 348	NETF NYSA NFTL NVSL NQTL NLSV
cAMP- or cGMP-dependent protein kinase phosphorylation site	455 to 458	RRSS
Protein kinase C phosphorylation site	68 to 70 135 to 137 139 to 141 153 to 155 236 to 238 243 to 245 275 to 277 283 to 285 305 to 307 353 to 355 408 to 410 453 to 455 457 to 459	TLK TSK SNK SQR TGR TIR THR SGR SDR SSK SQR SLR SSR

Table II (Cont'd)

5	Casein kinase II phosphorylation site	28 to 31	SGPE
		257 to 260	TLFD
		321 to 324	TKEE
		335 to 338	TLAE
		384 to 387	TATE
10	N-myristoylation site	51 TO 56	GGKPCL
		118 TO 123	GNLGCY
		134 TO 139	GTSKTS
		160 TO 165	GMESGY
		212 TO 217	GACGGN
		391 TO 396	GLCTAW
		429 TO 434	GTVVSL
15	Kringle domain signature	83 to 88	YCRNPD
	Kringle Domain	32 to 114	See Fig. 1
	CUB domain	214 to 318	See Fig. 1

As used herein, the term "post-translational modification site" refers to a protein domain that includes about 3 to 10 amino acid residues, more preferably about 3 to 6 amino acid residues wherein the domain has an amino acid sequence which comprises a consensus sequence which is recognized and modified by a protein-modifying enzyme. Exemplary protein-modifying enzymes include amino acid glycosylases, cAMP- and cGMP-dependent protein kinases, protein kinase C, casein kinase II, myristoylases, and prenyl transferases. In various embodiments, the protein of the invention has at least 1, 2, 4, 6, 10, 15, or 20 or more of the post-translational modification sites described herein in Tables I and II.

Exemplary additional domains present in human and murine TANGO 202 protein include Kringle domains and CUB domains. In one embodiment, the protein of the invention has at least one domain that is at least 55%, preferably at least about 65%, more preferably at least about 75%, yet more preferably at least about 85%, and most preferably at least about 95% identical to one of the domains described herein in Tables I and II. Preferably, the protein of the invention has at least one Kringle domain and one CUB domain.

A Kringle domain has a characteristic profile that has been described in the art (Castellino and Beals (1987) *J. Mol. Evol.* 26:358-369; Patthy (1985) *Cell* 41:657-663; Ikeo et al. (1991) *FEBS Lett.* 287:146-148). Many, but not all, Kringle domains comprise a conserved hexapeptide signature sequence, namely

(F or Y) - C - R - N - P - (D or N or R).

The cysteine residue is involved in a disulfide bond.

Kringle domains are triple-looped, disulfide cross-linked domains found in a varying number of copies in, for example, some serine proteases and plasma proteins. Kringle domains have a role in binding mediators (e.g. membranes, other proteins, or phospholipids) and in regulation of proteolytic activity. Kringle domains have been identified in the following proteins, for example: apolipoprotein A, blood coagulation factor XII (Hageman factor), hepatocyte growth factor (HGF), HGF-like protein (Frieznier Degen et al., (1991) *Biochemistry* 30:9781-9791), HGF activator (Miyazawa et al., (1993) *J. Biol. Chem.* 268:10024-10028), plasminogen, thrombin, tissue plasminogen activator, urokinase-type plasminogen activator, and four influenza neuraminidases. The presence of a Kringle domain in each of human and murine TANGO 202 protein indicates that TANGO 202 is involved in one or more physiological processes in which these other Kringle domain-containing proteins are involved, has biological activity in common with one or more of these other Kringle domain-containing proteins, or both.

CUB domains are extracellular domains of about 110 amino acid residues which occur in functionally diverse, mostly developmentally regulated proteins (Bork and Beckmann (1993) *J. Mol. Biol.* 231:539-545; Bork (1991) *FEBS Lett.* 282:9-12). Many CUB domains contain four conserved cysteine residues, although some, like that of TANGO 202, contain only two of the conserved cysteine residues. The structure of the CUB domain has been predicted to assume a beta-barrel configuration, similar to that of immunoglobulins. Other proteins which have been found to comprise one or more CUB domains include, for example, mammalian complement sub-components C1s and C1r, hamster serine protease Casp, mammalian complement activating component of Ra-reactive factor, vertebrate enteropeptidase,

vertebrate bone morphogenic protein 1, sea urchin blastula proteins BP10 and SpAN, *Caenorhabditis elegans* hypothetical proteins F42A10.8 and R151.5, neuropilin (A5 antigen), sea urchin fibropellins I and III, mammalian hyaluronate-binding protein TSG-6 (PS4), mammalian spermadhesins, and *Xenopus* embryonic protein UVS.2. The presence of a CUB domain in each of human and murine TANGO 202 protein indicates that TANGO 202 is involved in one or more physiological processes in which these other CUB domain-containing proteins are involved, has biological activity in common with one or more of these other CUB domain-containing proteins, or both.

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that human TANGO 202 protein includes a 19 amino acid signal peptide (amino acid residues 1 to 19 of SEQ ID NO: 3; SEQ ID NO: 4) preceding the mature TANGO 202 protein (amino acid residues 20 to 475 of SEQ ID NO: 3; SEQ ID NO: 5). Human TANGO 202 protein includes an extracellular domain (amino acid residues 20 to 392 of SEQ ID NO: 3; SEQ ID NO: 6); a transmembrane domain (amino acid residues 393 to 415 of SEQ ID NO: 3; SEQ ID NO: 7); and a cytoplasmic domain (amino acid residues 416 to 475 of SEQ ID NO: 3; SEQ ID NO: 8). The murine homolog of TANGO 202 protein is predicted to be a secreted protein. Thus, it is recognized that human TANGO 202 can also exist in the form of a secreted protein, likely being translated from an alternatively spliced TANGO 202 mRNA. In a variant form of the protein, an extracellular portion of TANGO 202 protein (e.g. amino acid residues 20 to 392 of SEQ ID NO: 3) can be cleaved from the mature protein to generate a soluble fragment of TANGO 202.

Figure 1G depicts a hydrophilicity plot of human TANGO 202 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 19 of SEQ ID NO: 3 is the signal sequence of human TANGO 202 (SEQ ID NO: 4). The hydrophobic region which corresponds to amino acid residues 393 to 415 of SEQ ID NO: 3 is the transmembrane domain of human TANGO 202 (SEQ ID NO: 7). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein,

and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human TANGO 202 protein from about amino acid residue 61 to about amino acid residue 95 appears to be located at or near the surface of the protein, while the region from about amino acid residue 395 to about amino acid residue 420 appears not to be located at or near the surface.

The predicted molecular weight of human TANGO 202 protein without modification and prior to cleavage of the signal sequence is about 51.9 kilodaltons. The predicted molecular weight of the mature human TANGO 202 protein without modification and after cleavage of the signal sequence is about 50.1 kilodaltons.

The full length of the cDNA encoding murine TANGO 202 protein (Figure 1; SEQ ID NO: 67) is 4928 nucleotide residues. The ORF of this cDNA, nucleotide residues 81 to 1490 of SEQ ID NO: 67 (i.e. SEQ ID NO: 68), encodes a 470-amino acid secreted protein (Figure 1; SEQ ID NO: 69).

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that murine TANGO 202 protein includes a 19 amino acid signal peptide (amino acid residues 1 to 19 of SEQ ID NO: 69; SEQ ID NO: 42) preceding the mature TANGO 202 protein (amino acid residues 20 to 470 of SEQ ID NO: 69; SEQ ID NO: 43). Murine TANGO 202 protein is a secreted protein.

Figure 1H depicts a hydrophilicity plot of murine TANGO 202 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 19 of SEQ ID NO: 69 is the signal sequence of murine TANGO 202 (SEQ ID NO: 42). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of murine TANGO 202 protein from about amino acid residue 61 to about amino acid residue 95 appears to be located at or near the surface of the protein, while the region from about amino acid residue 295 to about amino acid residue 305 appears not to be located at or near the surface

The predicted molecular weight of murine TANGO 202 protein without modification and prior to cleavage of the signal sequence is about 51.5 kilodaltons. The predicted molecular weight of the mature murine TANGO 202 protein without modification and after cleavage of the signal sequence is about 49.7 kilodaltons.

Human and murine TANGO 202 proteins exhibit considerable sequence similarity, as indicated herein in Figure 1F. Figure 1F depicts an alignment of human and murine TANGO 202 amino acid sequences (SEQ ID NOs: 3 and 69, respectively). In this alignment (made using the ALIGN software {Myers and Miller (1989) *CABIOS*, ver. 2.0}; pam120.mat scoring matrix; gap penalties -12/-4), the proteins are 76.5% identical. The human and murine ORFs encoding TANGO 202 are 87.4% identical, as assessed using the same software and parameters.

*In situ* hybridization experiments in mouse tissues indicated that mRNA corresponding to the cDNA encoding TANGO 202 is expressed in the tissues listed in Table III, wherein "+" indicates detectable expression and "++" indicates a greater level of expression than "+".

Table III

Animal	Tissue	Relative Level of Expression
Mouse (Adult)	bladder, especially in transitional epithelium	++
	renal glomeruli	+
	brain	+
	heart	+
	liver	+
	spleen	+
	placenta	+
Mouse (Embryo)	ubiquitous	+

Biological function of TANGO 202 proteins, nucleic acids encoding them, and modulators of these molecules

TANGO 202 proteins are involved in disorders which affect both tissues in which they are normally expressed and tissues in which they are normally not expressed. Based on the observation that TANGO 202 is expressed in human fetal skin, ubiquitously in fetal mouse tissues, in adult murine bone marrow stromal cells, and in cells of adult murine bladder, renal glomeruli, brain, heart, liver, spleen and placenta, TANGO 202 protein is involved in one or more biological processes which occur in these tissues. In particular, TANGO 202 is involved in modulating growth, proliferation, survival, differentiation, and activity of cells of these tissues including, but not limited to, hematopoietic and fetal cells. Thus, TANGO 202 has a role in disorders which affect these cells and their growth, proliferation, survival, differentiation, and activity. Ubiquitous expression of TANGO 202 in fetal murine tissues, contrasted with limited expression in adult murine tissues further indicates that TANGO 202 is involved in disorders in which it is inappropriately expressed (e.g. disorders in which TANGO 202 is expressed in adult murine tissues other than bone marrow stromal cells and disorders in which TANGO 202 is not expressed in one or more developing fetal tissues).

The occurrence of a Kringle domain in both the murine and human TANGO 202 proteins indicates that this protein is involved in modulating cellular binding to one or more mediators (e.g. proteins, phospholipids, intracellular organelles, or other cells), in modulating proteolytic activity, or both. The occurrence of a Kringle domain in other proteins (e.g. growth factors) indicates activities that these proteins share with TANGO 202 protein (e.g. modulating cell dissociation and migration into and through extracellular matrices). The occurrence of Kringle domains in numerous plasma proteins, particularly coupled with the observation that TANGO 202 is expressed in adult murine bone marrow stromal cells, indicates a role for TANGO 202 protein in modulating binding of blood or hematopoietic cells (or both) to one or more mediators. Thus, TANGO 202 is involved in disorders relating to aberrant cellular protease activity, inappropriate interaction or non-interaction of cells with mediators,

and in blood and hematopoietic cell-related disorders. Such disorders include, by way of example and not limitation, immune disorders, infectious diseases, auto-immune disorders, vascular and cardiovascular disorders, disorders related to mal-expression of growth factors, cancers, hematological disorders, and the like.

The occurrence of a CUB domain in both the murine and human TANGO 202 proteins indicates that this protein is involved in biological processes common to other CUB domain-containing proteins, such as developmental processes and binding to mediators. Therefore, TANGO 202 protein has a role in disorders which involve inappropriate developmental processes (e.g. abnormally high proliferation or un-differentiation of a differentiated tissue or abnormally low differentiation or proliferation of a non-developed or non-differentiated tissue) and modulation of cell growth, proliferation, survival, differentiation, and activity. Such disorders include, by way of example and not limitation, various cancers and birth and developmental defects.

Thus, proteins and nucleic acids of the invention which are identical to, similar to, or derived from human and murine TANGO 202 proteins and nucleic acids encoding them are useful for preventing, diagnosing, and treating, among others, vascular and cardiovascular disorders, hematological disorders, disorders related to mal-expression of growth factors, and cancer. Other uses for these proteins and nucleic acids of the invention relate to modulating cell growth (e.g. angiogenesis), proliferation (e.g. cancers), survival (e.g. apoptosis), differentiation (e.g. hematopoiesis), and activity (e.g. ligand-binding capacity). TANGO 202 proteins and nucleic acids encoding them are also useful for modulating cell dissociation and modulating migration of cells in extracellular matrices.

#### TANGO 234

A cDNA encoding at least a portion of human TANGO 234 protein was isolated from a human fetal spleen cDNA library. The human TANGO 234 protein is

predicted by structural analysis to be a transmembrane protein, although it can exist in a secreted form as well.

The full length of the cDNA encoding human TANGO 234 protein (Figure 2; SEQ ID NO: 9) is 4628 nucleotide residues. The ORF of this cDNA, nucleotide residues 28 to 4386 of SEQ ID NO: 9 (i.e. SEQ ID NO: 10), encodes a 1453-amino acid transmembrane protein (Figure 2; SEQ ID NO: 11).

The invention thus includes purified human TANGO 234 protein, both in the form of the immature 1453 amino acid residue protein (SEQ ID NO: 11) and in the form of the mature 1413 amino acid residue protein (SEQ ID NO: 13). Mature human TANGO 234 protein can be synthesized without the signal sequence polypeptide at the amino terminus thereof, or it can be synthesized by generating immature TANGO 234 protein and cleaving the signal sequence therefrom.

In addition to full length mature and immature human TANGO 234 proteins, the invention includes fragments, derivatives, and variants of these TANGO 234 proteins, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA molecule having the nucleotide sequence listed in SEQ ID NO: 9 or some portion thereof, such as the portion which encodes mature TANGO 234 protein, immature TANGO 234 protein, or a domain of TANGO 234 protein. These nucleic acids are collectively referred to as nucleic acids of the invention.

TANGO 234 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features, as indicated by the conservation of amino acid sequence between human TANGO 234 protein and bovine WC1 protein, as shown in Figures 2G through 2I, and the conservation of nucleotide sequence between the ORFs encoding human TANGO 234 protein and bovine WC1 protein, as shown in Figures 2J through 2Q.

A common domain present in TANGO 234 proteins is a signal sequence. As used herein, a signal sequence includes a peptide of at least about 10 amino acid residues in length which occurs at the amino terminus of membrane-bound proteins and which contains at least about 45% hydrophobic amino acid residues such as alanine, leucine, isoleucine, phenylalanine, proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least about 10 to 35 amino acid residues, preferably about 10 to 20 amino acid residues, and has at least about 35-60%, more preferably 40-50%, and more preferably at least about 45% hydrophobic residues. A signal sequence serves to direct a protein containing such a sequence to a lipid bilayer. Thus, in one embodiment, a TANGO 234 protein contains a signal sequence corresponding to amino acid residues 1 to 40 of SEQ ID NO: 11 (SEQ ID NO: 12). The signal sequence is cleaved during processing of the mature protein.

TANGO 234 proteins can include an extracellular domain. The human TANGO 234 protein extracellular domain is located from about amino acid residue 41 to about amino acid residue 1359 of SEQ ID NO: 3. TANGO 234 can alternately exist in a secreted form, such as a mature protein having the amino acid sequence of amino acid residues 41 to 1453 or residues 41 to about 1359 of SEQ ID NO: 11.

In addition, TANGO 234 include a transmembrane domain. In one embodiment, a TANGO 234 protein of the invention contains a transmembrane domain corresponding to about amino acid residues 1360 to 1383 of SEQ ID NO: 11 (SEQ ID NO: 15).

The present invention includes TANGO 234 proteins having a cytoplasmic domain, particularly including proteins having a carboxyl-terminal cytoplasmic domain. The human TANGO 234 cytoplasmic domain is located from about amino acid residue 1384 to amino acid residue 1453 of SEQ ID NO: 11 (SEQ ID NO: 16).

TANGO 234 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Table IV, as predicted by computerized sequence analysis of TANGO 234 proteins using amino acid sequence comparison software (comparing the

amino acid sequence of TANGO 234 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}). In certain embodiments, a protein of the invention has at least 1, 2, 4, 6, 10, 15, or 20 or more of the post-translational modification sites listed in Table IV.

Table IV

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 11	Amino Acid Sequence
N-glycosylation site	42 to 45	NGTD
	78 to 81	NTTA
	120 to 123	NESA
	161 to 164	NNSC
	334 to 337	NESF
	377 to 380	NCSG
	441 to 444	NESA
	548 to 551	NESN
	637 to 640	NAST
	972 to 975	NESL
	1013 to 1016	NVSD
	1084 to 1087	NATV
	1104 to 1107	NCTG
	1161 to 1164	NGTW
	1171 to 1174	NITT
	1318 to 1321	NESF
	1354 to 1357	NASS
Glycosaminoglycan attachment site	558 to 561	SGWG
	665 to 668	SGWG

Table IV (Continued)

cAMP- or cGMP-dependent protein kinase phosphorylation site	1229 to 1232 1399 to 1402	RRIS RRGS
Protein kinase C phosphorylation site	165 to 167	SGR
	268 to 270	TNR
	379 to 381	SGR
	419 to 421	SRR
	469 to 471	SDK
	506 to 508	STR
	589 to 591	SNR
	593 to 595	SGR
	661 to 663	SCR
	696 to 698	SSR
	746 to 748	TER
	805 to 807	SGR
	815 to 817	TWR
	959 to 961	SVR
	1256 to 1258	SGR
	1349 to 1351	SLK
	1396 to 1398	STR

Table IV (Continued)

Casein kinase II phosphorylation site	44 to 47	TDLE
	71 to 74	TVCD
5	178 to 181	TICD
	245 to 248	SHNE
	253 to 256	TCYD
	258 to 261	SDLE
10	319 to 322	SGSD
	332 to 335	SGNE
	392 to 395	TICD
	439 to 442	TGNE
	606 to 609	TVCD
15	622 to 625	SQLD
	673 to 676	SHSE
	686 to 689	SDME
	760 to 763	TGGE
	765 to 768	SLWD
20	818 to 821	SVCD
	845 to 848	SVGD
	857 to 860	TWAE
	907 to 910	SQCD
	923 to 926	SLCD
	927 to 930	THWD
25	974 to 977	SLLD
	1059 to 1062	TICD
	1106 to 1109	TGTE
	1145 to 1148	SETE
	1233 to 1236	SPAE
30	1241 to 1244	TCED
	1269 to 1272	TVCD
	1402 to 1405	SLEE
	1425 to 1428	TSDD

Table IV (Continued)

N-myristoylation site	67 to 72	GQWGTV
	90 to 95	GCPFSF
	101 to 106	GQAVTR
	119 to 124	GNESAL
	133 to 138	GSHNCY
	160 to 165	GNNSCS
	197 to 202	GCPSSF
	226 to 231	GNELAL
	240 to 245	GNHDCS
	267 to 272	GTNRCM
	304 to 309	GCGTAL
	328 to 333	GVSCSG
	374 to 379	GSNNCS
	411 to 416	GCPFSV
	418 to 423	GSRRAK
	440 to 445	GNESAL
	465 to 470	GVICSD
	547 to 552	GNESNI
	588 to 593	GSNRCS
	632 to 637	GMGLGN
	668 to 673	GNNDCS
	679 to 684	GVICSD
	695 to 700	GSSRCA
	712 to 717	GILCAN
	720 to 725	GMNIAE
	758 to 763	GCTGGE
	853 to 858	GNGLTW
	891 to 896	GVVCSR
	944 to 949	GTALST
	985 to 990	GAPPCI
	992 to 997	GNTVSV
	1078 to 1083	GCGVAF
	1121 to 1126	GQHDCR
	1132 to 1137	GVICSE
	1162 to 1167	GTWGSV
	1185 to 1190	GCGENG
	1265 to 1270	GSWGTV
	1288 to 1293	GCGSAL
	1302 to 1307	GQGTGT
	1331 to 1336	GQSDCG
	1342 to 1347	GVRCSG
	1422 to 1427	GTRTSD

Table IV (Continued)

N-myristoylation site (Cont'd)	1443 to 1438 1444 to 1449	GCEDAS GVLPAS
Amidation site	1167 to 1170	VGRR
Speract receptor repeated (SRR) domain signature	53 to 90 160 to 197 267 to 304 1041 to 1078 1251 to 1288	See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2
Scavenger receptor cysteine-rich (SRCR) domain	51 to 148 158 to 255 265 to 362 372 to 469 479 to 576 586 to 683 693 to 790 798 to 895 903 to 1000 1039 to 1136 1146 to 1243 1249 to 1346	See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2 See Fig. 2

Among the domains that occur in TANGO 234 protein are SRR domains and SRCR domains. In one embodiment, the protein of the invention has at least one domain that is at least 55%, preferably at least about 65%, more preferably at least about 75%, yet more preferably at least about 85%, and most preferably at least about 95% identical to one of these domains. In other embodiments, the protein has at least two of the SRR and SRCR domains described herein in Table IV. In other embodiments, the protein has at least one SRR domain and at least one SRCR domain.

The SRR domain is named after a receptor domain identified in a sea urchin egg protein designated speract. The consensus sequence of this domain (using standard one-letter amino acid codes, wherein X is any amino acid residue) is as follows.

-G-X<sub>5</sub>-G-X<sub>2</sub>-E-X<sub>6</sub>-W-G-X<sub>2</sub>-C-X<sub>3</sub>-(F or Y or W)-X<sub>8</sub>-C-X<sub>3</sub>-G-

Speract is a transmembrane glycoprotein of 500 amino acid residues (Dangott et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:2128-2132). Structurally, this receptor consists of a large extracellular domain of 450 residues, followed by a transmembrane region and a small cytoplasmic domain of 12 amino acid residues. The extracellular domain contains four repeats of an approximately 115 amino acid domain. There are 17 amino acid residues that are perfectly conserved in the four repeats in speract, including six cysteine residues, six glycine residues, and two glutamate residues. TANGO 234 has five SRR domains, in which 16 of the 17 conserved speract residues are present of four of the SRR domains and 15 are present in the remaining SRR domain. This domain is designated the speract receptor repeated domain. The amino acid sequence of mammalian macrophage scavenger receptor type I (MSRI) exhibits such a domain (Freeman et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8810-8814). MSRI proteins are membrane glycoproteins implicated in the pathologic deposition of cholesterol in arterial walls during atherogenesis. TANGO 234 is involved in one or more physiological processes related to cholesterol deposition and atherogenesis, as well as other vascular and cardiovascular disorders.

Scavenger receptor cysteine-rich (SRCR) domains are disulfide rich extracellular domains which are present in certain cell surface and secreted proteins. Proteins having SRCR domains exhibit diverse ligand binding specificity. For example, in addition to modified lipoproteins, some of these proteins bind a variety of surface components of pathogenic microorganisms, and some of the proteins bind apoptotic cells. SRCR domains are also involved in mediating immune development and response. Other SRCR-containing proteins are involved in binding of modified lipoproteins (e.g. oxidized low density lipoprotein {LDL}) by specialized macrophages, leading to the formation of macrophages filled with cholesteryl ester droplets (i.e. foam cells). TANGO 234 is involved in one or more physiological processes in which these other SRCR domain-containing proteins are involved, such as LDL uptake and metabolism, regulation of serum cholesterol level, atherogenesis, atherosclerosis,

bacterial or viral infections, immune development, and generation and perseverance of immune responses.

WC1 is a ruminant protein having an SRCR domain. WC1 and gamma delta T-cell receptor are the only known gamma delta T-cell specific antigens.

Antibodies which bind specifically with WC1 induce growth arrest in IL-2-dependent gamma delta T-cell and augment proliferation of gamma delta T-cells in an autologous mixed lymphocyte reaction or in the presence of anti-CD2 or anti-CD5 antibodies.

Injection of antibodies which bind specifically with WC1 into calves results in long-lasting depletion of gamma delta T-cells. Furthermore, antibodies which bind specifically with WC1 can be used to purify gamma delta T-cells.

Gamma delta T-cells are involved in a variety of physiological processes. For example, these cells are potential mediators of allergic airway inflammation and lyme disease. Furthermore, these cells are involved in natural resistance to viral infections and can mediate autoimmune diseases. Elimination of gamma delta T-cells by injection of antibodies which bind specifically therewith can affect the outcomes of these disorders.

TANGO 234 is likely the human orthologue of ruminant protein WC1, and thus is involved with the physiological processes described above in humans. An alignment of the amino acid sequences of (human) TANGO 234 and bovine WC1 protein is shown in Figures 2G, 2H, and 2I. In this alignment (made using the ALIGN software {Myers and Miller (1989) *CABIOS*, ver. 2.0}; pam120.mat scoring matrix; gap penalties -12/-4), the proteins are 40.4% identical. An alignment of the nucleotide sequences of the ORFs encoding (human) TANGO 234 and bovine WC1 protein is shown in Figures 2J through 2Q. The two ORFs are 54.3% identical, as assessed using the same software and parameters.

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that human TANGO 234 protein includes a 40 amino acid signal peptide (amino acid residues 1 to 40 of SEQ ID NO: 11; SEQ ID NO: 12) preceding the mature TANGO 234 protein (amino acid residues 41 to 4386 of SEQ ID NO: 11; SEQ ID NO: 13). Human TANGO 234 protein includes an

extracellular domain (amino acid residues 41 to 1359 of SEQ ID NO: 11; SEQ ID NO: 14); a transmembrane domain (amino acid residues 1360 to 1383 of SEQ ID NO: 11; SEQ ID NO: 15); and a cytoplasmic domain (amino acid residues 1384 to 1453 of SEQ ID NO: 11; SEQ ID NO: 16).

Figure 2F depicts a hydrophilicity plot of human TANGO 234 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 40 of SEQ ID NO: 11 is the signal sequence of human TANGO 234 (SEQ ID NO: 12). The hydrophobic region which corresponds to amino acid residues 1360 to 1383 of SEQ ID NO: 11 is the transmembrane domain of human TANGO 234 (SEQ ID NO: 15). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human TANGO 234 protein from about amino acid residue 225 to about amino acid residue 250 appears to be located at or near the surface of the protein, while the region from about amino acid residue 990 to about amino acid residue 1000 appears not to be located at or near the surface.

The predicted molecular weight of human TANGO 234 protein without modification and prior to cleavage of the signal sequence is about 159.3 kilodaltons. The predicted molecular weight of the mature human TANGO 234 protein without modification and after cleavage of the signal sequence is about 154.7 kilodaltons.

Chromosomal mapping to identify the location of the gene encoding human TANGO 234 protein indicated that the gene was located at chromosomal location h12p13 (with synteny to mo6). Flanking chromosomal markers include WI-6980 and GATA8A09.43. Nearby human loci include IBD2 (inflammatory bowel disease 2), FPF (familial periodic fever), and HPDR2 (hypophosphatemia vitamin D resistant rickets 2). Nearby genes are KLRC (killer cell receptor cluster), DRPLA (dentatorubro-pallidoluysian atrophy), GAPD (glyceraldehyde-3-phosphate dehydrogenase, and PXR1 (peroxisome receptor 1). Murine chromosomal mapping

indicated that the murine orthologue is located near the scr (scruffy) locus. Nearby mouse genes include drpla (dentatorubral phillidolusian atrophy), prp (proline rich protein), and kap (kidney androgen regulated protein).

Northern analysis experiments indicated that mRNA corresponding to the cDNA encoding TANGO 234 is expressed in the tissues listed in Table V, wherein "++" indicates moderate expression, "+" indicates lower expression, and "-" indicates no detectable expression.

Table V

Animal	Tissue	Relative Level of Expression
Human	spleen	++
	fetal lung	++
	lung	+
	thymus	+
	bone marrow	-
	peripheral blood leukocytes	-

Biological function of TANGO 234 proteins, nucleic acids encoding them, and modulators of these molecules

TANGO 234 proteins are involved in disorders which affect both tissues in which they are normally expressed and tissues in which they are normally not expressed. Based on the observation that TANGO 234 is expressed in human fetal lung, spleen, and, to a lesser extent in adult lung and thymus tissue, TANGO 234 protein is involved in one or more biological processes which occur in these tissues. In particular, TANGO 234 is involved in modulating growth, proliferation, survival, differentiation, and activity of cells including, but not limited to, lung, spleen, thymus bone marrow, hematopoietic, peripheral blood leukocytes, and fetal cells of the animal in which it is normally expressed. Thus, TANGO 234 has a role in disorders which affect these cells and their growth, proliferation, survival, differentiation, and activity. Expression of TANGO 234 in an animal is also involved in modulating growth,

proliferation, survival, differentiation, and activity of cells and viruses which are foreign to the host (i.e. bacterial, fungal, and viral infections).

Homology of human TANGO 234 with bovine WC1 protein indicates that TANGO 234 has physiological functions in humans analogous to the functions of WC1 in ruminants. Thus, TANGO 234 is involved in modulating growth, proliferation, survival, differentiation, and activity of gamma delta T cells. For example, TANGO 234 affects the ability of gamma delta T cells to interact with chemokines such as interleukin-2. TANGO 234 therefore is involved in the physiological processes associated with allergic airway inflammation, lyme arthritis, resistance to viral infection, auto-immune diseases, and the like.

In addition, occurrence in TANGO 234 of SRR and SRCR domains indicates that TANGO 234 is involved in physiological functions identical or analogous to the functions performed by other proteins having such domains. For example, like other SRR domain-containing proteins, TANGO 234 modulates cholesterol deposition in arterial walls, and is thus involved in development and persistence of atherogenesis and arteriosclerosis, as well as other vascular and cardiovascular disorders. Like other SRCR domain-containing proteins, TANGO 234 is involved in uptake and metabolism of LDL, regulation of serum cholesterol level, and can modulate these processes as well as the processes of atherogenesis, arteriosclerosis, immune development, and generation and perseverance of immune responses to bacterial, fungal, and viral infections.

#### TANGO 265

A cDNA encoding at least a portion of human TANGO 265 protein was isolated from a human fetal spleen cDNA library. The human TANGO 265 protein is predicted by structural analysis to be a transmembrane membrane protein, although it can exist in a secreted form as well.

The full length of the cDNA encoding human TANGO 265 protein (Figure 3; SEQ ID NO: 17) is 3104 nucleotide residues. The ORF of this cDNA,

nucleotide residues 32 to 2314 of SEQ ID NO: 17 (i.e. SEQ ID NO: 18), encodes a 761-amino acid transmembrane protein (Figure 3; SEQ ID NO: 19).

The invention thus includes purified TANGO 265 protein, both in the form of the immature 761 amino acid residue protein (SEQ ID NO: 19) and in the form of the mature 730 amino acid residue protein (SEQ ID NO: 21). Mature TANGO 265 protein can be synthesized without the signal sequence polypeptide at the amino terminus thereof, or it can be synthesized by generating immature TANGO 265 protein and cleaving the signal sequence therefrom.

In addition to full length mature and immature TANGO 265 proteins, the invention includes fragments, derivatives, and variants of TANGO 265 protein, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA molecule having the nucleotide sequence listed in SEQ ID NO: 17 or some portion thereof, such as the portion which encodes mature TANGO 265 protein, immature TANGO 265 protein, or a domain of TANGO 265 protein. These nucleic acids are collectively referred to as nucleic acids of the invention.

TANGO 265 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features.

A common domain present in TANGO 265 proteins is a signal sequence. As used herein, a signal sequence includes a peptide of at least about 10 amino acid residues in length which occurs at the amino terminus of membrane-bound proteins and which contains at least about 45% hydrophobic amino acid residues such as alanine, leucine, isoleucine, phenylalanine, proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least about 10 to 35 amino acid residues, preferably about 10 to 20 amino acid residues, and has at least about 35-60%, more preferably 40-50%, and more preferably at least about 45% hydrophobic residues. A signal sequence serves to direct a protein containing such a sequence to a

lipid bilayer. Thus, in one embodiment, a TANGO 265 protein contains a signal sequence corresponding to amino acid residues 1 to 31 of SEQ ID NO: 19 (SEQ ID NO: 20). The signal sequence is cleaved during processing of the mature protein.

TANGO 265 proteins can also include an extracellular domain. The human TANGO 265 protein extracellular domain is located from about amino acid residue 32 to about amino acid residue 683 of SEQ ID NO: 17. TANGO 265 can alternately exist in a secreted form, such as a mature protein having the amino acid sequence of amino acid residues 32 to 761 or residues 32 to about 683 of SEQ ID NO: 19.

TANGO 265 proteins can also include a transmembrane domain. In one embodiment, a TANGO 265 protein of the invention contains a transmembrane domain corresponding to about amino acid residues 684 to 704 of SEQ ID NO: 19 (SEQ ID NO: 23).

In addition, TANGO 265 proteins include a cytoplasmic domain, particularly including proteins having a carboxyl-terminal cytoplasmic domain. The human TANGO 265 cytoplasmic domain is located from about amino acid residue 705 to amino acid residue 761 of SEQ ID NO: 19 (SEQ ID NO: 24).

TANGO 265 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Table VI, as predicted by computerized sequence analysis of TANGO 265 proteins using amino acid sequence comparison software (comparing the amino acid sequence of TANGO 265 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}). In certain embodiments, a protein of the invention has at least 1, 2, 4, 6, 10, 15, or 20 or more of the post-translational modification sites listed in Table VI.

Table VI

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 19	Amino Acid Sequence
N-glycosylation site	120 to 123 135 to 138 496 to 499 607 to 610	NETQ NVTH NCSV NGLS
Glycosaminoglycan attachment site	70 to 73	SGDG
cAMP- or cGMP-dependent protein kinase phosphorylation site	108 to 111 116 to 119 281 to 284	RKKS KKKS KKWT
Protein kinase C phosphorylation site	106 to 108 262 to 264 361 to 363 366 to 368 385 to 387 533 to 535 555 to 557 721 to 723 738 to 740	SDR TSR TSR TYR SDK SWK SLR TLR SPK
Casein kinase II phosphorylation site	152 to 155 176 to 179 250 to 253 342 to 345 411 to 414 498 to 501 502 to 505 574 to 577 738 to 741 745 to 748	TFIE SPFD TASE SLLD SGVE SVYE SCVD SILE SPKE SASD

Table VI (Continued)

N-myristoylation site	79 to 84 191 to 196 331 to 336 412 to 417 437 to 442 620 to 625 671 to 676	GAREAI GMLYSG GGTRSS GVEYTR GTTTGS GLYQCW GAALAA
Sema domain	64 to 478	See Fig. 3

An exemplary domains which occurs in TANGO 265 proteins is a sema domain. In one embodiment, the protein of the invention has at least one domain that is at least 55%, preferably at least about 65%, more preferably at least about 75%, yet more preferably at least about 85%, and most preferably at least about 95% identical to one of the sema domains described herein in Table VI.

Sema domains occur in semaphorin proteins. Semaphorins are a large family of secreted and transmembrane proteins, some of which function as repellent signals during neural axon guidance. The sema domain and a variety of semaphorin proteins in which it occurs are described, for example, in Winberg et al. (1998 *Cell* 95:903-916). Sema domains also occur in human hepatocyte growth factor receptor (Swissprot Accession no. P08581) and the similar neuronal and epithelial transmembrane receptor protein (Swissprot Accession no. P51805). The presence of an sema domain in human TANGO 265 protein indicates that TANGO 265 is involved in one or more physiological processes in which the semaphorins are involved, has biological activity in common with one or more of the semaphorins, or both.

Human TANGO 265 protein exhibits considerable sequence similarity to murine semaphorin B protein (GenBank Accession no. X85991), as indicated herein in Figure 3D. Figure 3D depicts an alignment of the amino acid sequences of human TANGO 265 protein (SEQ ID NO: 19) and murine semaphorin B protein (SEQ ID NO: 76). In this alignment (pam120.mat scoring matrix, gap penalties -12/-4), the amino acid sequences of the proteins are 82.3% identical. Figures 3E through 3H depict an alignment of the nucleotide sequences of cDNA encoding human TANGO 265 protein

(SEQ ID NOs: 17) and murine cDNA encoding semaphorin B protein (SEQ ID NO: 77). In this alignment (pam120.mat scoring matrix, gap penalties -12/-4), the nucleic acid sequences of the cDNAs are 76.2% identical. Thus, TANGO 265 is the human orthologue of murine semaphorin B and shares functional similarities to that protein.

It is known that semaphorins are bifunctional, capable of functioning either as attractive axonal guidance proteins or as repellent axonal guidance proteins (Wong et al. (1997) *Development* 124:3597-3607). Furthermore, semaphorins bind with neuronal cell surface proteins designated plexins, which are expressed on both neuronal cells and cells of the immune system (Comeau et al. (1998) *Immunity* 8:473-482; Jin and Strittmatter (1997) *J. Neurosci.* 17:6256-6263).

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that human TANGO 265 protein includes a 31 amino acid signal peptide (amino acid residues 1 to 31 of SEQ ID NO: 19; SEQ ID NO: 20) preceding the mature TANGO 265 protein (amino acid residues 32 to 761 of SEQ ID NO: 19; SEQ ID NO: 21). Human TANGO 265 protein includes an extracellular domain (amino acid residues 32 to 683 of SEQ ID NO: 19; SEQ ID NO: 22); a transmembrane domain (amino acid residues 684 to 704 of SEQ ID NO: 19; SEQ ID NO: 23); and a cytoplasmic domain (amino acid residues 705 to 761 of SEQ ID NO: 19; SEQ ID NO: 24).

Figure 3I depicts a hydrophilicity plot of human TANGO 265 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 31 of SEQ ID NO: 19 is the signal sequence of human TANGO 265 (SEQ ID NO: 20). The hydrophobic region which corresponds to amino acid residues 684 to 704 of SEQ ID NO: 19 is the transmembrane domain of human TANGO 265 (SEQ ID NO: 23). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human TANGO 265 protein from about amino acid residue 350 to about amino acid residue 375 appears to be located at

or near the surface of the protein, while the region from about amino acid residue 230 to about amino acid residue 250 appears not to be located at or near the surface.

The predicted molecular weight of human TANGO 265 protein without modification and prior to cleavage of the signal sequence is about 83.6 kilodaltons.

The predicted molecular weight of the mature human TANGO 265 protein without modification and after cleavage of the signal sequence is about 80.2 kilodaltons.

Chromosomal mapping was performed by computerized comparison of TANGO 265 cDNA sequences against a chromosomal mapping database in order to identify the approximate location of the gene encoding human TANGO 265 protein.

This analysis indicated that the gene was located on chromosome 1 between markers D1S305 and D1S2635.

Biological function of TANGO 265 proteins, nucleic acids encoding them, and modulators of these molecules

TANGO 265 proteins are involved in disorders which affect both tissues in which they are normally expressed and tissues in which they are normally not expressed. Based on the observation that TANGO 265 is expressed in human fetal spleen, involvement of TANGO 202 protein in immune system development and modulation is indicated.

The presence of the sema domain in TANGO 265 indicates that this protein is involved in development of neuronal and epithelial tissues and also functions as a repellant protein which guides axonal development. TANGO 265 modulates nerve growth and regeneration and also modulates growth and regeneration of other epithelial tissues.

The observation that TANGO 265 shares significant identity with murine semaphorin B suggests that it has activity identical or analogous to the activity of this protein. These observations indicate that TANGO 265 modulates growth, proliferation, survival, differentiation, and activity of neuronal cells and immune system cells. Thus, TANGO 265 protein is useful, for example, for guiding neural axon development, for modulating differentiation of cells of the immune system, for

modulating cytokine production by cells of the immune system, for modulating reactivity of cells of the immune system toward cytokines, for modulating initiation and persistence of an inflammatory response, and for modulating proliferation of epithelial cells.

### TANGO 273

A cDNA encoding at least a portion of human TANGO 273 protein was isolated from a lipopolysaccharide- (LPS-)stimulated human osteoblast cDNA library. The corresponding murine cDNA was isolated from an LPS-stimulated murine osteoblast cDNA library. The human and murine TANGO 273 proteins are predicted by structural analysis to be transmembrane proteins.

The full length of the cDNA encoding human TANGO 273 protein (Figure 4; SEQ ID NO: 25) is 2964 nucleotide residues. The ORF of this cDNA, nucleotide residues 135 to 650 of SEQ ID NO: 25 (i.e. SEQ ID NO: 26), encodes a 172-amino acid transmembrane protein (Figure 4; SEQ ID NO: 27).

The invention thus includes purified human TANGO 273 protein, both in the form of the immature 172 amino acid residue protein (SEQ ID NO: 27) and in the form of the mature 150 amino acid residue protein (SEQ ID NO: 29). The invention also includes purified murine TANGO 273 protein, both in the form of the immature 172 amino acid residue protein (SEQ ID NO: 74) and in the form of the mature 150 amino acid residue protein (SEQ ID NO: 44). Mature human or murine TANGO 273 proteins can be synthesized without the signal sequence polypeptide at the amino terminus thereof, or they can be synthesized by generating immature TANGO 273 protein and cleaving the signal sequence therefrom.

In addition to full length mature and immature human and murine TANGO 273 proteins, the invention includes fragments, derivatives, and variants of these TANGO 273 proteins, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

5 The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA molecule having the nucleotide sequence listed in SEQ ID NO: 25 or some portion thereof or SEQ ID NO: 73 or some portion thereof, such as the portion which encodes mature TANGO 273 protein, immature TANGO 273 protein, or a domain of TANGO 273 protein. These nucleic acids are collectively referred to as nucleic acids of the invention.

10 TANGO 273 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features. This family includes, by way of example, the human and murine TANGO 273 proteins.

15 A common domain of TANGO 273 proteins is a signal sequence. As used herein, a signal sequence includes a peptide of at least about 10 amino acid residues in length which occurs at the amino terminus of membrane-bound proteins and which contains at least about 45% hydrophobic amino acid residues such as alanine, leucine, isoleucine, phenylalanine, proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least about 10 to 35 amino acid residues, preferably about 10 to 20 amino acid residues, and has at least about 35-60%, more preferably 40-50%, and more preferably at least about 45% hydrophobic residues. A signal sequence serves to direct a protein containing such a sequence to a lipid bilayer. Thus, in one embodiment, a TANGO 273 protein contains a signal sequence corresponding to amino acid residues 1 to 22 of SEQ ID NO: 27 (SEQ ID NO: 28) or to amino acid residues 1 to 22 of SEQ ID NO: 74. The signal sequence is cleaved during processing of the mature protein.

25 TANGO 273 proteins can also include an extracellular domain. The human TANGO 273 protein extracellular domain is located from about amino acid residue 23 to about amino acid residue 60 of SEQ ID NO: 27, and the murine TANGO 273 protein extracellular domain is located from about amino acid residue 23 to about amino acid residue 60 of SEQ ID NO: 74.

The present invention also includes TANGO 273 proteins having a transmembrane domain. As used herein, a “transmembrane domain” refers to an amino acid sequence having at least about 15 to 30 amino acid residues in length and which contains at least about 65-70% hydrophobic amino acid residues such as alanine, leucine, phenylalanine, protein, tyrosine, tryptophan, or valine. In a preferred embodiment, a transmembrane domain contains at least about 15 to 20 amino acid residues, preferably about 20 to 25 amino acid residues, and has at least about 60-80%, more preferably 65-75%, and more preferably at least about 70% hydrophobic residues. Thus, in one embodiment, a human TANGO 273 protein of the invention contains a transmembrane domain corresponding to about amino acid residues 61 to 81 of SEQ ID NO: 27 (SEQ ID NO: 31). In another embodiment, a murine TANGO 273 protein of the invention contains a transmembrane domain corresponding to about amino acid residues 61 to 81 of SEQ ID NO: 74.

In addition, TANGO 273 proteins include a cytoplasmic domain. The human TANGO 273 cytoplasmic domain is located from about amino acid residue 82 to amino acid residue 172 of SEQ ID NO: 27 (SEQ ID NO: 32), and the murine TANGO 273 cytoplasmic domain is located from about amino acid residue 82 to amino acid residue 172 of SEQ ID NO: 74.

TANGO 273 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Tables VII and VIII, as predicted by computerized sequence analysis of human and murine TANGO 273 proteins using amino acid sequence comparison software (comparing the amino acid sequence of TANGO 273 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}). In certain embodiments, a protein of the invention has at least 1, 2, 3, 4, 5, or all 6 of the post-translational modification sites listed in Table VII. In other embodiments, the protein of the invention has at least 1, 2, 3, 4, 5, 6, or all 7 of the post-translational modification sites listed in Table VIII.

Table VII

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 27	Amino Acid Sequence
N-glycosylation site	97 to 100	NVSY
Casein kinase II phosphorylation site	41 to 44	SYED
N-myristoylation site	31 to 36 47 to 52 70 to 75 131 to 136	GLYPTY GSRCCV GVLFCC GNSMAM
Src Homology 3 (SH3) domain binding site	86 to 90 103 to 107 113 to 117 121 to 125 140 to 145 151 to 155 160 to 164	YPPPL QPPNP QPGPP DPGGP VPPNSP CPPPP TPPPP

Table VIII

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 74	Amino Acid Sequence
N-glycosylation site	97 to 100	NVSY
Casein kinase II phosphorylation site	41 to 44	SYED
N-myristoylation site	31 to 36 47 to 52 70 to 75 131 to 136	GLYPTY GSRCCV GVLFCC GNTMAM

Table VIII (Cont'd)

Src Homology 3 (SH3) domain binding site	86 to 90	YPPPL
	103 to 107	QPPNP
	115 to 119	GPPYY
	121 to 125	DPGGP
	141 to 145	QPNSP
	151 to 155	YPPPP
	160 to 164	TPPPP
Amidation site	1 to 4	MGRR

The amino acid sequence of TANGO 273 protein includes about seven potential proline-rich Src homology 3 (SH3) domain binding sites nearer the cytoplasmic portion of the protein. SH3 domains mediate specific assembly of protein complexes, presumably by interacting with proline-rich protein domains (Morton and Campbell (1994) *Curr. Biol.* 4:615-617). SH3 domains also mediate interactions between proteins involved in transmembrane signal transduction. Coupling of proteins mediated by SH3 domains has been implicated in a variety of physiological systems, including those involving regulation of cell growth and proliferation, endocytosis, and activation of respiratory burst.

SH3 domains have been described in the art (e.g. Mayer et al. (1988) *Nature* 332:272-275; Musacchio et al. (1992) *FEBS Lett.* 307:55-61; Pawson and Schlessinger (1993) *Curr. Biol.* 3:434-442; Mayer and Baltimore (1993) *Trends Cell Biol.* 3:8-13; Pawson (1993) *Nature* 373:573-580), and occur in a variety of cytoplasmic proteins, including several (e.g. protein tyrosine kinases) involved in transmembrane signal transduction. Among the proteins in which one or more SH3 domains occur are protein tyrosine kinases such as those of the Src, Abl, Bkt, Csk and ZAP70 families, mammalian phosphatidylinositol-specific phospholipases C-gamma-1 and -2, mammalian phosphatidylinositol 3-kinase regulatory p85 subunit, mammalian Ras GTPase-activating protein (GAP), proteins which mediate binding of guanine

nucleotide exchange factors and growth factor receptors (e.g. vertebrate GRB2,  
*Caenorhabditis elegans* sem-5, and *Drosophila* DRK proteins), mammalian Vav  
oncoprotein, guanidine nucleotide releasing factors of the CDC 25 family (e.g. yeast  
CDC25, yeast SCD25, and fission yeast ste6 proteins), MAGUK proteins (e.g.  
5 mammalian tight junction protein ZO-1, vertebrate erythrocyte membrane protein p55,  
*C. elegans* protein lin-2, rat protein CASK, and mammalian synaptic proteins  
SAP90/PSD-95, CHAPSYN-110/PSD-93, SAP97/DLG1, and SAP102), proteins  
which interact with vertebrate receptor protein tyrosine kinases (e.g. mammalian  
cytoplasmic protein Nck and oncoprotein Crk), chicken Src substrate p80/85 protein  
10 (cortactin), human hemopoietic lineage cell specific protein Hs1, mammalian  
dihydrouridine-sensitive L-type calcium channel beta subunit, human myasthenic  
syndrome antigen B (MSYB), mammalian neutrophil cytosolic activators of NADPH  
oxidase (e.g. p47 {NCF-1}, p67 {NCF-2}, and *C. elegans* protein B0303.7) myosin  
heavy chains (MYO3) from amoebae, from slime molds, and from yeast, vertebrate and  
15 *Drosophila* spectrin and fodrin alpha chain proteins, human amphiphysin, yeast actin-  
binding proteins ABP1 and SLA3, yeast protein BEM1, fission yeast protein scd2  
(ral3), yeast BEM1-binding proteins BOI2 (BEB1) and BOB1 (BOI1), yeast fusion  
protein FUS1, yeast protein RSV167, yeast protein SSU81, yeast hypothetical proteins  
YAR014c, YFR024c, YHL002w, YHR016c, YJL020C, and YHR114w, hypothetical  
20 fission yeast protein SpAC12C2.05c, and *C. elegans* hypothetical protein F42H10.3.  
Of these proteins, multiple SH3 domains occur in vertebrate GRB2 protein, *C. elegans*  
sem-5 protein, *Drosophila* DRK protein, oncoprotein Crk, mammalian neutrophil  
cytosolic activators of NADPH oxidase p47 and p67, yeast protein BEM1, fission yeast  
protein scd2, yeast hypothetical protein YHR114w, mammalian cytoplasmic protein  
25 Nck, *C. elegans* neutrophil cytosolic activator of NADPH oxidase B0303.7, and yeast  
actin-binding protein SLA1. Of these proteins, three or more SH3 domains occur in  
mammalian cytoplasmic protein Nck, *C. elegans* neutrophil cytosolic activator of  
NADPH oxidase B0303.7, and yeast actin-binding protein SLA1. The presence of SH3  
domain binding sites in TANGO 273 indicates that TANGO 273 interacts with one or  
30 more of these and other SH3 domain-containing proteins and is thus involved in

physiological processes in which one or more of these or other SH3 domain-containing proteins are involved.

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that human TANGO 273 protein includes a 22 amino acid signal peptide (amino acid residues 1 to 22 of SEQ ID NO: 27; SEQ ID NO: 28) preceding the mature TANGO 273 protein (amino acid residues 23 to 172 of SEQ ID NO: 27; SEQ ID NO: 29). Human TANGO 273 protein includes an extracellular domain (amino acid residues 23 to 60 of SEQ ID NO: 27; SEQ ID NO: 30); a transmembrane domain (amino acid residues 61 to 81 of SEQ ID NO: 27; SEQ ID NO: 31); and a cytoplasmic domain (amino acid residues 82 to 172 of SEQ ID NO: 27; SEQ ID NO: 32).

Figure 4F depicts a hydrophilicity plot of human TANGO 273 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 22 of SEQ ID NO: 27 is the signal sequence of human TANGO 273 (SEQ ID NO: 28). The hydrophobic region which corresponds to amino acid residues 61 to 81 of SEQ ID NO: 27 is the transmembrane domain of human TANGO 273 (SEQ ID NO: 31). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human TANGO 273 protein from about amino acid residue 100 to about amino acid residue 120 appears to be located at or near the surface of the protein, while the region from about amino acid residue 130 to about amino acid residue 140 appears not to be located at or near the surface.

Chromosomal mapping was performed by computerized comparison of TANGO 273 cDNA sequences against a chromosomal mapping database in order to identify the approximate location of the gene encoding human TANGO 265 protein. This analysis indicated that the gene was located on chromosome 7 between markers D7S2467 and D7S2552.

The predicted molecular weight of human TANGO 273 protein without modification and prior to cleavage of the signal sequence is about 19.2 kilodaltons. The predicted molecular weight of the mature human TANGO 273 protein without modification and after cleavage of the signal sequence is about 16.8 kilodaltons.

5 The full length of the cDNA encoding murine TANGO 273 protein (Figure 4; SEQ ID NO: 72) is 2915 nucleotide residues. The ORF of this cDNA, nucleotide residues 137 to 650 of SEQ ID NO: 72 (i.e. SEQ ID NO: 73), encodes a 172-amino acid transmembrane protein (Figure 4; SEQ ID NO: 74).

10 The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that murine TANGO 273 protein includes a 22 amino acid signal peptide (amino acid residues 1 to 22 of SEQ ID NO: 74) preceding the mature TANGO 273 protein (amino acid residues 23 to 172 of SEQ ID NO: 74; SEQ ID NO: 44). Murine TANGO 273 protein includes an extracellular domain (amino acid residues 23 to 60 of SEQ ID NO: 74); a transmembrane domain (amino acid residues 61 to 81 of SEQ ID NO: 74); and a cytoplasmic domain (amino acid residues 82 to 172 of SEQ ID NO: 74).

15 Figure 4G depicts a hydrophilicity plot of murine TANGO 273 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 22 of SEQ ID NO: 74 is the signal sequence of murine TANGO 273. As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of murine TANGO 273 protein from about amino acid residue 100 to about amino acid residue 120 appears to be located at or near the surface of the protein, while the region from about amino acid residue 130 to about amino acid residue 140 appears not to be located at or near the surface.

25 The predicted molecular weight of murine TANGO 273 protein without modification and prior to cleavage of the signal sequence is about 19.4 kilodaltons.

The predicted molecular weight of the mature murine TANGO 273 protein without modification and after cleavage of the signal sequence is about 17.1 kilodaltons.

Human and murine TANGO 273 proteins exhibit considerable sequence similarity, as indicated herein in Figure 4E. Figure 4E depicts an alignment of human and murine TANGO 273 protein amino acid sequences (SEQ ID NOs: 27 and 74, respectively). In this alignment (pam120.mat scoring matrix, gap penalties -12/-4), the proteins are 89.5% identical. Alignment of the ORF encoding human TANGO 273 protein and the ORF encoding murine TANGO 273 protein using the same software and parameters indicated that the nucleotide sequences are 84.1% identical.

Biological function of TANGO 273 proteins, nucleic acids encoding them, and modulators of these molecules

cDNAs encoding the human and murine TANGO 273 proteins were each isolated from LPS-stimulated osteoblast cDNA libraries. These proteins are involved in bone-related metabolism, homeostasis, and development disorders. Thus, proteins and nucleic acids of the invention which are identical to, similar to, or derived from human and murine TANGO 273 proteins and nucleic acids encoding them are useful for preventing, diagnosing, and treating, among others, bone-related disorders such as osteoporosis, cancer, skeletal development disorders, bone fragility, and the like.

The fact that TANGO 273 is expressed in tissues which were exposed to LPS indicates that TANGO 273 mediates one or more physiological responses of cells to bacterial infection. Thus, TANGO 273 is involved in one or more of detection of bacteria in a tissue in which it is expressed, movement of cells with relation to sites of bacterial infection, production of biological molecules which inhibit bacterial infection, and production of biological molecules which alleviate cellular or other physiological damage wrought by bacterial infection.

Occurrence in TANGO 273 protein of multiple SH3 domain binding sites indicates that TANGO 273 protein interacts with one or more SH3 domain-

containing proteins. Thus, TANGO 273 protein mediates binding of proteins (i.e. binding of proteins to TANGO 273 and to one another to form protein complexes) in cells in which it is expressed. TANGO 273 is also involved in transduction of signals between the exterior environment of cells (i.e. including from other cells) and the interior of cells in which it is expressed. TANGO 273 mediates regulation of cell growth and proliferation, endocytosis, activation of respiratory burst, and other physiological processes triggered by transmission of a signal via a protein with which TANGO 273 interacts.

#### TANGO 286

A cDNA encoding at least a portion of human TANGO 286 protein was isolated from a human keratinocyte cDNA library. The human TANGO 286 protein is predicted by structural analysis to be a secreted protein.

The full length of the cDNA encoding TANGO 286 protein (Figure 5; SEQ ID NO: 33) is 1980 nucleotide residues. The ORF of this cDNA, nucleotide residues 133 to 1497 of SEQ ID NO: 33 (i.e. SEQ ID NO: 34), encodes a 455-amino acid secreted protein (Figure 5; SEQ ID NO: 35).

The invention thus includes purified TANGO 286 protein, both in the form of the immature 455 amino acid residue protein (SEQ ID NO: 35) and in the form of the mature 432 amino acid residue protein (SEQ ID NO: 37). Mature TANGO 286 protein can be synthesized without the signal sequence polypeptide at the amino terminus thereof, or it can be synthesized by generating immature TANGO 286 protein and cleaving the signal sequence therefrom.

In addition to full length mature and immature TANGO 286 proteins, the invention includes fragments, derivatives, and variants of these TANGO 286 proteins, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA

molecule having the nucleotide sequence listed in SEQ ID NO: 33 or some portion thereof, such as the portion which encodes mature TANGO 286 protein, immature TANGO 286 protein, or a domain of TANGO 286 protein. These nucleic acids are collectively referred to as nucleic acids of the invention.

5 TANGO 286 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features.

10 A common domain of TANGO 286 proteins is a signal sequence. As used herein, a signal sequence includes a peptide of at least about 10 amino acid residues in length which occurs at the amino terminus of membrane-bound proteins and which contains at least about 45% hydrophobic amino acid residues such as alanine, leucine, isoleucine, phenylalanine, proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least about 10 to 35 amino acid residues, preferably about 10 to 20 amino acid residues, and has at least about 35-60%, more preferably 40-50%, and more preferably at least about 45% hydrophobic residues. A signal sequence serves to direct a protein containing such a sequence to a lipid bilayer. Thus, in one embodiment, a TANGO 286 protein contains a signal sequence corresponding to amino acid residues 1 to 23 of SEQ ID NO: 35 (SEQ ID NO: 36). The signal sequence is cleaved during processing of the mature protein.

20 TANGO 286 is a secreted soluble protein (i.e. a secreted protein having a single extracellular domain), as indicated by computerized sequence analysis and comparison of the amino acid sequence of TANGO 286 with related proteins, such as the soluble proteins designated bactericidal permeability increasing (BPI) protein and recombinant endotoxin neutralizing polypeptide (RENK).

25 TANGO 286 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Table IX, as predicted by computerized sequence analysis of TANGO 286 proteins using amino acid sequence comparison software (comparing the amino acid sequence of TANGO 286 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}). In

30

certain embodiments, a protein of the invention has at least 1, 2, 4, 6, 10, 15, or 20 or more of the post-translational modification sites listed in Table IX.

Table IX

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 35	Amino Acid Sequence
N-glycosylation site	79 to 82	NFSN
	92 to 95	NTSL
	113 to 116	NIST
	161 to 164	NLST
	173 to 176	NYTL
	205 to 208	NLTD
	249 to 252	NLTL
	303 to 306	NFTL
	320 to 323	NSTV
	363 to 366	NRSN
Protein kinase C phosphorylation site	35 to 37	TQR
	362 to 364	SNR
	429 to 431	SSK
Casein kinase II phosphorylation site	63 to 66	SGSE
	130 to 133	SFAE
	163 to 166	STLE
	169 to 172	TKID
	175 to 178	TLLD
	183 to 186	SSPE
	253 to 256	STEE
	321 to 324	STVE
	365 to 368	SNIE
	409 to 412	SDIE

Table IX (Cont'd)

N-myristoylation site	42 to 47	GVQAGM
	269 to 274	GNVLSR
Lipid-binding serum glycoprotein domain	12 to 427	see Fig. 5

Certain lipid-binding serum glycoproteins, such as LPS-binding protein (LBP), bactericidal permeability-increasing protein (BPI), cholesteryl ester transfer protein (CETP), and phospholipid transfer protein (PLTP), share regions of sequence similarity which are herein designated a lipid-binding serum glycoprotein domain (Schumann et al., (1990) *Science* 249:1429-1431; Gray et al., (1989) *J. Biol. Chem.* 264:9505-9509; Day et al., (1994) *J. Biol. Chem.* 269:9388-9391). The consensus pattern of lipid-binding serum glycoprotein domains is as follows (using standard single letter amino acid abbreviations wherein X is any amino acid residue).

-(P or A)-(G or A)-(L or I or V or M or C)-X<sub>2</sub>-R-(I or V)-(S or T)-  
X<sub>3</sub>-L-X<sub>(4 or 5)</sub>-(E or Q)-X<sub>4</sub>-(L or I or V or M)-X<sub>(0 or 1)</sub>-(E or Q or K)-X<sub>8</sub>-P-  
(e.g. amino acid residues 28-60 of SEQ ID NO: 35)

Proteins in which a lipid-binding serum glycoprotein domain occurs are often structurally related and exhibit related physiological activities. LBP binds to lipid A moieties of bacterial LPS and, once bound thereto, induces secretion of  $\alpha$ -tumor necrosis factor, apparently by interacting with the CD14 receptor. BPI also binds LPS and exerts a cytotoxic effect on Gram-negative bacteria (Elsbach, (1998) *J. Leukoc. Biol.* 64:14-18). CETP is involved in transfer of insoluble cholesteryl esters during reverse cholesterol transport. PLTP appears to be involved in phospholipid transport and modulation of serum HDL particles.

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that TANGO 286 protein includes a 23 amino acid signal peptide (amino acid residues 1 to 23 of SEQ ID NO: 35; SEQ ID NO: 36) preceding the mature TANGO 286 protein (amino acid residues 24 to 455 of SEQ ID NO: 35; SEQ ID NO: 37). Human TANGO 286 protein is a secreted soluble protein.

Figure 5C depicts a hydrophilicity plot of TANGO 286 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human TANGO 286 protein from about amino acid residue 420 to about amino acid residue 435 appears to be located at or near the surface of the protein, while the region from about amino acid residue 325 to about amino acid residue 345 appears not to be located at or near the surface.

The predicted molecular weight of TANGO 286 protein without modification and prior to cleavage of the signal sequence is about 50.9 kilodaltons. The predicted molecular weight of the mature TANGO 286 protein without modification and after cleavage of the signal sequence is about 48.2 kilodaltons.

The gene encoding human TANGO 286 protein was determined to be located on chromosome 22 by comparison of matching genomic clones such as the clones assigned GenBank Accession numbers W16806 and AL021937.

A portion of TANGO 286 protein exhibits significant amino acid homology with a region of the human chromosome region 22q12-13 genomic nucleotide sequence having GenBank Accession number AL021937. Alignment of a 45 kilobase nucleotide sequence encoding TANGO 286 with AL021937, however, indicated the presence in TANGO 286 of exons which differ from those disclosed in L021937 (pam120.mat scoring matrix; gap penalties -12/-4). This region of chromosome 22 comprises an immunoglobulin lambda chain C (IGLC) pseudogene, the Ret finger protein-like 3 (RFPL3) and Ret finger protein-like 3 antisense (RFPL3S) genes, a gene encoding a novel immunoglobulin lambda chain V family protein, a novel gene encoding a protein similar both to mouse RGDS protein (RALGDS, RALGEF, guanine nucleotide dissociation stimulator A) and to rabbit oncogene RSC, a novel gene encoding the human orthologue of worm F16A11.2 protein, a novel gene encoding a protein similar both to BPI and to rabbit liposaccharide-binding protein, and a 5'-portion of a novel gene. This region also comprises various ESTs, STSs, GSSs,

genomic marker D22S1175, a ca repeat polymorphism and putative CpG islands. TANGO 286 protein thus shares one or more structural or functional features of these molecules.

TANGO 286 protein exhibits considerable sequence similarity with BPI protein, having 23.9% amino acid sequence identity therewith, as assessed using the ALIGN v. 2.0 computer software using a pam120.mat scoring matrix and gap penalties of -12/-4. TANGO 286 protein also exhibits considerable sequence similarity with recombinant endotoxin neutralizing polypeptide (RENP), having 24.5% amino acid sequence identity therewith, as assessed using the ALIGN software. Physiological activities of BPI protein and RENP have been described (e.g. Gabay et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:5610-5614; Elsbach, (1998) *J. Leukoc. Biol.* 64:14-18; Mahadeva et al., (1997) *Chest* 112:1699-1701; International patent application WO96/34873). RENP, for example, binds LPS and neutralizes bacterial endotoxins. BPI, RENP, and other proteins in which a lipid-binding serum glycoprotein domain occurs bind LPS and neutralize bacterial endotoxins, and are therefore useful for preventing, detecting, and treating LPS-related disorders such as shock, disseminated intravascular coagulation, anemia, thrombocytopenia, adult respiratory distress syndrome, renal failure, liver disease, and disorders associated with Gram negative bacterial infections. In addition to the physiological conditions described above, BPI protein is known to be involved in vasculitis and bronchiectasis, in that antibodies which bind specifically with BPI protein are present in at least some patients afflicted with these disorders (Mahadeva et al., *supra*).

Biological function of TANGO 286 proteins, nucleic acids encoding them, and modulators of these molecules

Expression of TANGO 286 in keratinocyte library indicates that this protein is involved in a disorders which involve keratinocytes. Such disorders include, for example, disorders involving extracellular matrix abnormalities, dermatological disorders, ocular disorders, inappropriate hair growth (e.g. baldness), infections of the nails of the fingers and toes, scalp disorders (e.g. dandruff), and the like.

The fact that TANGO 286 protein contains a lipid-binding serum glycoprotein domain indicates that TANGO 286 is involved in one or more physiological processes in which these other lipid-binding serum glycoprotein domain-containing proteins are involved. Thus, TANGO 286 is involved in one or more of lipid transport, metabolism, serum lipid particle regulation, host antimicrobial defensive mechanisms, and the like.

Human TANGO 286 shares physiological functionality with other proteins in which a lipid-binding serum glycoprotein domains occurs (e.g. LBP, BPI protein, CETP, and PLTP). Based on the amino acid sequence similarity of TANGO 286 with BPI protein and with RENP, TANGO 286 protein exhibits physiological activities exhibited by these proteins. Thus, TANGO 286 proteins are useful for preventing, diagnosing, and treating, among others, lipid transport disorders, lipid metabolism disorders, disorders of serum lipid particle regulation, obesity, disorders involving insufficient or inappropriate host antimicrobial defensive mechanisms, vasculitis, bronchiectasis, LPS-related disorders such as shock, disseminated intravascular coagulation, anemia, thrombocytopenia, adult respiratory distress syndrome, renal failure, liver disease, and disorders associated with Gram negative bacterial infections, such as bacteremia, endotoxemia, sepsis, and the like.

TANGO 294

A cDNA encoding at least a portion of human TANGO 294 protein was isolated from a human pulmonary artery smooth muscle cell cDNA library. The human

TANGO 294 protein is predicted by structural analysis to be a transmembrane membrane protein. No expression of DNA encoding TANGO 294 was detected in human heart, brain, placenta, lung, liver, skeletal muscle, kidney, or pancreas tissues.

The full length of the cDNA encoding TANGO 294 protein (Figure 6; SEQ ID NO: 45) is 2044 nucleotide residues. The ORF of this cDNA, nucleotide residues 126 to 1394 of SEQ ID NO: 45 (i.e. SEQ ID NO: 46), encodes a 423-amino acid transmembrane protein (Figure 6; SEQ ID NO: 47).

The invention includes purified TANGO 294 protein, both in the form of the immature 423 amino acid residue protein (SEQ ID NO: 47) and in the form of the mature 390 amino acid residue protein (SEQ ID NO: 49). Mature TANGO 294 protein can be synthesized without the signal sequence polypeptide at the amino terminus thereof, or it can be synthesized by generating immature TANGO 294 protein and cleaving the signal sequence therefrom.

In addition to full length mature and immature TANGO 294 proteins, the invention includes fragments, derivatives, and variants of TANGO 294 protein, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA molecule having the nucleotide sequence listed in SEQ ID NO: 45 or some portion thereof, such as the portion which encodes mature TANGO 294 protein, immature TANGO 294 protein, or a domain of TANGO 294 protein. These nucleic acids are collectively referred to as nucleic acids of the invention.

TANGO 294 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features.

Also included within the scope of the invention are TANGO 294 proteins having a signal sequence. As used herein, a signal sequence includes a peptide of at least about 10 amino acid residues in length which occurs at the amino terminus of membrane-bound proteins and which contains at least about 45% hydrophobic

amino acid residues such as alanine, leucine, isoleucine, phenylalanine, proline, tyrosine, tryptophan, or valine. In a preferred embodiment, a signal sequence contains at least about 10 to 35 amino acid residues, preferably about 10 to 20 amino acid residues, and has at least about 35-60%, more preferably 40-50%, and more preferably at least about 45% hydrophobic residues. A signal sequence serves to direct a protein containing such a sequence to a lipid bilayer. Thus, in one embodiment, a TANGO 294 protein contains a signal sequence corresponding to amino acid residues 1 to 33 of SEQ ID NO: 47 (SEQ ID NO: 48). The signal sequence is cleaved during processing of the mature protein.

The naturally-occurring form of TANGO 294 protein is a secreted protein (i.e. not comprising the predicted signal sequence). However, in variant forms, TANGO 294 proteins can be transmembrane proteins which include an extracellular domain. In this transmembrane variant form, the predicted TANGO 294 protein extracellular domain is located from about amino acid residue 34 to about amino acid residue 254 of SEQ ID NO: 47, the predicted cytoplasmic domain is located from about amino acid residue 280 to amino acid residue 423 of SEQ ID NO: 47 (SEQ ID NO: 52), and the predicted transmembrane domain is located from about amino acid residues 255 to 279 of SEQ ID NO: 47 (SEQ ID NO: 51).

TANGO 294 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Table X, as predicted by computerized sequence analysis of TANGO 294 proteins using amino acid sequence comparison software (comparing the amino acid sequence of TANGO 294 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}). In certain embodiments, a protein of the invention has at least 1, 2, 4, 6, 10, 15, or 20 or more of the post-translational modification sites listed in Table X.

Table X

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 47	Amino Acid Sequence
N-glycosylation site	48 to 51	NISE
	113 to 116	NNSL
	285 to 288	NMSR
	413 to 416	NLSQ
Protein kinase C phosphorylation site	12 to 14	SHR
	138 to 140	SRK
	217 to 219	TVK
Casein kinase II phosphorylation site	155 to 158	SYDE
	175 to 178	TGQE
	198 to 201	TMPE
	360 to 363	SNPE
Tyrosine kinase phosphorylation site	174 to 182	KTGQEKIYY
N-myristoylation site	99 to 104	GLVGGA
	130 to 135	GNSRGN
	188 to 193	GTTMGF
	277 to 282	GGFNTN
Amidation site	240 to 243	FGKK
Lipase serine active site	180 to 189	IYYVGYSQGT
Alpha/beta hydrolase fold domain	125 to 404	See Fig. 6

Alpha/beta hydrolase fold domains occur in a wide variety of enzymes (Ollis et al., (1992) *Protein Eng.* 5:197-211). The alpha/beta fold domain is a conserved topological domain in which sequence homology is not necessarily conserved. Conservation of topology in the alpha/beta fold domain preserves arrangement of catalytic residues, even though those residues, and the reactions they catalyze, can vary. In many enzymes, particularly including alpha/beta hydrolases, this

domain encompasses the active site of the enzyme. In one embodiment, the protein of the invention has at least one domain that is at least 55%, preferably at least about 65%, more preferably at least about 75%, yet more preferably at least about 85%, and most preferably at least about 95% identical to the alpha/beta hydrolase fold domain described herein in Table X.

The signal peptide prediction program SIGNALP (Nielsen et al. (1997) *Protein Engineering* 10:1-6) predicted that human TANGO 294 protein includes a 33 amino acid signal peptide (amino acid residues 1 to 33 of SEQ ID NO: 47; SEQ ID NO: 48) preceding the mature TANGO 294 protein (amino acid residues 34 to 423 of SEQ ID NO: 47; SEQ ID NO: 49). Human TANGO 294 protein is a soluble secreted protein. However, in the transmembrane variant form, human TANGO 294 protein includes an extracellular domain (amino acid residues 34 to 254 of SEQ ID NO: 47; SEQ ID NO: 50); a transmembrane domain (amino acid residues 255 to 279 of SEQ ID NO: 47; SEQ ID NO: 51); and a cytoplasmic domain (amino acid residues 280 to 423 of SEQ ID NO: 47; SEQ ID NO: 52).

Figure 6D depicts a hydrophilicity plot of human TANGO 294 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic region which corresponds to amino acid residues 1 to 33 of SEQ ID NO: 47 is the signal sequence of human TANGO 294 (SEQ ID NO: 49). The hydrophobic region which corresponds to amino acid residues 255 to 279 of SEQ ID NO: 47 is the predicted transmembrane domain of human TANGO 294 (SEQ ID NO: 51). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human TANGO 294 protein from about amino acid residue 130 to about amino acid residue 150 appears to be located at or near the surface of the protein, while the region from about amino acid residue 90 to about amino acid residue 100 appears not to be located at or near the surface.

The predicted molecular weight of human TANGO 294 protein without modification and prior to cleavage of the signal sequence is about 48.2 kilodaltons. The predicted molecular weight of the mature human TANGO 294 protein without modification and after cleavage of the signal sequence is about 44.2 kilodaltons.

It may be that amino acid residues 1 to 15 of SEQ ID NO: 47 do not occur in TANGO 294 protein. However, it is recognized that amino acid residues 16 to 33 of SEQ ID NO: 47 form a functional signal sequence even in the absence of residues 1 to 15. The amino acid sequence (and hence the properties) of mature TANGO 294 protein are unaffected by occurrence or non-occurrence of amino acid residues 1 to 15 of immature TANGO 294 protein.

Human TANGO 294 protein exhibits considerable sequence similarity (i.e. about 75% amino acid sequence identity) to lingual and gastric lipase proteins of rat (Swissprot Accession no. P04634; Docherty et al. (1985) *Nucleic Acids Res.* 13:1891-1903), dog (Swissprot Accession no. P80035; Carriere et al. (1991) *Eur. J. Biochem.* 202:75-83), and human (Swissprot Accession no. P07098; Bernbaeck and Blaekberg (1987) *Biochim. Biophys. Acta* 909:237-244), as assessed using the ALIGN v. 2.0 computer software using a pam12.mat scoring matrix and gap penalties of -12/-4. TANGO 294 is distinct from the known human lipase, as indicated in Figure 6C. Figure 6C depicts an alignment of the amino acid sequences of human TANGO 294 protein (SEQ ID NO: 47) and the known human lipase protein (SEQ ID NO: 75), as assessed using the same software and parameters. In this alignment (pam120.mat scoring matrix, gap penalties -12/-4), the amino acid sequences of the proteins are 49.8% identical. TANGO 294 also is distinct from the known human lysosomal acid lipase, as indicated in Figure 6E. Figure 6E depicts an alignment of the amino acid sequences of human TANGO 294 protein (SEQ ID NO: 47) and the known human lysosomal acid lipase protein (SEQ ID NO: 41). In this alignment (pam120.mat scoring matrix, gap penalties -12/-4), the amino acid sequences of the proteins are 56.9% identical.

TANGO 294 is a human lipase distinct from the known human lipase and the known human lysosomal acid lipase. Furthermore, in view of the comparisons

of the amino acid sequences of TANGO 294 and the two human lipases and the nature of transcriptional initiation sites, it is recognized that the transcriptional start site can correspond to either of the methionine residues located at residues 1 and 15 of SEQ ID NO: 47. The present invention thus includes proteins in which the initially transcribed amino acid residue is the methionine residue at position 1 of SEQ ID NO: 47 and proteins in which the initially transcribed amino acid residue is the methionine residue at position 15 of SEQ ID NO: 47 (i.e. proteins in which the amino acid sequence of TANGO 294 does not include residues 1 to 14 of SEQ ID NO: 47). Furthermore, because amino acid residues 1 to 14 of SEQ ID NO: 47 are predicted to be part of a signal sequence, it is recognized that the protein not comprising this portion of the amino acid sequence will nonetheless exhibit a functional signal sequence at its amino terminus.

Biological function of TANGO 294 proteins, nucleic acids encoding them, and modulators of these molecules

The sequence similarity of TANGO 294 and mammalian lingual, gastric, and lysosomal acid lipase proteins indicates that TANGO 294 is involved in physiological processes identical or analogous to those involving these lipases. Thus, TANGO 294 is involved in facilitating absorption and metabolism of fat. TANGO 294 can thus be used, for example, to prevent, detect, and treat disorders relating to fat absorption and metabolism, such as inadequate expression of gastric/pancreatic lipase, cystic fibrosis, exocrine pancreatic insufficiency, obesity, medical treatments which alter fat absorption, and the like.

TANGO 294 protein is known to be expressed in human pulmonary artery smooth muscle tissue. This indicates that TANGO 294 protein is involved in transportation and metabolism of fats and lipids in the human vascular and cardiovascular systems. Thus, TANGO 294 proteins of the invention can be used to prevent, detect, and treat disorders involving these body systems.

## INTERCEPT 296

A cDNA encoding at least a portion of human INTERCEPT 296 protein was isolated from a human esophagus cDNA library. The human INTERCEPT 296 protein is predicted by structural analysis to be a transmembrane protein having three or more transmembrane domains. Expression of DNA encoding INTERCEPT 296 tissue has been detected by northern analysis of human lung tissue. In human lung tissue, two moieties corresponding to INTERCEPT 296 have been identified in Northern blots. It is recognized that these two moieties may represent alternatively polyadenylated INTERCEPT 296 mRNAs or alternatively spliced INTERCEPT 296 mRNAs. It has furthermore been observed that INTERCEPT 296 does not appear to be expressed in any of heart, brain, placenta, skeletal muscle, kidney, and pancreas tissues.

The full length of the cDNA encoding INTERCEPT 296 protein (Figure 7; SEQ ID NO: 53) is 2133 nucleotide residues. The ORF of this cDNA, nucleotide residues 70 to 1098 of SEQ ID NO: 53 (i.e. SEQ ID NO: 54), encodes a 343-amino acid transmembrane protein (Figure 7; SEQ ID NO: 55).

The invention includes purified INTERCEPT 296 protein, which has the amino acid sequence listed in SEQ ID NO: 55. In addition to full length INTERCEPT 296 proteins, the invention includes fragments, derivatives, and variants of these INTERCEPT 296 proteins, as described herein. These proteins, fragments, derivatives, and variants are collectively referred to herein as polypeptides of the invention or proteins of the invention.

The invention also includes nucleic acid molecules which encode a polypeptide of the invention. Such nucleic acids include, for example, a DNA molecule having the nucleotide sequence SEQ ID NO: 53 or some portion thereof, such as the portion which encodes INTERCEPT 296 protein or a domain thereof. These nucleic acids are collectively referred to as nucleic acids of the invention.

INTERCEPT 296 proteins and nucleic acid molecules encoding them comprise a family of molecules having certain conserved structural and functional features, such as the five transmembrane domains which occur in the protein.

INTERCEPT 296 comprises at least five transmembrane domains, at least three cytoplasmic domains, and at least two extracellular domains. INTERCEPT 296 does not appear to comprise a cleavable signal sequence. Amino acid residues 1 to 70 of SEQ ID NO: 55 likely directs insertion of the protein into the cytoplasmic membrane. There are at least two mechanisms by which this can occur. Sequence analysis of residues 1 to 70 of SEQ ID NO: 55 indicates that this entire region may represent a signal sequence or that residues 1 to 47 represent a signal sequence, with residues 48-70 representing a transmembrane region. Human INTERCEPT 296 protein extracellular domains are located from about amino acid residue 70 to about amino acid residue 182 (SEQ ID NO: 57) and from about amino acid residue 228 to about amino acid residue 249 (SEQ ID NO: 58) of SEQ ID NO: 55. Human INTERCEPT 296 cytoplasmic domains are located from about amino acid residue 43 to amino acid residue 50 (SEQ ID NO: 64), from about amino acid residue 205 to amino acid residue 210 (SEQ ID NO: 65), and from amino acid residue 272 to amino acid residue 343 (SEQ ID NO: 66) of SEQ ID NO: 55. The five transmembrane domains of INTERCEPT 296 are located from about amino acid residues 24 to 42 (SEQ ID NO: 59), 51 to 70 (SEQ ID NO: 60), 183 to 204 (SEQ ID NO: 61), 211 to 227 (SEQ ID NO: 62), and 250 to 271 (SEQ ID NO: 63) of SEQ ID NO: 55.

INTERCEPT 296 proteins typically comprise a variety of potential post-translational modification sites (often within an extracellular domain), such as those described herein in Table XI, as predicted by computerized sequence analysis of INTERCEPT 296 proteins using amino acid sequence comparison software (comparing the amino acid sequence of INTERCEPT 296 with the information in the PROSITE database {rel. 12.2; Feb, 1995} and the Hidden Markov Models database {Rel. PFAM 3.3}). In certain embodiments, a protein of the invention has at least 1, 2, 4, 6, 10, 15, or 20 or more of the post-translational modification sites listed in Table XI.

Table XI

Type of Potential Modification Site or Domain	Amino Acid Residues of SEQ ID NO: 55	Amino Acid Sequence
N-glycosylation site	71 to 74	NFSS
	84 to 87	NTSY
	109 to 112	NITL
	121 to 124	NETI
	284 to 287	NQSV
Protein kinase C phosphorylation site	86 to 88	SYK
	131 to 133	TWR
	162 to 164	TPR
	304 to 306	SPR
	313 to 315	SPK
	326 to 328	STK
Casein kinase II phosphorylation site	286 to 289	SVDE
	296 to 299	SPEE
	309 to 312	SMAD
Tyrosine kinase phosphorylation site	148 to 156	KGLPDPVL Y
N-myristoylation site	79 to 84	GQVSTN
	100 to 105	GLQVGL
	107 to 112	GVNITL
	265 to 270	GLAMAV

Figure 7C depicts a hydrophilicity plot of INTERCEPT 296 protein. Relatively hydrophobic regions are above the dashed horizontal line, and relatively hydrophilic regions are below the dashed horizontal line. The hydrophobic regions which corresponds to amino acid residues 24 to 42, 51 to 70, 183 to 204, 211 to 227, and 250 to 271 of SEQ ID NO: 55 are the transmembrane domains of human

INTERCEPT 296 (SEQ ID NOs: 59 through 63, respectively). As described elsewhere herein, relatively hydrophilic regions are generally located at or near the surface of a protein, and are more frequently effective immunogenic epitopes than are relatively hydrophobic regions. For example, the region of human INTERCEPT 296 protein from about amino acid residue 120 to about amino acid residue 140 appears to be located at or near the surface of the protein, while the region from about amino acid residue 95 to about amino acid residue 110 appears not to be located at or near the surface.

The predicted molecular weight of INTERCEPT 296 protein without modification and prior to cleavage of the signal sequence is about 37.8 kilodaltons. The predicted molecular weight of the mature INTERCEPT 296 protein without modification and after cleavage of the signal sequence is about 30.2 kilodaltons.

Figure 7D depicts an alignment of the amino acid sequences of human INTERCEPT 296 protein (SEQ ID NO: 55) and *Caenorhabditis elegans* C06E1.3 related protein (SEQ ID NO: 399). In this alignment (pam120.mat scoring matrix, gap penalties -12/-4), the amino acid sequences of the proteins are 26.8% identical. The *C. elegans* protein has five predicted transmembrane domains.

Biological function of INTERCEPT 296 proteins, nucleic acids encoding them, and modulators of these molecules

The cDNA encoding INTERCEPT 296 protein was obtained from a human esophagus cDNA library, and INTERCEPT 296 is expressed in lung tissue. The INTERCEPT 296-related proteins and nucleic acids of the invention are therefore useful for prevention, detection, and treatment of disorders of the human lung and esophagus. Such disorders include, for example, various cancers, bronchitis, cystic fibrosis, respiratory infections (e.g. influenza, bronchiolitis, pneumonia, and tuberculosis), asthma, emphysema, chronic bronchitis, bronchiectasis, pulmonary edema, pleural effusion, pulmonary embolus, adult and infant respiratory distress syndromes, heartburn, and gastric reflux esophageal disease.

Tables A and B summarize sequence data corresponding to the human proteins herein designated TANGO 202, TANGO 234, TANGO 265, TANGO 273, TANGO 286, TANGO 294, and INTERCEPT 296.

Table A

Protein Designation	SEQ ID NOs			Depicted in Figure #	ATCC Accession #
	cDNA	ORF	Protein		
TANGO 202	1	2	3	1	207219
TANGO 234	9	10	11	2	207184
TANGO 265	17	18	19	3	207228
TANGO 273	25	26	27	4	207185
TANGO 286	33	34	35	5	207220
TANGO 294	45	46	47	6	207220
INTERCEPT 296	53	54	55	7	207220

Table B

Protein Desig.	Signal Sequence	Mature Protein	Extracellular Domain(s)	Transmembrane Domain(s)	Cytoplamic Domain(s)
SEQ ID NOs					
TANGO 202 (variant)	1 to 19 (1 to 19)	4 (4)	20 to 475 (20 to 475)	5 (5)	6 (6)
TANGO 234	1 to 40	12	41 to 1453	13	14
TANGO 265	1 to 31	20	32 to 761	21	22
TANGO 273	1 to 22	28	23 to 172	29	30
TANGO 286	1 to 23	36	24 to 455	37	37
TANGO 294 (variant 1)	1 to 33 (15 to 33)	48 (40)	34 to 423 (34 to 423)	49 (49)	50 (50)
<variant 2> {variant 3}	<1 to 33> {15 to 33}	<48> {40}	<34 to 423> {34 to 423}	<49> {49}	<49> {49}
INTERCEPT 296	N/A		1 to 343	55	56
					57
					58
					59
					60
					61
					62
					63

Amino Acid Residues

Various aspects of the invention are described in further detail in the following subsections.

#### I. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules that encode a polypeptide of the invention or a biologically active portion thereof, as well as nucleic acid molecules sufficient for use as hybridization probes to identify nucleic acid molecules encoding a polypeptide of the invention and fragments of such nucleic acid molecules suitable for use as PCR primers for the amplification or mutation of nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid molecule. Preferably, an "isolated" nucleic acid molecule is free of sequences (preferably protein-encoding sequences) which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kB, 4 kB, 3 kB, 2 kB, 1 kB, 0.5 kB or 0.1 kB of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule having the nucleotide sequence of all or a portion of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or a complement thereof, or which

has a nucleotide sequence comprising one of these sequences, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of the nucleic acid sequences of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, as a hybridization probe, nucleic acid molecules of the invention can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook et al., eds., *Molecular Cloning: A Laboratory Manual, 2nd ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

A nucleic acid molecule of the invention can be amplified using cDNA, mRNA or genomic DNA as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to all or a portion of a nucleic acid molecule of the invention can be prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or a portion thereof. A nucleic acid molecule which is complementary to a given nucleotide sequence is one which is sufficiently complementary to the given nucleotide sequence that it can hybridize to the given nucleotide sequence thereby forming a stable duplex.

Moreover, a nucleic acid molecule of the invention can comprise only a portion of a nucleic acid sequence encoding a full length polypeptide of the invention for example, a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a polypeptide of the invention. The nucleotide sequence determined from the cloning one gene allows for the generation of probes and primers designed for use in identifying and/or cloning homologs in other cell types, e.g., from other tissues, as well as homologs from other mammals. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically

comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 15, preferably about 25, more preferably about 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, or 400 or more consecutive nucleotides of the sense or anti-sense sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or of a naturally occurring mutant of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, or 73.

Probes based on the sequence of a nucleic acid molecule of the invention can be used to detect transcripts or genomic sequences encoding the same protein molecule encoded by a selected nucleic acid molecule. The probe comprises a label group attached thereto, e.g., a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as part of a diagnostic test kit for identifying cells or tissues which mis-express the protein, such as by measuring levels of a nucleic acid molecule encoding the protein in a sample of cells from a subject, e.g., detecting mRNA levels or determining whether a gene encoding the protein has been mutated or deleted.

A nucleic acid fragment encoding a biologically active portion of a polypeptide of the invention can be prepared by isolating a portion of SEQ ID NO: 2, 10, 18, 26, 34, 46, 54, 68, or 73, expressing the encoded portion of the polypeptide protein (e.g., by recombinant expression *in vitro*), and assessing the activity of the encoded portion of the polypeptide.

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, or 73 due to degeneracy of the genetic code and thus encode the same protein as that encoded by the nucleotide sequence of SEQ ID NO: 2, 10, 18, 26, 34, 46, 54, 68, or 73.

In addition to the nucleotide sequences of SEQ ID NOS: 2, 10, 18, 26, 34, 46, 54, 68, or 73, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequence can exist within a population (e.g., the human population). Such genetic polymorphisms can

exist among individuals within a population due to natural allelic variation. An allele is one of a group of genes which occur alternatively at a given genetic locus.

As used herein, the phrase "allelic variant" refers to a nucleotide sequence which occurs at a given locus or to a polypeptide encoded by the nucleotide sequence.

As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a polypeptide of the invention. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of a given gene. Alternative alleles can be identified by sequencing the gene of interest in a number of different individuals. This can be readily carried out by using hybridization probes to identify the same genetic locus in a variety of individuals. Any and all such nucleotide variations and resulting amino acid polymorphisms or variations that are the result of natural allelic variation and that do not alter the functional activity are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding proteins of the invention from other species (homologs), which have a nucleotide sequence which differs from that of the rat protein described herein are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologs of a cDNA of the invention can be isolated based on their identity to human nucleic acid molecules using the rat cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. For example, a cDNA encoding a soluble form of a membrane-bound protein of the invention isolated based on its hybridization to a nucleic acid molecule encoding all or part of the membrane-bound form. Likewise, a cDNA encoding a membrane-bound form can be isolated based on its hybridization to a nucleic acid molecule encoding all or part of the soluble form.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 (25, 40, 60, 80, 100, 150, 200, 250, 300, 350, 400, 450, 550, 650, 700, 800, 900, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000, 3500, 4000, 4500, or 4928) nucleotides in length and hybridizes under stringent

conditions to the nucleic acid molecule comprising the nucleotide sequence, preferably the coding sequence, of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, or 73, or a complement thereof. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% (65%, 70%, preferably 75%) identical to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6× sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2× SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, or 73, or a complement thereof, corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein).

In addition to naturally-occurring allelic variants of a nucleic acid molecule of the invention sequence that can exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation thereby leading to changes in the amino acid sequence of the encoded protein, without altering the biological activity of the protein. For example, one can make nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are not conserved or only semi-conserved among homologs of various species may be non-essential for activity and thus would be likely targets for alteration. Alternatively, amino acid residues that are conserved among the homologs of various species (e.g., murine and human) may be essential for activity and thus would not be likely targets for alteration.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding a polypeptide of the invention that contain changes in amino acid residues that are not essential for activity. Such polypeptides differ in amino acid sequence from SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, or 74, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule includes a nucleotide sequence encoding a protein that includes an amino acid sequence that is at least about 40% identical, 50%, 60%, 70%, 80%, 90%, 95%, or 98% identical to the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, or 74.

An isolated nucleic acid molecule encoding a variant protein can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, or 73, such that one or more amino acid residue substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), non-polar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

In a preferred embodiment, a mutant polypeptide that is a variant of a polypeptide of the invention can be assayed for: (1) the ability to form protein:protein interactions with the polypeptide of the invention; (2) the ability to bind a ligand of the polypeptide of the invention (e.g. another protein identified herein); (3) the ability to bind to a modulator or substrate of the polypeptide of the invention; or (4) the ability to modulate a physiological activity of the protein, such as one of those disclosed herein.

The present invention encompasses antisense nucleic acid molecules, i.e., molecules which are complementary to a sense nucleic acid encoding a polypeptide of the invention, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be complementary to an entire coding strand, or to only a portion thereof, e.g., all or part of the protein coding region (or open reading frame). An antisense nucleic acid molecule can be antisense to all or part of a non-coding region of the coding strand of a nucleotide sequence encoding a polypeptide of the invention. The non-coding regions ("5' and 3' untranslated regions") are the 5' and 3' sequences which flank the coding region and are not translated into amino acids.

An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine,

5 N<sub>6</sub>-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N<sub>6</sub>-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-  
10 N<sub>6</sub>-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an  
15 expression vector into which a nucleic acid has been sub-cloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

20 The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a selected polypeptide of the invention to thereby inhibit expression, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such  
25 that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the

antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

An antisense nucleic acid molecule of the invention can be an  $\alpha$ -anomeric nucleic acid molecule. An  $\alpha$ -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\alpha$ -units, the strands run parallel to each other (Gaultier et al. (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue et al. (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue et al. (1987) *FEBS Lett.* 215:327-330).

The invention also encompasses ribozymes. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes as described in Haselhoff and Gerlach (1988) *Nature* 334:585-591) can be used to catalytically cleave mRNA transcripts to thereby inhibit translation of the protein encoded by the mRNA. A ribozyme having specificity for a nucleic acid molecule encoding a polypeptide of the invention can be designed based upon the nucleotide sequence of a cDNA disclosed herein. For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a Cech et al. U.S. Patent No. 4,987,071; and Cech et al. U.S. Patent No. 5,116,742. Alternatively, an mRNA encoding a polypeptide of the invention can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel and Szostak (1993) *Science* 261:1411-1418.

The invention also encompasses nucleic acid molecules which form triple helical structures. For example, expression of a polypeptide of the invention can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the gene encoding the polypeptide (e.g., the promoter and/or enhancer) to form triple helical structures that prevent transcription of the gene in target cells. See generally Helene (1991) *Anticancer Drug Des.* 6(6):569-84; Helene (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14(12):807-15.

In various embodiments, the nucleic acid molecules of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup et al. (1996) *Bioorganic & Medicinal Chemistry* 4(1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, e.g., DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup et al. (1996), *supra*; Perry-O'Keefe et al. (1996) *Proc. Natl. Acad. Sci. USA* 93: 14670-675.

PNAs can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, e.g., inducing transcription or translation arrest or inhibiting replication. PNAs can also be used, e.g., in the analysis of single base pair mutations in a gene by, e.g., PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, e.g., S1 nucleases (Hyrup (1996), *supra*; or as probes or primers for DNA sequence and hybridization (Hyrup (1996), *supra*; Perry-O'Keefe et al. (1996) *Proc. Natl. Acad. Sci. USA* 93: 14670-675).

In another embodiment, PNAs can be modified, e.g., to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras can be generated which can combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, e.g., RNASE H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and

orientation (Hyrup (1996), *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996), *supra*, and Finn et al. (1996) *Nucleic Acids Res.* 24(17):3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and modified nucleoside analogs. Compounds such as 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite can be used as a link between the PNA and the 5' end of DNA (Mag et al. (1989) *Nucleic Acids Res.* 17:5973-88). PNA monomers are then coupled in a step-wise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn et al. (1996) *Nucleic Acids Res.* 24(17):3357-63). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser et al. (1975) *Bioorganic Med. Chem. Lett.* 5:1119-1124).

In other embodiments, the oligonucleotide can include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (*see, e.g.*, Letsinger et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. WO 88/09810) or the blood-brain barrier (*see, e.g.*, PCT Publication No. WO 89/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (*see, e.g.*, Krol et al. (1988) *Bio/Techniques* 6:958-976) or intercalating agents (*see, e.g.*, Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide can be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

## II. Isolated Proteins and Antibodies

One aspect of the invention pertains to isolated proteins, and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise antibodies directed against a polypeptide of the invention. In one embodiment, the native polypeptide can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, polypeptides of the invention are produced by recombinant DNA

techniques. Alternative to recombinant expression, a polypeptide of the invention can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or substantially free of chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. Thus, protein that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, or 5% (by dry weight) of heterologous protein (also referred to herein as a "contaminating protein"). When the protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, 10%, or 5% of the volume of the protein preparation. When the protein is produced by chemical synthesis, it is preferably substantially free of chemical precursors or other chemicals, i.e., it is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. Accordingly such preparations of the protein have less than about 30%, 20%, 10%, 5% (by dry weight) of chemical precursors or compounds other than the polypeptide of interest.

Biologically active portions of a polypeptide of the invention include polypeptides comprising amino acid sequences sufficiently identical to or derived from the amino acid sequence of the protein (e.g., the amino acid sequence shown in any of SEQ ID NOS: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, or 74), which include fewer amino acids than the full length protein, and exhibit at least one activity of the corresponding full-length protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the corresponding protein. A biologically active portion of a protein of the invention can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by

recombinant techniques and evaluated for one or more of the functional activities of the native form of a polypeptide of the invention.

Preferred polypeptides have the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, or 74. Other useful proteins are substantially identical (e.g., at least about 40%, preferably 50%, 60%, 70%, 80%, 90%, 95%, or 99%) to any of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, or 74 and retain the functional activity of the protein of the corresponding naturally-occurring protein yet differ in amino acid sequence due to natural allelic variation or mutagenesis.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions (e.g., overlapping positions)  $\times$  100). In one embodiment the two sequences are the same length.

The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990) *J. Mol. Biol.* 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino

acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules. *Id.* When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, (1988) *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Yet another useful algorithm for identifying regions of local sequence similarity and alignment is the FASTA algorithm as described in Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85:2444-2448. When using the FASTA algorithm for comparing nucleotide or amino acid sequences, a PAM120 weight residue table can, for example, be used with a *k*-tuple value of 2.

The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, only exact matches are counted.

The invention also provides chimeric or fusion proteins. As used herein, a "chimeric protein" or "fusion protein" comprises all or part (preferably biologically active) of a polypeptide of the invention operably linked to a heterologous polypeptide (i.e., a polypeptide other than the same polypeptide of the invention). Within the fusion protein, the term "operably linked" is intended to indicate that the polypeptide of the invention and the heterologous polypeptide are fused in-frame to each other. The heterologous polypeptide can be fused to the amino-terminus or the carboxyl-terminus of the polypeptide of the invention.

One useful fusion protein is a GST fusion protein in which the polypeptide of the invention is fused to the carboxyl terminus of GST sequences. Such

fusion proteins can facilitate the purification of a recombinant polypeptide of the invention.

In another embodiment, the fusion protein contains a heterologous signal sequence at its amino terminus. For example, the native signal sequence of a polypeptide of the invention can be removed and replaced with a signal sequence from another protein. For example, the gp67 secretory sequence of the baculovirus envelope protein can be used as a heterologous signal sequence (*Current Protocols in Molecular Biology*, Ausubel et al., eds., John Wiley & Sons, 1992). Other examples of eukaryotic heterologous signal sequences include the secretory sequences of melittin and human placental alkaline phosphatase (Stratagene; La Jolla, California). In yet another example, useful prokaryotic heterologous signal sequences include the phoA secretory signal (Sambrook et al., *supra*) and the protein A secretory signal (Pharmacia Biotech; Piscataway, New Jersey).

In yet another embodiment, the fusion protein is an immunoglobulin fusion protein in which all or part of a polypeptide of the invention is fused to sequences derived from a member of the immunoglobulin protein family. The immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a ligand (soluble or membrane-bound) and a protein on the surface of a cell (receptor), to thereby suppress signal transduction *in vivo*. The immunoglobulin fusion protein can be used to affect the bioavailability of a cognate ligand of a polypeptide of the invention. Inhibition of ligand/receptor interaction can be useful therapeutically, both for treating proliferative and differentiative disorders and for modulating (e.g. promoting or inhibiting) cell survival. Moreover, the immunoglobulin fusion proteins of the invention can be used as immunogens to produce antibodies directed against a polypeptide of the invention in a subject, to purify ligands and in screening assays to identify molecules which inhibit the interaction of receptors with ligands.

Chimeric and fusion proteins of the invention can be produced by standard recombinant DNA techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers.

Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (*see, e.g.,* Ausubel et al., *supra*). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A nucleic acid encoding a polypeptide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the polypeptide of the invention.

A signal sequence of a polypeptide of the invention (e.g. the signal sequence in one of SEQ ID NO: 3, 4, 11, 12, 19, 20, 27, 28, 35, 36, 47, 48, 69, and 74) can be used to facilitate secretion and isolation of the secreted protein or other proteins of interest. Signal sequences are typically characterized by a core of hydrophobic amino acids which are generally cleaved from the mature protein during secretion in one or more cleavage events. Such signal peptides contain processing sites that allow cleavage of the signal sequence from the mature proteins as they pass through the secretory pathway. Thus, the invention pertains to the described polypeptides having a signal sequence, as well as to the signal sequence itself and to the polypeptide in the absence of the signal sequence (i.e., the cleavage products). In one embodiment, a nucleic acid sequence encoding a signal sequence of the invention can be operably linked in an expression vector to a protein of interest, such as a protein which is ordinarily not secreted or is otherwise difficult to isolate. The signal sequence directs secretion of the protein, such as from a eukaryotic host into which the expression vector is transformed, and the signal sequence is subsequently or concurrently cleaved. The protein can then be readily purified from the extracellular medium by art recognized methods. Alternatively, the signal sequence can be linked to the protein of interest using a sequence which facilitates purification, such as with a GST domain.

In another embodiment, the signal sequences of the present invention can be used to identify regulatory sequences, e.g., promoters, enhancers, repressors. Since signal sequences are the most amino-terminal sequences of a peptide, the nucleic acids which flank the signal sequence on its amino-terminal side are likely regulatory

sequences which affect transcription. Thus, a nucleotide sequence which encodes all or a portion of a signal sequence can be used as a probe to identify and isolate signal sequences and their flanking regions, and these flanking regions can be studied to identify regulatory elements therein.

The present invention also pertains to variants of the polypeptides of the invention. Such variants have an altered amino acid sequence which can function as either agonists (mimetics) or as antagonists. Variants can be generated by mutagenesis, e.g., discrete point mutation or truncation. An agonist can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of the protein. An antagonist of a protein can inhibit one or more of the activities of the naturally occurring form of the protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the protein of interest. Thus, specific biological effects can be elicited by treatment with a variant of limited function. Treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein can have fewer side effects in a subject relative to treatment with the naturally occurring form of the protein.

Variants of a protein of the invention which function as either agonists (mimetics) or as antagonists can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of the protein of the invention for agonist or antagonist activity. In one embodiment, a variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential protein sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display). There are a variety of methods which can be used to produce libraries of potential variants of the polypeptides of the invention from a degenerate oligonucleotide sequence. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang (1983) *Tetrahedron* 39:3; Itakura et al. (1984) *Annu. Rev. Biochem.*

53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477).

In addition, libraries of fragments of the coding sequence of a polypeptide of the invention can be used to generate a variegated population of polypeptides for screening and subsequent selection of variants. For example, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of the coding sequence of interest with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes amino terminal and internal fragments of various sizes of the protein of interest.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify variants of a protein of the invention (Arkin and Yourvan (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave et al. (1993) *Protein Engineering* 6(3):327-331).

An isolated polypeptide of the invention, or a fragment thereof, can be used as an immunogen to generate antibodies using standard techniques for polyclonal and monoclonal antibody preparation. The full-length polypeptide or protein can be

used or, alternatively, the invention provides antigenic peptide fragments for use as immunogens. The antigenic peptide of a protein of the invention comprises at least 8 (preferably 10, 15, 20, or 30 or more) amino acid residues of the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, or 74, and encompasses an epitope of the protein such that an antibody raised against the peptide forms a specific immune complex with the protein.

Preferred epitopes encompassed by the antigenic peptide are regions that are located on the surface of the protein, e.g., hydrophilic regions. Figures 1G, 1H, 2F, 3I, 4F, 4G, 5C, 6D, and 7C are hydrophobicity plots of the proteins of the invention. These plots or similar analyses can be used to identify hydrophilic regions.

An immunogen typically is used to prepare antibodies by immunizing a suitable (i.e. immunocompetent) subject such as a rabbit, goat, mouse, or other mammal or vertebrate. An appropriate immunogenic preparation can contain, for example, recombinantly-expressed or chemically-synthesized polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or a similar immunostimulatory agent.

Accordingly, another aspect of the invention pertains to antibodies directed against a polypeptide of the invention. The terms "antibody" and "antibody substance" as used interchangeably herein refer to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site which specifically binds an antigen, such as a polypeptide of the invention. A molecule which specifically binds to a given polypeptide of the invention is a molecule which binds the polypeptide, but does not substantially bind other molecules in a sample, e.g., a biological sample, which naturally contains the polypeptide. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab')<sub>2</sub> fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody

molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope.

Polyclonal antibodies can be prepared as described above by immunizing a suitable subject with a polypeptide of the invention as an immunogen.

5 The antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized polypeptide. If desired, the antibody molecules can be harvested or isolated from the subject (e.g., from the blood or serum of the subject) and further purified by well-known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the specific antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497, the human B cell hybridoma technique (Kozbor et al. (1983) *Immunol. Today* 4:72), the EBV-hybridoma technique (Cole et al. (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing hybridomas is well known (*see generally Current Protocols in Immunology* (1994) Coligan et al. (eds.) John Wiley & Sons, Inc., New York, NY). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind the polypeptide of interest, e.g., using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal antibody directed against a polypeptide of the invention can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with the polypeptide of interest. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, U.S. Patent No. 5,223,409; PCT Publication No.

WO 92/18619; PCT Publication No. WO 91/17271; PCT Publication No. WO 92/20791; PCT Publication No. WO 92/15679; PCT Publication No. WO 93/01288; PCT Publication No. WO 92/01047; PCT Publication No. WO 92/09690; PCT Publication No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J.* 12:725-734.

Additionally, recombinant antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in PCT Publication No. WO 87/02671; European Patent Application 184,187; European Patent Application 171,496; European Patent Application 173,494; PCT Publication No. WO 86/01533; U.S. Patent No. 4,816,567; European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura et al. (1987) *Canc. Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; and Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison (1985) *Science* 229:1202-1207; Oi et al. (1986) *Bio/Techniques* 4:214; U.S. Patent 5,225,539; Jones et al. (1986) *Nature* 321:552-525; Verhoeyan et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060.

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Such antibodies can be produced using transgenic mice which are incapable of expressing endogenous immunoglobulin heavy and light chains genes, but which can express human heavy and light chain genes. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and

subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar (1995, *Int. Rev. Immunol.* 13:65-93). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., U.S. Patent 5,625,126; U.S. Patent 5,633,425; U.S. Patent 5,569,825; U.S. Patent 5,661,016; and U.S. Patent 5,545,806. In addition, companies such as Abgenix, Inc. (Freemont, CA), can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a murine antibody, is used to guide the selection of a completely human antibody recognizing the same epitope (Jespers et al. (1994) *Bio/technology* 12:899-903).

An antibody directed against a polypeptide of the invention (e.g., monoclonal antibody) can be used to isolate the polypeptide by standard techniques, such as affinity chromatography or immunoprecipitation. Moreover, such an antibody can be used to detect the protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the polypeptide. The antibodies can also be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or

phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ .

### III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a polypeptide of the invention (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors, expression vectors, are capable of directing the expression of genes to which they are operably linked. In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids (vectors). However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell. This means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for

expression of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, and the like. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein.

The recombinant expression vectors of the invention can be designed for expression of a polypeptide of the invention in prokaryotic (e.g., *E. coli*) or eukaryotic cells (e.g., insect cells (using baculovirus expression vectors), yeast cells or mammalian cells). Suitable host cells are discussed further in Goeddel, *supra*. Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein.

Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification

of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann et al., (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a co-expressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident  $\lambda$  prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada et al. (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari et al. (1987) *EMBO J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al. (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and pPicZ (Invitrogen Corp, San Diego, CA).

Alternatively, the expression vector is a baculovirus expression vector. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith et al. (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed (1987) *Nature* 329:840) and pMT2PC (Kaufman et al. (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook et al., *supra*.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert et al. (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji et al. (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund et al. (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the  $\alpha$ -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an

antisense orientation. That is, the DNA molecule is operably linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an RNA molecule which is antisense to the mRNA encoding a polypeptide of the invention. Regulatory sequences operably linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid, or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub et al. (*Reviews - Trends in Genetics*, Vol. 1(1) 1986).

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic (e.g., *E. coli*) or eukaryotic cell (e.g., insect cells, yeast or mammalian cells).

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection,

lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, et al. (*supra*), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., for resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce a polypeptide of the invention. Accordingly, the invention further provides methods for producing a polypeptide of the invention using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding a polypeptide of the invention has been introduced) in a suitable medium such that the polypeptide is produced. In another embodiment, the method further comprises isolating the polypeptide from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which a sequences encoding a polypeptide of the invention have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous sequences encoding a polypeptide of the invention have been introduced into their genome or homologous recombinant animals in which endogenous encoding a polypeptide of the invention sequences have been altered. Such animals are useful for studying the function and/or activity of the polypeptide and for identifying and/or evaluating modulators of polypeptide activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or

more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, an "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing nucleic acid encoding a polypeptide of the invention (or a homologue thereof) into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the polypeptide of the invention to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, U.S. Patent No. 4,873,191 and in Hogan, *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of mRNA encoding the transgene in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying the transgene can further be bred to other transgenic animals carrying other transgenes.

To create an homologous recombinant animal, a vector is prepared which contains at least a portion of a gene encoding a polypeptide of the invention into

which a deletion, addition or substitution has been introduced to thereby alter, e.g.,  
 functionally disrupt, the gene. In a preferred embodiment, the vector is designed such  
 that, upon homologous recombination, the endogenous gene is functionally disrupted  
 (i.e., no longer encodes a functional protein; also referred to as a "knock out" vector).  
 Alternatively, the vector can be designed such that, upon homologous recombination,  
 the endogenous gene is mutated or otherwise altered but still encodes functional protein  
 (e.g., the upstream regulatory region can be altered to thereby alter the expression of the  
 endogenous protein). In the homologous recombination vector, the altered portion of  
 the gene is flanked at its 5' and 3' ends by additional nucleic acid of the gene to allow  
 for homologous recombination to occur between the exogenous gene carried by the  
 vector and an endogenous gene in an embryonic stem cell. The additional flanking  
 nucleic acid sequences are of sufficient length for successful homologous  
 recombination with the endogenous gene. Typically, several kilobases of flanking  
 DNA (both at the 5' and 3' ends) are included in the vector (*see, e.g.*, Thomas and  
 Capecchi (1987) *Cell* 51:503 for a description of homologous recombination vectors).  
 The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and  
 cells in which the introduced gene has homologously recombined with the endogenous  
 gene are selected (*see, e.g.*, Li et al. (1992) *Cell* 69:915). The selected cells are then  
 injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras  
 (*see, e.g.*, Bradley in *Teratocarcinomas and Embryonic Stem Cells: A Practical  
 Approach*, Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can  
 then be implanted into a suitable pseudopregnant female foster animal and the embryo  
 brought to term. Progeny harboring the homologously recombined DNA in their germ  
 cells can be used to breed animals in which all cells of the animal contain the  
 homologously recombined DNA by germline transmission of the transgene. Methods  
 for constructing homologous recombination vectors and homologous recombinant  
 animals are described further in Bradley (1991) *Current Opinion in Bio/Technology*  
 2:823-829 and in PCT Publication NOS. WO 90/11354, WO 91/01140, WO 92/0968,  
 and WO 93/04169.

In another embodiment, transgenic non-human animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, *see, e.g.*, Lakso et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman et al. (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut et al. (1997) *Nature* 385:810-813 and PCT Publication NOS. WO 97/07668 and WO 97/07669.

#### IV. Pharmaceutical Compositions

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

5 The invention includes methods for preparing pharmaceutical compositions for modulating the expression or activity of a polypeptide or nucleic acid of the invention. Such methods comprise formulating a pharmaceutically acceptable carrier with an agent which modulates expression or activity of a polypeptide or nucleic acid of the invention. Such compositions can further include additional active agents. Thus, the invention further includes methods for preparing a pharmaceutical composition by formulating a pharmaceutically acceptable carrier with an agent which modulates expression or activity of a polypeptide or nucleic acid of the invention and one or more additional active compounds.

10 The agent which modulates expression or activity can, for example, be a small molecule other than a nucleic acid, polypeptide, or antibody of the invention. For example, such small molecules include peptides, peptidomimetics, amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (i.e. including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

15 It is understood that appropriate doses of small molecule agents and protein or polypeptide agents depends upon a number of factors within the ken of the ordinarily skilled physician, veterinarian, or researcher. The dose(s) of these agents will vary, for example, depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the composition is to be administered, if applicable, and the effect which the practitioner desires the agent to have upon the nucleic acid or polypeptide of the invention. Exemplary doses of a small molecule include milligram or microgram amounts per kilogram of subject or sample weight (e.g. about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or

about 1 microgram per kilogram to about 50 micrograms per kilogram). Exemplary doses of a protein or polypeptide include gram, milligram or microgram amounts per kilogram of subject or sample weight (e.g. about 1 microgram per kilogram to about 5 grams per kilogram, about 100 micrograms per kilogram to about 500 milligrams per kilogram, or about 1 milligram per kilogram to about 50 milligrams per kilogram). It is furthermore understood that appropriate doses of one of these agents depend upon the potency of the agent with respect to the expression or activity to be modulated. Such appropriate doses can be determined using the assays described herein. When one or more of these agents is to be administered to an animal (e.g. a human) in order to modulate expression or activity of a polypeptide or nucleic acid of the invention, a physician, veterinarian, or researcher can, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific agent employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediamine-tetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The

parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL (BASF; Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium, and then incorporating the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a

powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches, and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes having monoclonal antibodies incorporated therein or thereon) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

For antibodies, the preferred dosage is 0.1 mg/kg to 100 mg/kg of body weight (generally 10 mg/kg to 20 mg/kg). If the antibody is to act in the brain, a dosage of 50 mg/kg to 100 mg/kg is usually appropriate. Generally, partially human antibodies and fully human antibodies have a longer half-life within the human body than other antibodies. Accordingly, lower dosages and less frequent administration is often possible. Modifications such as lipidation can be used to stabilize antibodies and to enhance uptake and tissue penetration (e.g., into the brain). A method for lipidation of

antibodies is described by Cruikshank et al. ((1997) *J. Acquired Immune Deficiency Syndromes and Human Retrovirology* 14:193).

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (U.S. Patent 5,328,470), or by stereotactic injection (*see, e.g.*, Chen et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

#### V. Uses and Methods of the Invention

The nucleic acid molecules, proteins, protein homologs, and antibodies described herein can be used in one or more of the following methods: a) screening assays; b) detection assays (e.g., chromosomal mapping, tissue typing, forensic biology); c) predictive medicine (e.g., diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenomics); and d) methods of treatment (e.g., therapeutic and prophylactic). For example, polypeptides of the invention can be used for all of the purposes identified herein in portions of the disclosure relating to individual types of protein of the invention (e.g. TANGO 202 proteins, TANGO 234 proteins, TANGO 265 proteins, TANGO 273 proteins, TANGO 286 proteins, TANGO 294 proteins, and INTERCEPT 296 proteins). The isolated nucleic acid molecules of the invention can be used to express proteins (e.g., via a recombinant expression vector in a host cell in gene therapy applications), to detect mRNA (e.g., in a biological sample) or a genetic lesion, and to modulate activity of a polypeptide of the invention. In addition, the polypeptides of the invention can be used to screen drugs or compounds which

modulate activity or expression of a polypeptide of the invention as well as to treat disorders characterized by insufficient or excessive production of a protein of the invention or production of a form of a protein of the invention which has decreased or aberrant activity compared to the wild type protein. In addition, the antibodies of the invention can be used to detect and isolate a protein of the and modulate activity of a protein of the invention.

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

#### A. Screening Assays

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, i.e., candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other drugs) which bind to polypeptide of the invention or have a stimulatory or inhibitory effect on, for example, expression or activity of a polypeptide of the invention.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of the membrane-bound form of a polypeptide of the invention or biologically active portion thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam (1997) *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6909; Erb et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann et al.

(1994). *J. Med. Chem.* 37:2678; Cho et al. (1993) *Science* 261:1303; Carrell et al. (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell et al. (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and Gallop et al. (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds can be presented in solution (e.g., Houghten (1992) *Bio/Techniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (U.S. Patent No. 5,223,409), spores (Patent NOS. 5,571,698; 5,403,484; and 5,223,409), plasmids (Cull et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:1865-1869) or phage (Scott and Smith (1990) *Science* 249:386-390; Devlin (1990) *Science* 249:404-406; Cwirla et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6378-6382; and Felici (1991) *J. Mol. Biol.* 222:301-310).

In one embodiment, an assay is a cell-based assay in which a cell which expresses a membrane-bound form of a polypeptide of the invention, or a biologically active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to the polypeptide determined. The cell, for example, can be a yeast cell or a cell of mammalian origin. Determining the ability of the test compound to bind to the polypeptide can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the polypeptide or biologically active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^3\text{H}$ , either directly or indirectly, and the radioisotope detected by direct counting of radio-emission or by scintillation counting. Alternatively, test compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In a preferred embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of a polypeptide of the invention, or a biologically active portion thereof, on the cell surface with a known compound which binds the polypeptide to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the

polypeptide, wherein determining the ability of the test compound to interact with the polypeptide comprises determining the ability of the test compound to preferentially bind to the polypeptide or a biologically active portion thereof as compared to the known compound.

5 In another embodiment, the assay involves assessment of an activity characteristic of the polypeptide, wherein binding of the test compound with the polypeptide or a biologically active portion thereof alters (i.e. increases or decreases) the activity of the polypeptide.

10 In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of a polypeptide of the invention, or a biologically active portion thereof, on the cell surface with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the polypeptide or biologically active portion thereof. Determining the ability of the test compound to modulate the activity of the polypeptide or a  
15 biologically active portion thereof can be accomplished, for example, by determining the ability of the polypeptide to bind to or interact with a target molecule or to transport molecules across the cytoplasmic membrane.

20 Determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. As used herein, a "target molecule" is a molecule with which a selected polypeptide (e.g., a polypeptide of the invention binds or interacts with in nature, for example, a molecule on the surface of a cell which expresses the selected protein, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell  
25 membrane or a cytoplasmic molecule. A target molecule can be a polypeptide of the invention or some other polypeptide or protein. For example, a target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal (e.g., a signal generated by binding of a compound to a polypeptide of the invention) through the cell membrane and into the cell or a second intercellular

protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with a polypeptide of the invention. Determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., an mRNA, intracellular  $\text{Ca}^{2+}$ , diacylglycerol, IP3, and the like), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g. luciferase), or detecting a cellular response, for example, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the present invention is a cell-free assay comprising contacting a polypeptide of the invention or biologically active portion thereof with a test compound and determining the ability of the test compound to bind to the polypeptide or biologically active portion thereof. Binding of the test compound to the polypeptide can be determined either directly or indirectly as described above. In a preferred embodiment, the assay includes contacting the polypeptide of the invention or biologically active portion thereof with a known compound which binds the polypeptide to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the polypeptide, wherein determining the ability of the test compound to interact with the polypeptide comprises determining the ability of the test compound to preferentially bind to the polypeptide or biologically active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-free assay comprising contacting a polypeptide of the invention or biologically active portion thereof with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the polypeptide or biologically active portion thereof. Determining the ability of the test compound to modulate the activity of the

polypeptide can be accomplished, for example, by determining the ability of the polypeptide to bind to a target molecule by one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of the polypeptide can be accomplished by determining the ability of the polypeptide of the invention to further modulate the target molecule. For example, the catalytic activity, the enzymatic activity, or both, of the target molecule on an appropriate substrate can be determined as previously described.

In yet another embodiment, the cell-free assay comprises contacting a polypeptide of the invention or biologically active portion thereof with a known compound which binds the polypeptide to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the polypeptide, wherein determining the ability of the test compound to interact with the polypeptide comprises determining the ability of the polypeptide to preferentially bind to or modulate the activity of a target molecule.

The cell-free assays of the present invention are amenable to use of both a soluble form or the membrane-bound form of a polypeptide of the invention. In the case of cell-free assays comprising the membrane-bound form of the polypeptide, it can be desirable to utilize a solubilizing agent such that the membrane-bound form of the polypeptide is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-octylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton X-100, Triton X-114, Thesit, Isotridecypoly(ethylene glycol ether)<sub>n</sub>, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl-N,N-dimethyl-3-ammonio-1-propane sulfonate.

In one or more embodiments of the above assay methods of the present invention, it can be desirable to immobilize either the polypeptide of the invention or its target molecule to facilitate separation of complexed from non-complexed forms of

one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to the polypeptide, or interaction of the polypeptide with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione Sepharose beads (Sigma Chemical; St. Louis, MO) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or A polypeptide of the invention, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of the polypeptide of the invention can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the polypeptide of the invention or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with the polypeptide of the invention or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes,

in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the polypeptide of the invention or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the polypeptide of the invention or target molecule.

In another embodiment, modulators of expression of a polypeptide of the invention are identified in a method in which a cell is contacted with a candidate compound and the expression of the selected mRNA or protein (i.e., the mRNA or protein corresponding to a polypeptide or nucleic acid of the invention) in the cell is determined. The level of expression of the selected mRNA or protein in the presence of the candidate compound is compared to the level of expression of the selected mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of expression of the polypeptide of the invention based on this comparison. For example, when expression of the selected mRNA or protein is greater (i.e. statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of the selected mRNA or protein expression. Alternatively, when expression of the selected mRNA or protein is less (i.e. statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of the selected mRNA or protein expression. The level of the selected mRNA or protein expression in the cells can be determined by methods described herein.

In yet another aspect of the invention, a polypeptide of the inventions can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (*see, e.g.*, U.S. Patent No. 5,283,317; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J. Biol. Chem.* 268:12046-12054; Bartel et al. (1993) *Bio/Techniques* 14:920-924; Iwabuchi et al. (1993) *Oncogene* 8:1693-1696; and PCT Publication No. WO 94/10300), to identify other proteins, which bind to or interact with the polypeptide of the invention and modulate activity of the polypeptide of the invention. Such binding proteins are also likely to be involved in the propagation of signals by the polypeptide

of the inventions as, for example, upstream or downstream elements of a signaling pathway involving the polypeptide of the invention.

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

## B. Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

### 1. Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. Accordingly, nucleic acid molecules described herein or fragments thereof, can be used to map the location of the corresponding genes on a chromosome. The mapping of the sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the sequence of a gene of the invention. Computer analysis of the sequence of a gene of the invention can be used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the gene sequences will

yield an amplified fragment. For a review of this technique, see D'Eustachio et al. ((1983) *Science* 220:919-924).

5 PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using a single thermal cycler. Using the nucleic acid sequences of the invention to design oligonucleotide primers, sub-localization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map a gene to its chromosome include *in situ* hybridization (described in Fan et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6223-27), pre-screening 10 with labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries. Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. For a review of this technique, see Verma et al. (Human Chromosomes: A Manual of Basic Techniques (Pergamon Press, New York, 1988)). 15

Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to non-coding regions of the genes actually are preferred for mapping purposes. 20 Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man, available on-line through Johns Hopkins University Welch Medical Library). 25 The relationship between genes and disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, e.g., Egeland et al. (1987) *Nature* 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with a gene of the invention can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

## 2. Tissue Typing

The nucleic acid sequences of the present invention can also be used to identify individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The sequences of the present invention are useful as additional DNA markers for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an alternative technique which determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the nucleic acid sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have

a unique set of such DNA sequences due to allelic differences. The sequences of the present invention can be used to obtain such identification sequences from individuals and from tissue. The nucleic acid sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the non-coding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the non-coding regions, fewer sequences are necessary to differentiate individuals. The non-coding sequences of SEQ ID NO: 1, 9, 17, 25, 33, 45, or 53 can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers which each yield a non-coding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO: 2, 10, 18, 26, 34, 46, or 54 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

If a panel of reagents from the nucleic acid sequences described herein is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

### 3. Use of Partial Gene Sequences in Forensic Biology

DNA-based identification techniques can also be used in forensic biology. Forensic biology is a scientific field employing genetic typing of biological evidence found at a crime scene as a means for positively identifying, for example, a perpetrator of a crime. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, or semen found at a crime scene. The

amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e. another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to non-coding regions are particularly appropriate for this use as greater numbers of polymorphisms occur in the non-coding regions, making it easier to differentiate individuals using this technique. Examples of polynucleotide reagents include the nucleic acid sequences of the invention or portions thereof, e.g., fragments derived from non-coding regions having a length of at least 20 or 30 bases.

The nucleic acid sequences described herein can further be used to provide polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue, e.g., brain tissue. This can be very useful in cases where a forensic pathologist is presented with a tissue of unknown origin. Panels of such probes can be used to identify tissue by species and/or by organ type.

#### C. Predictive Medicine

The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the present invention relates to diagnostic assays for determining expression of a polypeptide or nucleic acid of the invention and/or activity of a polypeptide of the invention, in the context of a biological sample (e.g., blood, serum, cells, tissue) to thereby determine whether an individual is afflicted

with a disease or disorder, or is at risk of developing a disorder, associated with aberrant expression or activity of a polypeptide of the invention. The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with aberrant expression or activity of a polypeptide of the invention. For example, mutations in a gene of the invention can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with aberrant expression or activity of a polypeptide of the invention.

Another aspect of the invention provides methods for expression of a nucleic acid or polypeptide of the invention or activity of a polypeptide of the invention in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (e.g., drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (e.g., the genotype of the individual examined to determine the ability of the individual to respond to a particular agent).

Yet another aspect of the invention pertains to monitoring the influence of agents (e.g., drugs or other compounds) on the expression or activity of a polypeptide of the invention in clinical trials. These and other agents are described in further detail in the following sections.

#### 1. Diagnostic Assays

An exemplary method for detecting the presence or absence of a polypeptide or nucleic acid of the invention in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting a polypeptide or nucleic acid (e.g., mRNA, genomic DNA) of the invention such that the presence of a polypeptide or nucleic acid of the invention is detected in the biological sample. A preferred agent for detecting mRNA or genomic DNA encoding a polypeptide of the invention is a labeled nucleic

acid probe capable of hybridizing to mRNA or genomic DNA encoding a polypeptide of the invention. The nucleic acid probe can be, for example, a full-length cDNA, such as the nucleic acid of SEQ ID NO: 1, 9, 17, 25, 33, 45, 53, 67, or 72, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to a mRNA or genomic DNA encoding a polypeptide of the invention. Other suitable probes for use in the diagnostic assays of the invention are described herein.

A preferred agent for detecting a polypeptide of the invention is an antibody capable of binding to a polypeptide of the invention, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')<sub>2</sub>) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of a polypeptide of the invention include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of a polypeptide of the invention include introducing into a subject a labeled antibody directed against the polypeptide. For

example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting a polypeptide of the invention or mRNA or genomic DNA encoding a polypeptide of the invention, such that the presence of the polypeptide or mRNA or genomic DNA encoding the polypeptide is detected in the biological sample, and comparing the presence of the polypeptide or mRNA or genomic DNA encoding the polypeptide in the control sample with the presence of the polypeptide or mRNA or genomic DNA encoding the polypeptide in the test sample.

The invention also encompasses kits for detecting the presence of a polypeptide or nucleic acid of the invention in a biological sample (a test sample). Such kits can be used to determine if a subject is suffering from or is at increased risk of developing a disorder associated with aberrant expression of a polypeptide of the invention (e.g., one of the disorders described in the section of this disclosure wherein the individual polypeptide of the invention is discussed). For example, the kit can comprise a labeled compound or agent capable of detecting the polypeptide or mRNA encoding the polypeptide in a biological sample and means for determining the amount of the polypeptide or mRNA in the sample (e.g., an antibody which binds the polypeptide or an oligonucleotide probe which binds to DNA or mRNA encoding the polypeptide). Kits can also include instructions for observing that the tested subject is suffering from or is at risk of developing a disorder associated with aberrant expression of the polypeptide if the amount of the polypeptide or mRNA encoding the polypeptide is above or below a normal level.

For antibody-based kits, the kit can comprise, for example: (1) a first antibody (e.g., attached to a solid support) which binds to a polypeptide of the invention; and, optionally, (2) a second, different antibody which binds to either the polypeptide or the first antibody and is conjugated to a detectable agent.

For oligonucleotide-based kits, the kit can comprise, for example: (1) an oligonucleotide, e.g., a detectably labeled oligonucleotide, which hybridizes to a nucleic acid sequence encoding a polypeptide of the invention or (2) a pair of primers useful for amplifying a nucleic acid molecule encoding a polypeptide of the invention. The kit can also comprise, e.g., a buffering agent, a preservative, or a protein stabilizing agent. The kit can also comprise components necessary for detecting the detectable agent (e.g., an enzyme or a substrate). The kit can also contain a control sample or a series of control samples which can be assayed and compared to the test sample contained. Each component of the kit is usually enclosed within an individual container and all of the various containers are within a single package along with instructions for observing whether the tested subject is suffering from or is at risk of developing a disorder associated with aberrant expression of the polypeptide.

## 2. Prognostic Assays

The methods described herein can furthermore be utilized as diagnostic or prognostic assays to identify subjects having or at risk of developing a disease or disorder associated with aberrant expression or activity of a polypeptide of the invention. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with aberrant expression or activity of a polypeptide of the invention (e.g., one of the disorders described in the section of this disclosure wherein the individual polypeptide of the invention is discussed). Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing such a disease or disorder. Thus, the present invention provides a method in which a test sample is obtained from a subject and a polypeptide or nucleic acid (e.g., mRNA,

genomic DNA) of the invention is detected, wherein the presence of the polypeptide or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant expression or activity of the polypeptide. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (e.g., serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant expression or activity of a polypeptide of the invention. For example, such methods can be used to determine whether a subject can be effectively treated with a specific agent or class of agents (e.g., agents of a type which decrease activity of the polypeptide). Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant expression or activity of a polypeptide of the invention in which a test sample is obtained and the polypeptide or nucleic acid encoding the polypeptide is detected (e.g., wherein the presence of the polypeptide or nucleic acid is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant expression or activity of the polypeptide).

The methods of the invention can also be used to detect genetic lesions or mutations in a gene of the invention, thereby determining if a subject with the lesioned gene is at risk for a disorder characterized aberrant expression or activity of a polypeptide of the invention. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion or mutation characterized by at least one of an alteration affecting the integrity of a gene encoding the polypeptide of the invention, or the mis-expression of the gene encoding the polypeptide of the invention. For example, such genetic lesions or mutations can be detected by ascertaining the existence of at least one of: 1) a deletion of one or more nucleotides from the gene; 2) an addition of one or more nucleotides to

the gene; 3) a substitution of one or more nucleotides of the gene; 4) a chromosomal rearrangement of the gene; 5) an alteration in the level of a messenger RNA transcript of the gene; 6) an aberrant modification of the gene, such as of the methylation pattern of the genomic DNA; 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of the gene; 8) a non-wild type level of the protein encoded by the gene; 9) an allelic loss of the gene; and 10) an inappropriate post-translational modification of the protein encoded by the gene. As described herein, there are a large number of assay techniques known in the art which can be used for detecting lesions in a gene.

In certain embodiments, detection of the lesion involves the use of a probe/primer in a polymerase chain reaction (PCR) (*see, e.g.*, U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (*see, e.g.*, Landegran et al. (1988) *Science* 241:1077-1080; and Nakazawa et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:360-364), the latter of which can be particularly useful for detecting point mutations in a gene (*see, e.g.*, Abravaya et al. (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to the selected gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. PCR and/or LCR can be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self-sustained sequence replication (Guatelli et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh, et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi et al. (1988) *Bio/Technology* 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified

molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

5 In an alternative embodiment, mutations in a selected gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, (optionally) amplified, digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of  
10 sequence specific ribozymes (*see, e.g.*, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

15 In other embodiments, genetic mutations can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin et al. (1996) *Human Mutation* 7:244-255; Kozal et al. (1996) *Nature Medicine* 2:753-759). For example, genetic mutations can be identified in two-dimensional arrays containing light-generated DNA probes as described in Cronin et al., *supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear  
20 arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other  
25 complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the selected gene and detect mutations by comparing the sequence of the sample nucleic acids with the

corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert ((1977) *Proc. Natl. Acad. Sci. USA* 74:560) or Sanger ((1977) *Proc. Natl. Acad. Sci. USA* 74:5463). It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Bio/Techniques* 19:448), including sequencing by mass spectrometry (*see, e.g.*, PCT Publication No. WO 94/16101; Cohen et al. (1996) *Adv. Chromatogr.* 36:127-162; and Griffin et al. (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in a selected gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers et al. (1985) *Science* 230:1242). In general, the technique of mismatch cleavage entails providing heteroduplexes formed by hybridizing (labeled) RNA or DNA containing the wild-type sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to base pair mismatches between the control and sample strands. RNA/DNA duplexes can be treated with RNASE to digest mismatched regions, and DNA/DNA hybrids can be treated with S1 nuclease to digest mismatched regions.

In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. *See, e.g.*, Cotton et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:4397; Saleeba et al. (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called DNA mismatch repair enzymes) in defined systems for detecting and mapping

point mutations in cDNAs obtained from samples of cells. For example, the mutY enzyme of E. coli cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu et al. (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on a selected sequence, e.g., a wild-type sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, e.g., U.S. Patent No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in genes. For example, single strand conformation polymorphism (SSCP) can be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:2766; see also Cotton (1993) *Mutat. Res.* 285:125-144; Hayashi (1992) *Genet. Anal. Tech. Appl.* 9:73-79). Single-stranded DNA fragments of sample and control nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, and the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments can be labeled or detected with labeled probes. The sensitivity of the assay can be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen et al. (1991) *Trends Genet.* 7:5).

In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers et al. (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a 'GC clamp' of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature

gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) *Biophys. Chem.* 265:12753).

5 Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers can be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions which permit hybridization only if a perfect match is found (Saiki et al. (1986) *Nature* 324:163); Saiki et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:6230). Such  
10 allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification can be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification can carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization; Gibbs et al. (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme  
15 3' end of one primer where, under appropriate conditions, mismatching can prevent or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition, it can be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini et al. (1992) *Mol. Cell Probes* 6:1). Amplification  
20 can also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci. USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a  
25 known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein can be performed, for example, using pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which can be conveniently used, e.g., in clinical settings to diagnose

patients exhibiting symptoms or family history of a disease or illness involving a gene encoding a polypeptide of the invention. Furthermore, any cell type or tissue, preferably peripheral blood leukocytes, in which the polypeptide of the invention is expressed can be utilized in the prognostic assays described herein.

5

### 3. Pharmacogenomics

Agents, or modulators which have a stimulatory or inhibitory effect on activity or expression of a polypeptide of the invention as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders associated with aberrant activity of the polypeptide. In conjunction with such treatment, the pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of the individual may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (e.g., drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to determine appropriate dosages and therapeutic regimens. Accordingly, the activity of a polypeptide of the invention, expression of a nucleic acid of the invention, or mutation content of a gene of the invention in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. *See, e.g., Linder (1997) Clin. Chem. 43(2):254-266.* In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body are referred to as "altered drug action." Genetic conditions transmitted as single factors altering the way the body acts on drugs are referred to as "altered drug metabolism". These pharmacogenetic

conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical complication is hemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (e.g., N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, a PM will show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Thus, the activity of a polypeptide of the invention, expression of a nucleic acid encoding the polypeptide, or mutation content of a gene encoding the polypeptide in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection,

can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a modulator of activity or expression of the polypeptide, such as a modulator identified by one of the exemplary screening assays described herein.

5

#### 4. Monitoring of Effects During Clinical Trials

10

Monitoring the influence of agents (e.g., drug compounds) on the expression or activity of a polypeptide of the invention (e.g., the ability to modulate aberrant cell proliferation chemotaxis, and/or differentiation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent, as determined by a screening assay as described herein, to increase gene expression, protein levels, or protein activity, can be monitored in clinical trials of subjects exhibiting decreased gene expression, protein levels, or protein activity. Alternatively, the effectiveness of an agent, as determined by a screening assay, to decrease gene expression, protein levels or protein activity, can be monitored in clinical trials of subjects exhibiting increased gene expression, protein levels, or protein activity. In such clinical trials, expression or activity of a polypeptide of the invention and preferably, that of other polypeptide that have been implicated in for example, a cellular proliferation disorder, can be used as a marker of the immune responsiveness of a particular cell.

15

20

25

For example, and not by way of limitation, genes, including those of the invention, that are modulated in cells by treatment with an agent (e.g., compound, drug or small molecule) which modulates activity or expression of a polypeptide of the invention (e.g., as identified in a screening assay described herein) can be identified. Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of a gene of the invention and other genes implicated in the disorder. The levels of gene expression (i.e., a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount

of protein produced, by one of the methods as described herein, or by measuring the levels of activity of a gene of the invention or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state can be determined before, and at various points during, treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of the polypeptide or nucleic acid of the invention in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level the of the polypeptide or nucleic acid of the invention in the post-administration samples; (v) comparing the level of the polypeptide or nucleic acid of the invention in the pre-administration sample with the level of the polypeptide or nucleic acid of the invention in the post-administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent can be desirable to increase the expression or activity of the polypeptide to higher levels than detected, i.e., to increase the effectiveness of the agent. Alternatively, decreased administration of the agent can be desirable to decrease expression or activity of the polypeptide to lower levels than detected, i.e., to decrease the effectiveness of the agent.

### C. Methods of Treatment

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant expression or activity of a polypeptide of the invention and/or in which the polypeptide of the invention is involved. Disorders

characterized by aberrant expression or activity of the polypeptides of the invention are described elsewhere in this disclosure.

1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant expression or activity of a polypeptide of the invention, by administering to the subject an agent which modulates expression or at least one activity of the polypeptide. Subjects at risk for a disease which is caused or contributed to by aberrant expression or activity of a polypeptide of the invention can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of aberrancy, for example, an agonist or antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein.

2. Therapeutic Methods

Another aspect of the invention pertains to methods of modulating expression or activity of a polypeptide of the invention for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of the polypeptide. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more of the biological activities of the polypeptide. Examples of such stimulatory agents include the active polypeptide of the invention and a nucleic acid molecule encoding the polypeptide of the invention that has been introduced into the cell. In another embodiment, the agent inhibits one or more of the biological activities of the

polypeptide of the invention. Examples of such inhibitory agents include antisense nucleic acid molecules and antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of a polypeptide of the invention. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., up-regulates or down-regulates) expression or activity. In another embodiment, the method involves administering a polypeptide of the invention or a nucleic acid molecule of the invention as therapy to compensate for reduced or aberrant expression or activity of the polypeptide.

Stimulation of activity is desirable in situations in which activity or expression is abnormally low or down-regulated and/or in which increased activity is likely to have a beneficial effect, e.g., in wound healing. Conversely, inhibition of activity is desirable in situations in which activity or expression is abnormally high or up-regulated and/or in which decreased activity is likely to have a beneficial effect.

The contents of all references, patents, and published patent applications cited throughout this application are hereby incorporated by reference.

#### Deposits of Clones

Each of these deposits was made merely as a convenience to those of skill in the art. These deposits are not an admission that a deposit is required under 35 U.S.C. §112.

Clone EpT202, encoding human TANGO 202 was deposited with the American Type Culture Collection (ATCC, 10801 University Boulevard, Manassas, VA 20110-2209) on April 21, 1999 and was assigned Accession Number 207219. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

Clone EpTm202, encoding murine TANGO 202 was deposited with

ATCC on April 21, 1999 and was assigned (composite) Accession Number 207221. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

5                   Clone EpT234, encoding human TANGO 234 was deposited with ATCC on April 2, 1999 and was assigned Accession Number 207184. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

10                   Clone EpT265, encoding human TANGO 265 was deposited with ATCC on April 28, 1999 and was assigned Accession Number 207228. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

15                   Clone EpT273, encoding human TANGO 273 was deposited with ATCC on April 2, 1999 and was assigned Accession Number 207185. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

20                   Clone EpTm273, encoding murine TANGO 273 was deposited with ATCC on April 2, 1999 and was assigned (composite) Accession Number 207221. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

25                   Clone EpT286, encoding human TANGO 286 was deposited with ATCC on April 20, 1999 and was assigned (composite) Accession Number 207220. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

                  Clone EpT294, encoding human TANGO 294 was deposited with ATCC on April 20, 1999 and was assigned (composite) Accession Number 207220. This deposit will be maintained under the terms of the Budapest Treaty on the

International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

Clone EpT296, encoding human INTERCEPT 296 was deposited with ATCC on April 20, 1999 and was assigned (composite) Accession Number 207220.

This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

Clones containing cDNA molecules encoding human TANGO 286, human TANGO 294, and INTERCEPT 296 were deposited with ATCC on April 21, 1999 as Accession Number 207220, as part of a composite deposit representing a mixture of five strains, each carrying one recombinant plasmid harboring a particular cDNA clone.

To distinguish the strains and isolate a strain harboring a particular cDNA clone, an aliquot of the mixture is streaked out to single colonies on nutrient medium (e.g., LB plates) supplemented with 100 mg/ml ampicillin, single colonies are grown, and then plasmid DNA is extracted using a standard miniprep procedure. Next, a sample of the DNA miniprep is digested with a combination of the restriction enzymes *Sall*, *NotI*, and *DraII* and the resulting products are resolved on a 0.8% agarose gel using standard DNA electrophoresis conditions. This digestion procedure liberates fragments as follows:

1. human TANGO 286 (clone EpT286): 1.85 kB and .1 kB (human TANGO 286 has a *DraII* cut site at about bp 1856).

2. human TANGO 294 (clone EpT294): 1.4 kB and .6 kB (human TANGO 294 has a *DraII* cut site at about bp 1447).

3. human INTERCEPT 296 (clone EpT296): .4 kB, 1.6 kB, and .1 kB (human INTERCEPT 296 has *DraII* cut sites at about bp 410 and at about bp 1933).

The identity of the strains can be inferred from the fragments liberated.

Clones containing cDNA molecules encoding mouse TANGO 202 and mouse TANGO 273 were deposited with ATCC on April 21, 1999 and were assigned Accession Number 207221, as part of a composite deposit representing a mixture of five strains, each carrying one recombinant plasmid harboring a particular cDNA clone.

To distinguish the strains and isolate a strain harboring a particular cDNA clone, an aliquot of the mixture is streaked out to single colonies on nutrient medium (e.g., LB plates) supplemented with 100 mg/ml ampicillin, single colonies are grown, and then plasmid DNA is extracted using a standard miniprep procedure. Next, a sample of the DNA miniprep is digested with a combination of the restriction enzymes *Sal* I, *Not* I, and *Apa* I, and the resultant products are resolved on a 0.8% agarose gel using standard DNA electrophoresis conditions. This digestion procedure liberates fragments as follows:

1. mouse TANGO 202 (clone EpTm202): 3.5 kB and 1.4 kB (mouse TANGO 202 has a *Apa* I cut site at about bp 3519).

2. mouse TANGO 273 (clone EpTm273): .3 kB and 2.6 kB (mouse TANGO 273 has a *Apa* I cut site at about bp 298).

The identity of the strains can be inferred from the fragments liberated.

Human TANGO 202, human TANGO 234, human TANGO 265, and human TANGO 273 were each deposited as single deposits. Their clone names, deposit dates, and accession numbers are as follows:

1. human TANGO 202: clone EpT202 was deposited with ATCC on April 21, 1999, and was assigned Accession Number 207219.

2. human TANGO 234: clone EpT234 was deposited with ATCC on April 2, 1999, and was assigned Accession Number 207184.

3. human TANGO 265: clone EpT265 was deposited with ATCC on April 28, 1999, and was assigned Accession Number 207228.

4. human TANGO 273: clone EpT273 was deposited with ATCC on April 2, 1999, and was assigned Accession Number 207185.

### Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed is:

1                   1. An isolated nucleic acid molecule selected from the group consisting  
2 of:

3                   a) a nucleic acid molecule having a nucleotide sequence which is at least 40%  
4 identical to the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34,  
5 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone  
6 deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a  
7 complement thereof;

8                   b) a nucleic acid molecule comprising at least 15 nucleotide residues and  
9 having a nucleotide sequence identical to at least 15 consecutive nucleotide residues of  
10 SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the  
11 nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184,  
12 207228, 207185, 207220, or 207221, or a complement thereof;

13                   c) a nucleic acid molecule comprising at least 15 nucleotide residues and  
14 having a nucleotide sequence identical to at least 15 consecutive nucleotide residues of  
15 SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the  
16 nucleotide sequence of a cDNA of a clone deposited as ATCC 207219, 207184,  
17 207228, 207185, 207220, or 207221, or a complement thereof;

18                   d) a nucleic acid molecule which encodes a polypeptide comprising the amino  
19 acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or  
20 the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219,  
21 207184, 207228, 207185, 207220, or 207221, or a complement thereof;

22                   e) a nucleic acid molecule which encodes a fragment of a polypeptide  
23 comprising the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44,  
24 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone  
25 deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, wherein the  
26 fragment comprises at least 8 consecutive amino acid residues of SEQ ID NO: 3-8, 11-  
27 16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a  
28 cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or  
29 207221; and

30 f) a nucleic acid molecule which encodes a naturally occurring allelic variant of  
31 a polypeptide comprising the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24,  
32 27-32, 35-44, 47-52, 55-66, 69, 74, wherein the nucleic acid molecule hybridizes to a  
33 nucleic acid molecule consisting of the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10,  
34 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a  
35 cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or  
36 207221, or a complement thereof under stringent conditions.

1 2. The isolated nucleic acid molecule of claim 1, which is selected from  
2 the group consisting of:

3 a) a nucleic acid having the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17,  
4 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA  
5 of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221,  
6 or a complement thereof; and

7 b) a nucleic acid molecule which encodes a polypeptide having the amino acid  
8 sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the  
9 amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219,  
10 207184, 207228, 207185, 207220, or 207221, or a complement thereof.

1 3. The nucleic acid molecule of claim 1, further comprising vector  
2 nucleic acid sequences.

1 4. The nucleic acid molecule of claim 1 further comprising nucleic acid  
2 sequences encoding a heterologous polypeptide.

1 5. A host cell which contains the nucleic acid molecule of claim 1.

1 6. The host cell of claim 5 which is a mammalian host cell.

1 7. A non-human mammalian host cell containing the nucleic acid  
2 molecule of claim 1.

1 8. An isolated polypeptide selected from the group consisting of:

2 a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID  
3 NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence  
4 encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185,  
5 207220, or 207221, wherein the fragment comprises at least 8 contiguous amino acids  
6 of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino  
7 acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184,  
8 207228, 207185, 207220, or 207221;

9 b) a naturally occurring allelic variant of a polypeptide comprising the amino  
10 acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or  
11 the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219,  
12 207184, 207228, 207185, 207220, or 207221, wherein the polypeptide is encoded by a  
13 nucleic acid molecule which hybridizes to a nucleic acid molecule consisting of the  
14 nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54,  
15 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC  
16 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof under  
17 stringent conditions; and

18 c) a polypeptide which is encoded by a nucleic acid molecule comprising a  
19 nucleotide sequence which is at least 40% identical to a nucleic acid consisting of the  
20 nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54,  
21 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a clone deposited as ATCC  
22 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof.

1 9. The isolated polypeptide of claim 8 having the amino acid sequence  
2 of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino

acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof.

10. The polypeptide of claim 8, wherein the amino acid sequence of the polypeptide further comprises heterologous amino acid residues.

11. An antibody which selectively binds to the polypeptide of claim 8.

12. A method for producing a polypeptide selected from the group consisting of:

a) a polypeptide comprising the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof;

b) a polypeptide comprising a fragment of the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof, wherein the fragment comprises at least 8 contiguous amino acids of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof; and

c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO: 3-8, 11-16, 19-24, 27-32, 35-44, 47-52, 55-66, 69, 74, or the amino acid sequence encoded by a cDNA of a clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule consisting of the nucleotide sequence of SEQ ID NO: 1, 2, 9, 10, 17, 18, 25, 26, 33, 34, 45, 46, 53, 54, 67, 68, 72, 73, or the nucleotide sequence of a cDNA of a

clone deposited as ATCC 207219, 207184, 207228, 207185, 207220, or 207221, or a complement thereof under stringent conditions;

the method comprising culturing the host cell of claim 5 under conditions in which the nucleic acid molecule is expressed.

13. A method for detecting the presence of a polypeptide of claim 8 in a sample, comprising:

a) contacting the sample with a compound which selectively binds to a polypeptide of claim 8; and

b) determining whether the compound binds to the polypeptide in the sample.

14. The method of claim 13, wherein the compound which binds to the polypeptide is an antibody.

15. A kit comprising a compound which selectively binds to a polypeptide of claim 8 and instructions for use.

16. A method for detecting the presence of a nucleic acid molecule of claim 1 in a sample, comprising the steps of:

a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and

b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample.

17. The method of claim 16, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

18. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 and instructions for use.

3 19. A method for identifying a compound which binds to a polypeptide  
4 of claim 8 comprising the steps of:

5 a) contacting a polypeptide, or a cell expressing a polypeptide of claim 8 with a  
6 test compound; and

7 b) determining whether the polypeptide binds to the test compound.

1 20. The method of claim 19, wherein the binding of the test compound  
2 to the polypeptide is detected by a method selected from the group consisting of:

3 a) detection of binding by direct detecting of test compound/polypeptide  
4 binding;

5 b) detection of binding using a competition binding assay;

6 c) detection of binding using an assay for an activity characteristic of the  
7 polypeptide.

1 21. A method for modulating the activity of a polypeptide of claim 8  
2 comprising contacting a polypeptide or a cell expressing a polypeptide of claim 8 with  
3 a compound which binds to the polypeptide in a sufficient concentration to modulate  
4 the activity of the polypeptide.

1 22. A method for identifying a compound which modulates the activity  
2 of a polypeptide of claim 8, comprising:

3 a) contacting a polypeptide of claim 8 with a test compound; and

4 b) determining the effect of the test compound on the activity of the  
5 polypeptide to thereby identify a compound which modulates the activity of the  
6 polypeptide.

1 23. An antibody substance which selectively binds to the polypeptide of  
2 claim 8, wherein the antibody substance is made by providing the polypeptide to an

3 immunocompetent vertebrate and thereafter harvesting blood or serum from the  
4 vertebrate.

RECEIVED

NOVEL GENES ENCODING PROTEINS HAVING  
DIAGNOSTIC, PREVENTIVE, THERAPEUTIC, AND OTHER USES

Abstract of the Disclosure

The invention provides isolated nucleic acids encoding a variety of proteins having diagnostic, preventive, therapeutic, and other uses. These nucleic and proteins are useful for diagnosis, prevention, and therapy of a number of human and other animal disorders. The invention also provides antisense nucleic acid molecules, expression vectors containing the nucleic acid molecules of the invention, host cells into which the expression vectors have been introduced, and non-human transgenic animals in which a nucleic acid molecule of the invention has been introduced or disrupted. The invention still further provides isolated polypeptides, fusion polypeptides, antigenic peptides and antibodies. Diagnostic, screening, and therapeutic methods utilizing compositions of the invention are also provided. The nucleic acids and polypeptides of the present invention are useful as modulating agents in regulating a variety of cellular processes.

Patent for the Invention

Fig. 1A

M A P P A A R L A L L																			11
GTCGACCCACGCGTCCGCCCCACGCGTCCGGCCC ATG GCG CCG CCC GCC GCC CGC CTC GCC CTG CTC																			66
S A A A L T L A A R P A P S P G L G P G	31																		
TCC GCC GCG GCG CTC ACG CTG GCG GCC CGG CCC GCG CCT AGC CCC GGC CTC GGC CCC GGA	126																		
P E C F T A N G A D Y R G T Q N W T A L	51																		
CCC GAG TGT TTC ACA GCC AAT GGT GCG GAT TAT AGG GGA ACA CAG AAC TGG ACA GCA CTA	186																		
Q G G K P C L F W N E T F Q H P Y N T L	71																		
CAA GGC GGG AAG CCA TGT CTG TTT TGG AAC GAG ACT TTC CAG CAT CCA TAC AAC ACT CTG	246																		
K Y P N G E G G L G E H N Y C R N P D G	91																		
AAA TAC CCC AAC GGG GAG GGG GGC CTG GGT GAG CAC AAC TAT TGC AGA AAT CCA GAT GGA	306																		
D V S P W C Y V A E H E D G V Y W K Y C	111																		
GAC GTG AGC CCC TGG TGC TAT GTG GCA GAG CAC GAG GAT GGT GTC TAC TGG AAG TAC TGT	366																		
E I P A C Q M P G N L G C Y K D H G N P	131																		
GAG ATA CCT GCT TGC CAG ATG CCT GGA AAC CTT GGC TGC TAC AAG GAT CAT GGA AAC CCA	426																		
P P L T G T S K T S N K L T I Q T C I S	151																		
CCT CCT CTA ACT GGC ACC AGT AAA ACG TCC AAC AAA CTC ACC ATA CAA ACT TGC ATC AGT	486																		
F C R S Q R F K F A G M E S G Y A C F C	171																		
TTT TGT CGG AGT CAG AGG TTC AAG TTT GCT GGG ATG GAG TCA GGC TAT GCT TGC TTC TGT	546																		
G N N P D Y W K Y G E A A S T E C N S V	191																		
GGA AAC AAT CCT GAT TAC TGG AAG TAC GGG GAG GCA GCC AGT ACC GAA TGC AAC AGC GTC	606																		
C F G D H T Q P C G G D G R I I L F D T	211																		
TGC TTC GGG GAT CAC ACC CAA CCC TGT GGT GGC GAT GGC AGG ATC ATC CTC TTT GAT ACT	666																		
L V G A C G G N Y S A M S S V V Y S P D	231																		
CTC GTG GGC GCC TGC GGT GGG AAC TAC TCA GCC ATG TCT TCT GTG GTC TAT TCC CCT GAC	726																		
F P D T Y A T G R V C Y W T I R V P G A	251																		
TTC CCC GAC ACC TAT GCC ACG GGG AGG GTC TGC TAC TGG ACC ATC CGG GTT CCG GGG GCC	786																		
S H I H F S F P L F D I R D S A D M V E	271																		
TCC CAC ATC CAC TTC AGC TTC CCC CTA TTT GAC ATC AGG GAC TCG GCG GAC ATG GTG GAG	846																		
L L D G Y T H R V L A R F H G R S R P P	291																		
CTT CTG GAT GGC TAC ACC CAC CGT GTC CTA GCC CGC TTC CAC GGG AGG AGC CGC CCA CCT	906																		
L S F N V S L D F V I L Y F F S D R I N	311																		
CTG TCC TTC AAC GTC TCT CTG GAC TTC GTC ATC TTG TAT TTC TTC TCT GAT CGC ATC AAT	966																		
Q A Q G F A V L Y Q A V K E E L P Q E R	331																		
CAG GCC CAG GGA TTT GCT GTT TTA TAC CAA GCC GTC AAG GAA GAA CTG CCA CAG GAG AGG	1026																		
P A V N Q T V A E V I T E Q A N L S V S	351																		
CCC GCT GTC AAC CAG ACG GTG GCC GAG GTG ATC ACG GAG CAG GCC AAC CTC AGT GTC AGC	1086																		
A A R S S K V L Y V I T T S P S H P P Q	371																		
GCT GCC CGG TCC TCC AAA GTC CTC TAT GTC ATC ACC ACC AGC CCC AGC CAC CCA CCT CAG	1146																		
T V P G S N S W A P P M G A G S H R V E	391																		
ACT GTC CCA GGT AGC AAT TCC TGG GCG CCA CCC ATG GGG GCT GGA AGC CAC AGA GTT GAA	1206																		
G W T V Y G L A T L L I L T V T A I V A	411																		

AAACCCCACTGTGCCTAGGACTTGAGGTCCCTCTTTGAGCTCAAGGCTGCCGTGGTCAACCTCTCCTGTGGTTCTTCTC	1540
TGACAGACTCTTCCCTCCTCTCCCTCTGCCTCGGCCCTCTTCGGGGAAACCTCCTCCTACAGACTAGGAAGAGGCACC	1619
TGCTGCCAGGGCAGGCAGAGCCTGGATTCTCTCCTGCTT	1657

Fig. 1B

Fig. 1C

GTCGACCCACGCGTCCGCCCCGGCTCCCGGTGCTGCCCCCTCTGCCCCGGGCCGCGCCCGGGGTCCCGCACTGACGGCC 79  
 M A P P A A R L A L L S A A A L T L A 19  
 C ATG GCG CCG CCC GCC GCC CGT CTC GCG CTG CTC TCC GCC GCT GCG CTC ACT CTG GCG 137  
 A R P A P G P R S G P E C F T A N G A D 39  
 GCC CGG CCC GCG CCC GGT CCC CGC TCC GGC CCC GAG TGC TTC ACA GCC AAC GGT GCA GAT 197  
 Y R G T Q S W T A L Q G G K P C L F W N 59  
 TAC AGG GGA ACA CAG AGC TGG ACA GCG CTG CAA GGT GGG AAG CCA TGT CTG TTC TGG AAC 257  
 E T F Q H P Y N T L K Y P N G E G G L G 79  
 GAG ACT TTC CAG CAT CCG TAC AAC ACG CTG AAG TAC CCC AAC GGG GAA GGA GGA CTG GGC 317  
 E H N Y C R N P D G D V S P W C Y V A E 99  
 GAG CAC AAT TAT TGC AGA AAT CCA GAT GGA GAC GTG AGC CCT TGG TGC TAC GTG GCC GAG 377  
 H E D G V Y W K Y C E I P A C Q M P G N 119  
 CAT GAG GAC GGA GTC TAC TGG AAG TAC TGT GAA ATT CCT GCC TGC CAG ATG CCT GGA AAC 437  
 F G C Y K D H G N P P P L T G T S K T S 139  
 CTT GGC TGC TAC AAG GAT CAT GGA AAC CCA CCT CCT CTC ACG GGC ACC AGT AAA ACC TCT 497  
 M K L T I Q T C I S F C R S Q R F K F A 159  
 AAC AAG CTC ACC ATA CAA ACC TGT ATC AGC TTC TGT CGG AGT CAG AGA TTC AAG TTT GCT 557  
 G M E S G Y A C F C G N N P D Y W K H G 179  
 GGG ATG GAG TCA GGC TAT GCC TGC TTC TGT GGG AAC AAT CCT GAC TAC TGG AAG CAC GGG 617  
 E A A S T E C N S V C F G D H T Q P C G 199  
 GAG GCG GCC AGC ACC GAG TGC AAT AGT GTC TGC TTC GGG GAC CAC ACG CAG CCC TGC GGT 677  
 G D G R I I L F D T L V G A C G G N Y S 219  
 GGG GAC GGC AGG ATT ATC CTC TTT GAC ACT CTC GTG GGC GCC TGC GGT GGG AAC TAC TCA 737  
 A M A A V V Y S P D F P D T Y A T G R V 239  
 GCC ATG GCA GCC GTG GTG TAC TCC CCT GAC TTC CCT GAC ACC TAC GCC ACT GGC AGA GTC 797  
 C Y W T I R V P G A S R I H F N F T L F 259  
 TGC TAC TGG ACC ATC CGG GTT CCA GGA GCC TCT CGC ATC CAT TTC AAC TTC ACC CTG TTT 857  
 D I R D S A D M -V E L L D G Y T H R V L 279  
 GAT ATC AGG GAC TCT GCA GAC ATG GTG GAG CTG CTG GAC GGC TAC ACC CAC CGC GTC CTG 917  
 V R L S G R S R P P L S F N V S L D F V 299  
 GTC CGG CTC AGT GGG AGG AGC CGC CCG CCT CTG TCT TTC AAT GTC TCT CTG GAT TTT GTC 977  
 I L Y F F S D R I N Q A Q G F A V L Y Q 319  
 ATT TTG TAT TTC TTC TCT GAT CGC ATC AAT CAG GCC CAG GGA TTT GCT GTG TTG TAC CAA 1037  
 A T K E E P P Q E R P A V N Q T L A E V 339  
 GCC ACC AAG GAG GAA CCG CCA CAG GAG AGA CCT GCT GTC AAC CAG ACC CTG GCA GAG GTG 1097  
 I T E Q A N L S V S A A H S S K V L Y V 359

ATC ACC GAG CAA GCC AAC CTC AGT GTC AGC GCT GCC CAC TCC TCC AAA GTC CTC TAT GTC 1157  
 I T P S P S H P P Q T A Q V A I P G H R 379  
 ATC ACC CCC AGC CCC AGC CAC CCT CCG CAG ACT GCC CAG GTA GCC ATT CCT GGG CAC CGT 1217  
 Q L G P T A T E W K D G L C T A W R P S 399  
 CAG TTG GGG CCA ACA GCC ACA GAG TGG AAG GAT GGA CTG TGT ACG GCC TGG CGA CCC TCC 1277  
 S S S Q S Q Q L S Q R F F C M S H L N L 419  
 TCA TCC TCA CAG TCA CAG CAG TTG TCG CAA AGA TTC TTC TGC ATG TCA CAT TTA AAT CTC 1337  
 I E S L H Q E T L G T V V S L G L L E I 439  
 ATC GAG TCC CTG CAT CAG GAG ACC TTA GGG ACT GTC GTC AGC CTG GGG CTT CTG GAG ATA 1397  
 S G P F S M N L P L Q S P S L R R S S R 459  
 TCT GGA CCA TTT TCT ATG AAC CTT CCA CTA CAA TCT CCA TCT TTA AGA AGA AGC TCA AGG 1457  
 V R V N K M T A I P S \* 471  
 GTC AGA GTC AAC AAG ATG ACC GCA ATC CCC TCG TGA 1493  
 GTGACTGAAGCCACGCCTGCATGAGAGGCTCCGCTCCAAGCTCGAGTTTGCTCCCCTGAGTTCTCCTCTGATGAGTTC 1572  
 CCGGCCTTCCCATTCACCACCATCTCTTTTGGGAGCACCTGCTTTAGAGGCAGCCAGCCTGGGATCCTCCATCACAT 1651  
 GTACCAGCCTGGCTGCTCTGCTGGGGATGGTAAGACAGGCCCAGGCTGACAGGACACAGCTGGACCTGACTCCAGAAGA 1730  
 CTCTTGGGTGGTGGGGAGGTATAGTGTAGGATGAGTTTTCTTGCTTCTTCTCTGTTTTGTCCACATACAGATCGGTTTT 1809  
 CCGTGTCTTTACAGTTTGC AATAGAGCCAGACTGAAAGAACTGTCAGGTTTTCTAGGCTGGCCTGGTTCCCCACTAAGA 1888  
 GTGGCATTTGGCGCCCTAGAGGCCAGAGGCCAGTGTAGGCTTGGAGCTTTCTCTGCTGCCAACTACCATGTGTATCT 1967  
 AGTCCGAGGGGACTGAGAGCAGGGCCACACCAGATGTCATCTTTCTAGAGGGTTCTTTTTTAGTACCCACTGACCAATGG 2046  
 GGCAAGCCTGAGGATTGGTCCATCTGTTTGTCCATGGAACAGACACAGTGAACCTTCTGGATACTAGACTTAAGTAGCC 2125  
 TAGCCCTCAAGTAGTTGCCAATCCTGTGGAATCAGAATTCAGCCTGTCTTCTGTCTCAGCCCAAGCCTGTAGCCTAG 2204  
 AGCTGGGGCTGTAGCCTAGAGCTGGGGCTGTAGCCTAGAGCTGGGGCTGTAGCACAGAGCTGGGGCTGTAGCCTAGAGC 2283  
 TGGGGCTGTAGCACAGAGCTGGGGCTGTAGCCTAGAGCTGGGGCTGTAGCACAGAGCTGGGGCTGTAGCACAGAGCTGG 2362  
 GGCTGTAGCCTAGAGCTGGGGCTGTAGCACAGAGCTGGGGCTGTAACCTCAGCGATCAAGAGCTTGCTTTGTATACATCG 2441  
 GACCCTAGGTTCTATCCCAGCACTATCAGAAGGTGGGAGAGAAAAAGACTGCACCATAGCATGCGGGCAGCATCTGTGG 2520  
 TTCTTACGTGAGGTGTCATCATTTTAAAAGCAGATCAAACTACCGCGAGTTTTGTCCTTTGTCCCTTATCATGGGAGC 2599  
 AGAGTAGGAGTAAGGGCTCTGGTCTTGCTCATTTGTCCCCCAGACAGGGAGGCAGGAAAAGGTCAGGCTTGGGAAGTGA 2678  
 GATCCTCCCAGGAAAAGCTGCAAGATTGAGAGACCCAGCTGCAGTTGGGAGAGGAAGGGCCATCCCCGACTGAGAAGTC 2757  
 CTGCAGTCTGGAAGTGGCCTTTGTCTCAGCAGCAGCTGTGCCCTGAAGGTAGACCTTGGTCACTCTCCTGCCAGCCCTTGA 2836  
 GCCTCTGCTCTCCTGGGTACCCTCCTGGAACACCATGCTAACCTTCCCCGAGTCTCTCAGTCACTGCCATTGAGGCCTC 2915  
 TCCTCTAGCTGCTGCTCCCCAGGACTGTCTGGGGCCATCTGGGGATCAGGGAGAGGCAGCAGGAGTACTGACGAGGCAG 2994

TGACCTGAGCTGATGAGTCAACCAGAGGACACCAGAGTCTACAGTGGGCTGGCTGCTGGCTCAGCTCCTATGGGAGGCC 3073  
 TACAGGGGTACTAAGCTAGGGGGTCATCATCTCATTTGATCTGGGAAAGGCTACAGGCTCCTGGATGTGAAGACAGGCC 3152  
 CACTACATAAGAAGACCACTGGAAATAGACTGACAGGAGCAGGTTCCACTCTAGGCTGTCCATAGCGTTTGCAGGACTC 3231  
 CCCTGAGACCAAGTGTGTGAGTCACAGAGTGCCATGTGCGTAGTGCCATAAAGGATATGGGTTCCTTAACCAGGGAAGGCTC 3310  
 ATAGCAGGCCAGGACATTTTTTTCAGCTCAGAGCACTGGCCCCAGGCTTCCTCTAAGCCACCACTCACCTGTCTCTTCCT 3389  
 ATCTCGGACACAGGAAGCAAGCCCCAGTGTGGTGGCAGCTGCGGCTCAGCATTTGGTGTCCCCAGGAAGGGCGGTGGATG 3468  
 TGCCCCAGCTCCTTTTGTCTGTGGGCTGGCACAGCCCAACACTGCAGGGCCCCACCTTCTCTCTTGGGGGGTAGGGACAC 3547  
 ATAAGGAAAATAACCCACCTCCAACAACAGCAGAGGACAGTGGGAAGGAAGGGCTGTAAATCACCCAGGCCAGACCTC 3626  
 CAGAAATGACAGGCACAGTCTGTTAGAACCTGTAGGCAGCCAGTCACAGAGGGCCTTTGTGCTGGTAACACCCCTGCCTG 3705  
 GAGCATAGGGGTAAGCCGAGGGAGAAGAGCAGCCCTCAGAGACATCAGCTAAAAACATAGGTGCCCTATGTCCCTCCCT 3784  
 TCCTGTCACACTGCTTACAAAGCAGAGACAGAGTAGGAAAGAGGTCTTCATCCTCTCCACATCAGCAAGGATAGGGCT 3863  
 GCGGCTGCCTAAAGTGAGCAAGGAGAACAGAGCTCTGGACTTCTCTAAATGTGGGCTCTGGCTTCAGACTCCTCAGCCA 3942  
 AAAGCTCTTGAAGATCAAAGCTCTGGCGGGTACAGCTGTCTGGCCTGTGGGCCAGCCCATGGGATGTGCCTGGGCCAG 4021  
 GTGCCACCCACGGCTCACTGTCTATCCCAGGAGGGACCCACCTGATGCTCCTCATCATCCGCTGGCCTGACACTATCA 4100  
 GAGCTCGCGCCGGCTGTTGCCAGGGACAGACTGACTACACTTGACCTTCAAGAGCACTTAGAAGTGGATGGCCTCCAGA 4179  
 CTCTGTCAGCCTCTGCAGGGGCCACACAAGTCTCCCAGCCAAGTCCACAAGCCTCCATGGTTCCCTGGCTCCTCTCCT 4258  
 GTGGAGTGTCTGTTTGTATGTCTGAGGTCTGCTTTGGGTACCGCCCTGGGAACTGCTAACCTCCGATTGGTCCCTTTGT 4337  
 GTCTCTGTTTACTGTCCTCTTCTACCTCCAGGTCACTTAGCTCTGGCTGCTCTGGCTGGGAGTGGGGGGTGGGGATGCT 4416  
 GGCTGCACCCCCACCCTGGTCTGCCAACAGAACCTGGGGGCCTCACACGGGCTCCTGTCTTGCCAAGCTGGAGCTGAGC 4495  
 ACACCTGGCCCAGGCTGAGTGGGGCAGAGCAAACAAGTGAAGGGGATCTCTCTCCTTAGAGGGAGGTGGCCGAAGGTGT 4574  
 AGATCCAGCGAGGGAGCTGCCATCCCCGCCACCTTCATAGCAGCAAGACCTTCCCATTTCCAATCTCACCTCCAGCAG 4653  
 GGATATGACTTTGGACAACAAGGCTTTATTTGTAAATATGCTCTTAATATGCAACTTTGAGAATAAGATAGAAACATCA 4732  
 TGTATTTTAAATATAAAATGAAGTGTGACACACTGTATACAATTTAATATATATTTTTAGGATTTTGTATTTTAAGAA 4811  
 AATGGAATGTGATGGTACTTAACTTTTACAAAAGAGAGAAAATGTTATTTTACTGTTTGAAGAAAATAAATATTCTCA 4890  
 TTGTTGTAGAAAAAAAAAAAAAAAAAAGGGCGGCCGC 4928

Fig. 1E

Hum. 10 20 30 40 50 60 70  
 MAPPAARLALLSAAALTLAARPAPSPGLGPGPECFTANGADYRGTQNWTLQGGKPCLFWNETFQHPYNT  
 Mur. 10 20 30 40 50 60  
 MAPPAARLALLSAAALTLAARPAPGPR--SGPECFTANGADYRGTQSWTLQGGKPCLFWNETFQHPYNT

Hum. 80 90 100 110 120 130 140  
 LKYPNGEGGLGEHNYCRNPDGDVSPWCYVAEHEDGVYWKYCEIPACQMPGNLGCYKDHGNPPPLTGTSTKT  
 Mur. 70 80 90 100 110 120 130  
 LKYPNGEGGLGEHNYCRNPDGDVSPWCYVAEHEDGVYWKYCEIPACQMPGNLGCYKDHGNPPPLTGTSTKT

Hum. 150 160 170 180 190 200 210  
 SNKLTITQTCISFCRSQRFKFAGMESGYACFCGNNPDYWKYGEAASTECNSVCFGDHTQPCGGDGRIILFD  
 Mur. 140 150 160 170 180 190 200  
 SNKLTITQTCISFCRSQRFKFAGMESGYACFCGNNPDYWKYGEAASTECNSVCFGDHTQPCGGDGRIILFD

Hum. 220 230 240 250 260 270 280  
 TLVGACGGNYSAMSSVVYSPDFPDYATGRVCYWTIRVPGASHIHFSFPLFDIRDSADMVELLDGYTHRV  
 Mur. 210 220 230 240 250 260 270  
 TLVGACGGNYSAMAAVVYSPDFPDYATGRVCYWTIRVPGASRIHFNFTLFDIRDSADMVELLDGYTHRV

Hum. 290 300 310 320 330 340 350  
 LARFHGRSRPPLSFNVSLDFVILYFFSDRINQAQGFVLYQAVKEELPQERPAVNQTVAEVITEQANLSV  
 Mur. 280 290 300 310 320 330 340  
 LVRLSGRSRPPLSFNVSLDFVILYFFSDRINQAQGFVLYQATKEEPPQERPAVNQTLAEVITEQANLSV

Hum. 360 370 380 390 400 410 420  
 SAARSSKVLVYVITTPSHPPQTVPGSNSWAPPMGAGSHRVEGWTVYGLATLLILTVAIVAKILLHVTFK  
 Mur. 350 360 370 380 390 400 410  
 SAAHSSKVLVYVITTPSHPPQTAQVAIPGHRQLGPTA---TEWKD-GLCTAWRPSSSSSQQLSQRFFCM

Hum. 430 440 450 460 470  
 SHRVPASGDLRDCHQPGTSGEIWSIFYKPSTISISIFKKKLKGQSQ-QDDRNPLVSD  
 Mur. 420 430 440 450 460 470  
 SHLNLIESLHQETLGTVVSLGLLEISGPFSSMNLPLQSPSLRRSSRVVRVKNKMTAIPS

Fig. 1F

Fig. 1G





Fig. 1H



GCGGCCGCTCGCGATCTAGAACTAGTA																			M	M	L	P	Q	N	S	W	H	I	D	F	G	13
ATG ATG CTG CCT CAA AAC TCG TGG CAT ATT GAT TTT GGA																			66													
R	C	C	C	H	Q	N	L	F	S	A	V	V	T	C	I	L	L	L	N	33												
AGA	TGC	TGC	TGT	CAT	CAG	AAC	CTT	TTC	TCT	GCT	GTG	GTA	ACT	TGC	ATC	CTG	CTC	CTG	AAT	126												
S	C	F	L	I	S	S	F	N	G	T	D	L	E	L	R	L	V	N	G	53												
TCC	TGC	TTT	CTC	ATC	AGC	AGT	TTT	AAT	GGA	ACA	GAT	TTG	GAG	TTG	AGG	CTG	GTC	AAT	GGA	186												
D	G	P	C	S	G	T	V	E	V	K	F	Q	G	Q	W	G	T	V	C	73												
GAC	GGT	CCC	TGC	TCT	GGG	ACA	GTG	GAG	GTG	AAA	TTC	CAG	GGA	CAG	TGG	GGG	ACT	GTG	TGT	246												
D	D	G	W	N	T	T	A	S	T	V	V	C	K	Q	L	G	C	P	F	93												
GAT	GAT	GGG	TGG	AAC	ACT	ACT	GCC	TCA	ACT	GTC	GTG	TGC	AAA	CAG	CTT	GGA	TGT	CCA	TTT	306												
S	F	A	M	F	R	F	G	Q	A	V	T	R	H	G	K	I	W	L	D	113												
TCT	TTC	GCC	ATG	TTT	CGT	TTT	GGA	CAA	GCC	GTG	ACT	AGA	CAT	GGA	AAA	ATT	TGG	CTT	GAT	366												
D	V	S	C	Y	G	N	E	S	A	L	W	E	C	Q	H	R	E	W	G	133												
GAT	GTT	TCC	TGT	TAT	GGA	AAT	GAG	TCA	GCT	CTC	TGG	GAA	TGT	CAA	CAC	CGG	GAA	TGG	GGA	426												
S	H	N	C	Y	H	G	E	D	V	G	V	N	C	Y	G	E	A	N	L	153												
AGC	CAT	AAC	TGT	TAT	CAT	GGA	GAA	GAT	GTT	GGT	GTG	AAC	TGT	TAT	GGT	GAA	GCC	AAT	CTG	486												
G	L	R	L	V	D	G	N	N	S	C	S	G	R	V	E	V	K	F	Q	173												
GGT	TTG	AGG	CTA	GTG	GAT	GGA	AAC	AAC	TCC	TGT	TCA	GGG	AGA	GTG	GAG	GTG	AAA	TTC	CAA	546												
E	R	W	G	T	I	C	D	D	G	W	N	L	N	T	A	A	V	V	C	193												
GAA	AGG	TGG	GGG	ACT	ATA	TGT	GAT	GAT	GGG	TGG	AAC	TTG	AAT	ACT	GCT	GCC	GTG	GTG	TGC	606												
R	Q	L	G	C	P	S	S	F	I	S	S	G	V	V	N	S	P	A	V	213												
AGG	CAA	CTA	GGA	TGT	CCA	TCT	TCT	TTT	ATT	TCT	TCT	GGA	GTT	GTT	AAT	AGC	CCT	GCT	GTA	666												
L	R	P	I	W	L	D	D	I	L	C	Q	G	N	E	L	A	L	W	N	233												
TTG	CGC	CCC	ATT	TGG	CTG	GAT	GAC	ATT	TTA	TGC	CAG	GGG	AAT	GAG	TTG	GCA	CTC	TGG	AAT	726												
C	R	H	R	G	W	G	N	H	D	C	S	H	N	E	D	V	T	L	T	253												
TGC	AGA	CAT	CGT	GGA	TGG	GGA	AAT	CAT	GAC	TGC	AGT	CAC	AAT	GAG	GAT	GTC	ACA	TTA	ACT	786												
C	Y	D	S	S	D	L	E	L	R	L	V	G	G	T	N	R	C	M	G	273												
TGT	TAT	GAT	AGT	AGT	GAT	CTT	GAA	CTA	AGG	CTT	GTA	GGT	GGA	ACT	AAC	CGC	TGT	ATG	GGG	846												
R	V	E	L	K	I	Q	G	R	W	G	T	V	C	H	H	K	W	N	N	293												
AGA	GTA	GAG	CTG	AAA	ATC	CAA	GGA	AGG	TGG	GGG	ACC	GTA	TGC	CAC	CAT	AAG	TGG	AAC	AAT	906												
A	A	A	D	V	V	C	K	Q	L	G	C	G	T	A	L	H	F	A	G	313												
GCT	GCA	GCT	GAT	GTC	GTA	TGC	AAG	CAG	TTG	GGA	TGT	GGA	ACC	GCA	CTT	CAC	TTC	GCT	GGC	966												
L	P	H	L	Q	S	G	S	D	V	V	W	L	D	G	V	S	C	S	G	333												
TTG	CCT	CAT	TTG	CAG	TCA	GGG	TCT	GAT	GTT	GTA	TGG	CTT	GAT	GGT	GTC	TCC	TGC	TCC	GGT	1026												
N	E	S	F	L	W	D	C	R	H	S	G	T	V	N	F	D	C	L	H	353												
AAT	GAA	TCT	TTT	CTT	TGG	GAC	TGC	AGA	CAT	TCC	GGA	ACC	GTC	AAT	TTT	GAC	TGT	CTT	CAT	1086												

Q	N	D	V	S	V	I	C	S	D	G	A	D	L	E	L	R	L	A	D	373
CAA	AAC	GAT	GTG	TCT	GTG	ATC	TGC	TCA	GAT	GGA	GCA	GAT	TTG	GAA	CTG	CGA	CTA	GCA	GAT	1146
G	S	N	N	C	S	G	R	V	E	V	R	I	H	E	Q	W	W	T	I	393
GGA	AGT	AAC	AAT	TGT	TCA	GGG	AGA	GTA	GAG	GTG	AGA	ATT	CAT	GAA	CAG	TGG	TGG	ACA	ATA	1206
C	D	Q	N	W	K	N	E	Q	A	L	V	V	C	K	Q	L	G	C	P	413
TGT	GAC	CAG	AAC	TGG	AAG	AAT	GAA	CAA	GCC	CTT	GTG	GTT	TGT	AAG	CAG	CTA	GGA	TGT	CCG	1266
F	S	V	F	G	S	R	R	A	K	P	S	N	E	A	R	D	I	W	I	433
TTC	AGC	GTC	TTT	GGC	AGT	CGT	CGT	GCT	AAA	CCT	AGT	AAT	GAA	GCT	AGA	GAC	ATT	TGG	ATA	1326
N	S	I	S	C	T	G	N	E	S	A	L	W	D	C	T	Y	D	G	K	453
AAC	AGC	ATA	TCT	TGC	ACT	GGG	AAT	GAG	TCA	GCT	CTC	TGG	GAC	TGC	ACA	TAT	GAT	GGA	AAA	1386
A	K	R	T	C	F	R	R	S	D	A	G	V	I	C	S	D	K	A	D	473
GCA	AAG	CGA	ACA	TGC	TTC	CGA	AGA	TCA	GAT	GCT	GGA	GTA	ATT	TGT	TCT	GAT	AAG	GCA	GAT	1446
L	D	L	R	L	V	G	A	H	S	P	C	Y	G	R	L	E	V	K	Y	493
CTG	GAC	CTA	AGG	CTT	GTC	GGG	GCT	CAT	AGC	CCC	TGT	TAT	GGG	AGA	TTG	GAG	GTG	AAA	TAC	1506
Q	G	E	W	G	T	V	C	H	D	R	W	S	T	R	N	A	A	V	V	513
CAA	GGA	GAG	TGG	GGG	ACT	GTG	TGT	CAT	GAC	AGA	TGG	AGC	ACA	AGG	AAT	GCA	GCT	GTT	GTG	1566
C	K	Q	L	G	C	G	K	P	M	H	V	F	G	M	T	Y	F	K	E	533
TGT	AAA	CAA	TTG	GGA	TGT	GGA	AAG	CCT	ATG	CAT	GTG	TTT	GGT	ATG	ACC	TAT	TTT	AAA	GAA	1626
A	S	G	P	I	W	L	D	D	V	S	C	I	G	N	E	S	N	I	W	553
GCA	TCA	GGA	CCT	ATT	TGG	CTG	GAT	GAC	GTT	TCT	TGC	ATT	GGA	AAT	GAG	TCA	AAT	ATC	TGG	1686
D	C	E	H	S	G	W	G	K	H	N	C	V	H	R	E	D	V	I	V	573
GAC	TGT	GAA	CAC	AGT	GGA	TGG	GGA	AAG	CAT	AAT	TGT	GTA	CAC	AGA	GAG	GAT	GTG	ATT	GTA	1746
C	S	G	D	A	T	W	G	L	R	L	V	G	G	S	N	R	C	S		593
ACC	TGC	TCA	GGT	GAT	GCA	ACA	TGG	GGC	CTG	AGG	CTG	GTG	GGC	GGC	AGC	AAC	CGC	TGC	TCG	1806
G	R	L	E	V	Y	F	Q	G	R	W	G	T	V	C	D	D	G	W	N	613
GGA	AGA	CTG	GAG	GTG	TAC	TTT	CAA	GGA	CGG	TGG	GGC	ACA	GTG	TGT	GAT	GAC	GGC	TGG	AAC	1866
S	K	A	A	A	V	V	C	S	Q	L	D	C	P	S	S	I	I	G	M	633
AGT	AAA	GCT	GCA	GCT	GTG	GTG	TGT	AGC	CAG	CTG	GAC	TGC	CCA	TCT	TCT	ATC	ATT	GGC	ATG	1926
G	L	G	N	A	S	T	G	Y	G	K	I	W	L	D	D	V	S	C	D	653
GGT	CTG	GGA	AAC	GCT	TCT	ACA	GGA	TAT	GGA	AAA	ATT	TGG	CTC	GAT	GAT	GTT	TCC	TGT	GAT	1986
G	D	E	S	D	L	W	S	C	R	N	S	G	W	G	N	N	D	C	S	673
GGA	GAT	GAG	TCA	GAT	CTC	TGG	TCA	TGC	AGG	AAC	AGT	GGG	TGG	GGA	AAT	AAT	GAC	TGC	AGT	2046
H	S	E	D	V	G	V	I	C	S	D	A	S	D	M	E	L	R	L	V	693
CAC	AGT	GAA	GAT	GTT	GGA	GTG	ATC	TGT	TCT	GAT	GCA	TCG	GAT	ATG	GAG	CTG	AGG	CTT	GTG	2106
G	G	S	S	R	C	A	G	K	V	E	V	N	V	Q	G	A	V	G	I	713
GGT	GGA	AGC	AGC	AGG	TGT	GCT	GGA	AAA	GTT	GAG	GTG	AAT	GTC	CAG	GGT	GCC	GTG	GGA	ATT	2166
L	C	A	N	G	W	G	M	N	I	A	E	V	V	C	R	Q	L	E	C	733
CTG	TGT	GCT	AAT	GGC	TGG	GGA	ATG	AAC	ATT	GCT	GAA	GTT	GTT	TGC	AGG	CAA	CTT	GAA	TGT	2226

G	S	A	I	R	V	S	R	E	P	H	F	T	E	R	T	L	H	I	L	753
GGG	TCT	GCA	ATC	AGG	GTC	TCC	AGA	GAG	CCT	CAT	TTC	ACA	GAA	AGA	ACA	TTA	CAC	ATC	TTA	2286
M	S	N	S	G	C	T	G	G	E	A	S	L	W	D	C	I	R	W	E	773
ATG	TCG	AAT	TCT	GGC	TGC	ACT	GGA	GGG	GAA	GCC	TCT	CTC	TGG	GAT	TGT	ATA	CGA	TGG	GAG	2346
W	K	Q	T	A	C	H	L	N	M	E	A	S	L	I	C	S	A	H	R	793
TGG	AAA	CAG	ACT	GCG	TGT	CAT	TTA	AAT	ATG	GAA	GCA	AGT	TTG	ATC	TGC	TCA	GCC	CAC	AGG	2406
Q	P	R	L	V	G	A	D	M	P	C	S	G	R	V	E	V	K	H	A	813
CAG	CCC	AGG	CTG	GTT	GGA	GCT	GAT	ATG	CCC	TGC	TCT	GGA	CGT	GTT	GAA	GTG	AAA	CAT	GCA	2466
D	T	W	R	S	V	C	D	S	D	F	S	L	H	A	A	N	V	L	C	833
GAC	ACA	TGG	CGC	TCT	GTC	TGT	GAT	TCT	GAT	TTC	TCT	CTT	CAT	GCT	GCC	AAT	GTG	CTG	TGC	2526
R	E	L	N	C	G	D	A	I	S	L	S	V	G	D	H	F	G	K	G	853
AGA	GAA	TTA	AAT	TGT	GGA	GAT	GCC	ATA	TCT	CTT	TCT	GTG	GGA	GAT	CAC	TTT	GGA	AAA	GGG	2586
N	G	L	T	W	A	E	K	F	Q	C	E	G	S	E	T	H	L	A	L	873
AAT	GGT	CTA	ACT	TGG	GCC	GAA	AAG	TTC	CAG	TGT	GAA	GGG	AGT	GAA	ACT	CAC	CTT	GCA	TTA	2646
C	P	I	V	Q	H	P	E	D	T	C	I	H	S	R	E	V	G	V	V	893
TGC	CCC	ATT	GTT	CAA	CAT	CCG	GAA	GAC	ACT	TGT	ATC	CAC	AGC	AGA	GAA	GTT	GGA	GTT	GTC	2706
C	S	R	Y	T	D	V	R	L	V	N	G	K	S	Q	C	D	G	Q	V	913
TGT	TCC	CGA	TAT	ACA	GAT	GTC	CGA	CTT	GTG	AAT	GGC	AAA	TCC	CAG	TGT	GAC	GGG	CAA	GTG	2766
E	I	N	V	L	G	H	W	G	S	L	C	D	T	H	W	D	P	E	D	933
GAG	ATC	AAC	GTG	CTT	GGA	CAC	TGG	GGC	TCA	CTG	TGT	GAC	ACC	CAC	TGG	GAC	CCA	GAA	GAT	2826
A	R	V	L	C	R	Q	L	S	C	G	T	A	L	S	T	T	G	G	K	953
GCC	CGT	GTT	CTA	TGC	AGA	CAG	CTC	AGC	TGT	GGG	ACT	GCT	CTC	TCA	ACC	ACA	GGA	GGA	AAA	2886
I	G	E	R	S	V	R	V	W	G	H	R	F	H	C	L	G	N	E		973
TAT	ATT	GGA	GAA	AGA	AGT	GTT	CGT	GTG	TGG	GGA	CAC	AGG	TTT	CAT	TGC	TTA	GGG	AAT	GAG	2946
S	L	L	D	N	C	Q	M	T	V	L	G	A	P	P	C	I	H	G	N	993
TCA	CTT	CTG	GAT	AAC	TGT	CAA	ATG	ACA	GTT	CTT	GGA	GCA	CCT	CCC	TGT	ATC	CAT	GGA	AAT	3006
T	V	S	V	I	C	T	G	S	L	T	Q	P	L	F	P	C	L	A	N	1013
ACT	GTC	TCT	GTG	ATC	TGC	ACA	GGA	AGC	CTG	ACC	CAG	CCA	CTG	TTT	CCA	TGC	CTC	GCA	AAT	3066
V	S	D	P	Y	L	S	A	V	P	E	G	S	A	L	I	C	L	E	D	1033
GTA	TCT	GAC	CCA	TAT	TTG	TCT	GCA	GTT	CCA	GAG	GGC	AGT	GCT	TTG	ATC	TGC	TTA	GAG	GAC	3126
K	R	L	R	L	V	D	G	D	S	R	C	A	G	R	V	E	I	Y	H	1053
AAA	CGG	CTC	CGC	CTA	GTG	GAT	GGG	GAC	AGC	CGC	TGT	GCC	GGG	AGA	GTA	GAG	ATC	TAT	CAC	3186
D	G	F	W	G	T	I	C	D	D	G	W	D	L	S	D	A	H	V	V	1073
GAC	GGC	TTC	TGG	GGC	ACC	ATC	TGT	GAT	GAC	GGC	TGG	GAC	CTG	AGC	GAT	GCC	CAC	GTG	GTG	3246
C	Q	K	L	G	C	G	V	A	F	N	A	T	V	S	A	H	F	G	E	1093
TGT	CAA	AAG	CTG	GGC	TGT	GGA	GTG	GCC	TTC	AAT	GCC	ACG	GTC	TCT	GCT	CAC	TTT	GGG	GAG	3306
G	S	G	P	I	W	L	D	D	L	N	C	T	G	T	E	S	H	L	W	1113
GGG	TCA	GGG	CCC	ATC	TGG	CTG	GAT	GAC	CTG	AAC	TGC	ACA	GGA	ACG	GAG	TCC	CAC	TTG	TGG	3366

Q	C	P	S	R	G	W	G	Q	H	D	C	R	H	K	E	D	A	G	V	1133
CAG	TGC	CCT	TCC	CGC	GGC	TGG	GGG	CAG	CAC	GAC	TGC	AGG	CAC	AAG	GAG	GAC	GCA	GGG	GTC	3426
I	C	S	E	F	T	A	L	R	L	Y	S	E	T	E	T	E	S	C	A	1153
ATC	TGC	TCA	GAA	TTC	ACA	GCC	TTG	AGG	CTC	TAC	AGT	GAA	ACT	GAA	ACA	GAG	AGC	TGT	GCT	3486
G	R	L	E	V	F	Y	N	G	T	W	G	S	V	G	R	R	N	I	T	1173
GGG	AGA	TTG	GAA	GTC	TTC	TAT	AAC	GGG	ACC	TGG	GGC	AGC	GTC	GGC	AGG	AGG	AAC	ATC	ACC	3546
T	A	I	A	G	I	V	C	R	Q	L	G	C	G	E	N	G	V	V	S	1193
ACA	GCC	ATA	GCA	GGC	ATT	GTG	TGC	AGG	CAG	CTG	GGC	TGT	GGG	GAG	AAT	GGA	GTT	GTC	AGC	3606
L	A	P	L	S	K	T	G	S	G	F	M	W	V	D	D	I	Q	C	P	1213
CTC	GCC	CCT	TTA	TCT	AAG	ACA	GGC	TCT	GGT	TTC	ATG	TGG	GTG	GAT	GAC	ATT	CAG	TGT	CCT	3666
K	T	H	I	S	I	W	Q	C	L	S	A	P	W	E	R	R	I	S	S	1233
AAA	ACG	CAT	ATC	TCC	ATA	TGG	CAG	TGC	CTG	TCT	GCC	CCA	TGG	GAG	CGA	AGA	ATC	TCC	AGC	3726
P	A	E	E	T	W	I	T	C	E	D	R	I	R	V	R	G	G	D	T	1253
CCA	GCA	GAA	GAG	ACC	TGG	ATC	ACA	TGT	GAA	GAT	AGA	ATA	AGA	GTG	CGT	GGA	GGA	GAC	ACC	3786
E	C	S	G	R	V	E	I	W	H	A	G	S	W	G	T	V	C	D	D	1273
GAG	TGC	TCT	GGG	AGA	GTG	GAG	ATC	TGG	CAC	GCA	GGC	TCC	TGG	GGC	ACA	GTG	TGT	GAT	GAC	3846
S	W	D	L	A	E	A	E	V	V	C	Q	Q	L	G	C	G	S	A	L	1293
TCC	TGG	GAC	CTG	GCC	GAG	GCG	GAA	GTG	GTG	TGT	CAG	CAG	CTG	GGC	TGT	GGC	TCT	GCT	CTG	3906
A	A	L	R	D	A	S	F	G	Q	G	T	G	T	I	W	L	D	D	M	1313
GCT	GCC	CTG	AGG	GAC	GCT	TCG	TTT	GGC	CAG	GGA	ACT	GGA	ACC	ATC	TGG	TTG	GAT	GAC	ATG	3966
R	C	K	G	N	E	S	F	L	W	D	C	H	A	K	P	W	G	Q	S	1333
CGG	TGC	AAA	GGA	AAT	GAG	TCA	TTT	CTA	TGG	GAC	TGT	CAC	GCC	AAA	CCC	TGG	GGA	CAG	AGT	4026
D	C	G	H	K	E	D	A	G	V	R	C	S	G	Q	S	L	K	S	L	1353
GAC	TGT	GGA	CAC	AAG	GAA	GAT	GCT	GGC	GTG	AGG	TGC	TCT	GGA	CAG	TCG	CTG	AAA	TCA	CTG	4086
N	A	S	S	G	H	L	A	L	I	L	S	S	I	F	G	L	L	L	L	1373
AAT	GCC	TCC	TCA	GGT	CAT	TTA	GCA	CTT	ATT	TTA	TCC	AGT	ATC	TTT	GGG	CTC	CTT	CTC	CTG	4146
V	L	F	I	L	F	L	T	W	C	R	V	Q	K	Q	K	H	L	P	L	1393
GTT	CTG	TTT	ATT	CTA	TTT	CTC	ACG	TGG	TGC	CGA	GTT	CAG	AAA	CAA	AAA	CAT	CTG	CCC	CTC	4206
R	V	S	T	R	R	R	G	S	L	E	E	N	L	F	H	E	M	E	T	1413
AGA	GTT	TCA	ACC	AGA	AGG	AGG	GGT	TCT	CTC	GAG	GAG	AAT	TTA	TTC	CAT	GAG	ATG	GAG	ACC	4266
C	L	K	R	E	D	P	H	G	T	R	T	S	D	D	T	P	N	H	G	1433
TGC	CTC	AAG	AGA	GAG	GAC	CCA	CAT	GGG	ACA	AGA	ACC	TCA	GAT	GAC	ACC	CCC	AAC	CAT	GGT	

AC

4628

Fig. 2E

CHIRP 3000000000




Fig. 2F

Fig. 2G

	10	20	30	40	50	60	70
Hum	MMLPQNSWHIDFGRCCCHQNLFSVAVTICILLNSCFLISSFNGTDLELRLVNGDGPCSGTVEVKFQGQWG						
	:	:	:	:	:	:	:
WCI	MAL-----GR---HLSLRGL---CVLLLTGT--MVG---GQALELRLKDGVRCEGRVEVKHQGEWG						
		10		20		30	40
							50

	80	90	100	110	120	130
Hum	TVCDGWNNTASTVVKQLGCPFSFAMFRFGQAVTR-HGKIWLDDVSCYGNESALWECQH---REWGSHN					
	:	:	:	:	:	:
WCI	TVDGYRWTLLKDAVVCRLGCGAAIG-FPGGAYFGPGLPIWLLYTSCEGTSTVSDCEHSNIKDYRNDG					
	60	70	80	90	100	110

	140	150	160	170	180	190	200
Hum	CYHGEDVGVNCYGEANLGLRLVDGNNSCSGRVEVKFQERWGTICDDGWNLTAAVVCRLGCPSSFISSG						
	:	:	:	:	:	:	:
WCI	YNHGRDAGVVCSG----FVRLAGGDGPCSGRVEVHSGEAWIPVSDGNFTLATAQIICAELGCGKAVSVLG						
	120	130	140	150	160	170	180

	210	220	230	240	250	260	270
Hum	VWNPAVLRPIWLDDILCQGNELALWNCRRHGWGNHDCSHNEDVTLTCTYDSSDLELRLVGGTNRRCMGRVE						
	:	:	:	:	:	:	:
WCI	HELFPRESSAQVWAEFRCEGEEPELWVCPRVPCPGGTCHHSGSAQVVCSEVRL-MTNGSSQCEGQVE						
	190	200	210	220	230	240	250

	280	290	300	310	320	330	340
Hum	LKIQGRWGTVCHHKWNNAAADVCKQLGCGTALHFAGLPHLQSGSDVWLDGVSCSGNESFLWDCRHSGT						
	:	:	:	:	:	:	:
WCI	MNISQWRALCASHWSLANANVICRLGCGVAISTPGGPHLVEEGDQILTARFHCSGAESFLWSCPVTAL						
	260	270	280	290	300	310	320

	350	360	370	380	390	400	410
Hum	VNFDCLHQNDVSVICSDGADLELRLADGSNNCSGRVEVRIHEQWWTICDQNWKEQALVVKQLGCPFSV						
	:	:	:	:	:	:	:
WCI	GGPDCSHGNTASVICS-GNQI-----QVLPQCND-----SV						
	330	340					350

	420	430	440	450	460	470	480
Hum	FGSRRAKPSNEARDIWINISICTGNESALWDCTYDGKAKRTCRRSDAGVICSDKADLDLRLVGAHSPCY						
	:	:	:	:	:	:	:
WCI	-----SQPTGSA-----ASEDSA----PY-----CSDSRQL--RLVDGGGPCA						
		360				370	380

	490	500	510	520	530	540	550
Hum	GRLEVKYQGEWGTVCHDRWSTRNAAVVCRLGCGKPMHVFGMTYFKEASGPIWLDDVSCIGNESNIWDCE						
	:	:	:	:	:	:	:
WCI	GRVEILDQGSWGTICDDGWLDDARVVCRLGCGEALNATGSAHFGAGSGPIWLDNLNCTGKESHVWRCP						
	390	400	410	420	430	440	450

Fig. 2H

	1190	1200	1210	1220	1230	1240
Hum	CGENGVS LAPLSKTGSGFMWVDDIQCPKTHISIWQCLSAPWERRISSPAETWITCEDR-----					
WCI	CGDSGSLNTSVGLREGSRPRWVDLIQCRKMDTSLWQCPSPGWKYSSCSPKEEAYISCEGRRPKSCPTAAA					
	1080	1090	1100	1110	1120	1130 1140
	1250 1260 1270 1280 1290 1300					
Hum	-----IRVRGGDTECSGRVEIWHAGSWGTVCDSDWDLAAEAVVCQQLGCGSALAAALRDASFGQGTGTIWW					
WCI	CTDREKLRLRGGDSECSGRVEVWHNGSWGTVCDSDWSLAAEAVVCQQLGCGQALEAVRSAAFGPGNGSIW					
	1150	1160	1170	1180	1190	1200 1210
	1310 1320 1330 1340 1350 1360					
Hum	LDDMRCKGNESFLWDCHAKPWGQSDCGHKEDAGVRCSG-----QSLKSLNASSGHLALI					
WCI	LDEVQCGGRESSLWDCVAEPWGQSDCKHEEDAGVRCSGVRTTLPTTTAGTRTTNSNLPGIFSLPGVLCI					
	1220	1230	1240	1250	1260	1270 1280
	1370 1380 1390 1400 1410					
Hum	LSSIFGLLLLVLFLFTWCRVQK-----QKHLPLRVS-----TRRRG-----SLEENLFHEME					
WCI	LGSLFLVLVILVTQLLRW-RAERRALSSYEDALAEAVYEELDYLLTQKEGLGSPDQMTDVPDENYDDAE					
	1290	1300	1310	1320	1330	1340 1350
	1420 1430 1440					
Hum	TC-----LKREDPHGTRTSD-----DTPNHGCEDAS-----DTSLLGV					
WCI	EVPVPGTPSPSQGNREEVPPEKEDGVRSSQTGSFLNFSREAAANPGEGEESFWLLQKKGDAGYDDVELSA					
	1360	1370	1380	1390	1400	1410 1420
	1450					
Hum	LPASEAT-K					
WCI	LGTSPVTFSS					
	1430					

Fig. 2 I

Fig. 2J

	10	20	30	40	50	60	70
Hum	ATGATGCTGCCTCAAACTCGTGGCATATTGATTTTGAAGATGCTGCTGCATCAGAACCTTTTCTCTG						
	:::	....:::			.....:	::: : : :::	
WCI	ATG-----GCTC-TGG-----GCAGACA-----CCTCT-CCCTG						
		10				20	

	80	90	100	110	120	130	140
Hum	CTGTGGTAACTTGCATCCTGCTCCTGAATTCCTGCTTTCATCAGCAGTTTAAATGGAACAGATTGGA						
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	C-GGGGACTCT-GTGTCTCCTCCT-----CGGCA-----C-----CATGGTGGGTGGTCAAGCTCTGGA						
	30	40	50		60	70	80

	150	160	170	180	190	200	210
Hum	GTTGAGGCTGGTCAATGGAGACGGTCCCTGCTCTGGGACAGTGGAGGTGAAATTCCAGGGACAGTGGGGG						
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	GCTGAGGTTGAAGGATGGAGTCCATCGCTGTGAGGGGAGAGTGGAAAGTGAAGCACCAAGGAGAATGGGGC						
	90	100	110	120	130	140	150

	220	230	240	250	260	270
Hum	ACTGTGTGTGATGATGGGTGGAACACTACTGCCT-CAACTGTCGTGTGCAAACAGCTTGGATGTCCATTT					
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	ACAGTGGATGGTTACAGGTGGA-CATTGAAGGATGCATCTGTAGTGTGCAGACAGCTGGGGTGTGGAGCT					
	160	170	180	190	200	210

	280	290	300	310	320	330	340
Hum	TCTTTCGCCATGTTTCGTTTTGGACAAGCCGTGA--CTAGACATGGAAAAATTTGGCTTGATGATGTTTC						
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	GCCATTG--GTTTTCTGGAGGGGCTTATTTTGGGCCAGGACTTGGCCCCATTGGCTTTTGTATACTTC						
	220	230	240	250	260	270	280

	350	360	370	380	390	400	410
Hum	CTGTTATGGAATGAGTCAGCTCTCTGGGAATGTCAACACCGGAATGGGAAGCCATAACTGTTATCAT						
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	ATGTGAAGGGACAGAGTCAACTGTCAGTGACTGTGAGCAT-TCTAATATTAAAGAC-TATC-GTAATGAT						
	290	300	310	320	330	340	350

	420	430	440	450	460	470	480
Hum	GGAGAAGATGTTGGTGTGAACTGTTATGGTGAAGCCAA-TCTGGGTTTGAG--GCTAG-TGGATGGAAAC						
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	GGCTATAATCATGGTCGGGA---TGCTGGAGTAGTCTGCTCAGGATTGTGCGTCTGGCTGGAGGGGATG						
	360	370	380	390	400	410	420

	490	500	510	520	530	540	550
Hum	AACTCCTGTTTCAGGGAGAGTGGAGGTGAAATTCGAAGAAAGGTGGGGGACTATATGTGATGATGGGTGGA						
	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .	: : : : .
WCI	GAC-CCTGCTCAGGGCGAGTAGAAGTGCATT--CTGGAGAAGCTTGGATCCCAGTGT-CTGATGGGAACT						
	430	440	450	460	470	480	

Fig. 2K

1740            1750            1760            1770            1780            1790            1800  
 Hum ACATGGGGCCTGAGGCTGGTGGGCGGCAGCAACCGCTGCTCGGGAAGACTGGAGGTGTACTTTTCAAGGAC  
       :.    ::    ::    ::    ::    ::    ::    :    .    :    :    :    :    :    :    :    :    :    :    :    :    :    :    :    :  
 WCI -CT--GGCCCTCAGGATGGTGAGTGAGGACCAGCAGTGTGCTGGGTGGCTGGAAGTTTTCTACAATGGGA  
           1440            1450            1460            1470            1480            1490            1500

Fig. 2 M

Fig. 2N

		3060	3070	3080	3090	3100	3110	3120
Hum	:	ATATTTGTCTGCAGTTCCAGAGGGCAGTGC	TTTGATCTGCTTAGAGGACAAACGGCTCCGCCTAGTGGAT					
		.....	.....	... :	.....	.....	.....	.....
WCI	:	TGCAGGCTCTGCGGCCTCAGAGGAGAGT	TCTCCCTACTGCTCAGACAGCAGGCAGCTCCGCCTGGTGGAC					
		2740	2750	2760	2770	2780	2790	2800
		3130	3140	3150	3160	3170	3180	3190
Hum	:	GGGGACAGCCGCTGTGCCGGGAGAGTAGAGATCTATCACGACGGCTTCTGGGGCACCATCTGTGATGACG						
		.....	...	.....	...	.....	.....	.....
WCI	:	GGGGCGGTCCCTGCGGCGGGAGAGTGGAGATCCTTGACCAGGGCTCCTGGGGCACCATCTGTGATGATG						
		2810	2820	2830	2840	2850	2860	2870
		3200	3210	3220	3230	3240	3250	3260
Hum	:	GCTGGGACCTGAGCGATGCCACGTGGTGTGTCAAAAGCTGGGCTGTGGAGTGGCCTTCAATGCCACGGT						
		.....	.....	..	.....	.....	.....	.....
WCI	:	ACTGGGACCTGGACGATGCCCGTGTGGTGTGCAGGCAGCTGGGCTGTGGAGAAGCCCTCAATGCCACGGG						
		2880	2890	2900	2910	2920	2930	2940
		3270	3280	3290	3300	3310	3320	3330
Hum	:	CTCTGCTCACTTTGGGGAGGGGTGAGGGCCCATCTGGCTGGATGACCTGAACTGCACAGGAACGGAGTCC						
		.....	.....	.....	.....	.....	.....	.....
WCI	:	GTCTGCTCACTTCGGGGCAGGATCAGGGCCCATCTGGCTGGACGACCTGAACTGCACAGGAAAGGAGTCC						
		2950	2960	2970	2980	2990	3000	3010
		3340	3350	3360	3370	3380	3390	3400
Hum	:	CACTTGTGGCAGTGCCCTTCCCGCGGCTGGGGGCAGCAGCACTGCAGGCACAAGGAGGACGCAGGGGTCA						
		...	.....	.....	.....	.....	.....	.....
WCI	:	CACGTGTGGAGGTGCCCTTCCCGGGGCTGGGGGCGGCAGCACTGCAGACACAAGGAGGACGCCGGGGTCA						
		3020	3030	3040	3050	3060	3070	3080
		3410	3420	3430	3440	3450	3460	3470
Hum	:	TCTGCTCAGAATTCACAGCCTTGAGGCTCTACAGTGAAACTGAAACAGAGAGCTGTGCTGGGAGATTGGA						
		.....	.....	.....	.....	.....	.....	.....
WCI	:	TCTGCTCAGAGTTCTTGCCCTCAGGAT---	GGTGAG-CGAGGACCAGCAG-TGTGCTGGGTGGCTGGA					
		3090	3100	3110	3120	3130	3140	
		3480	3490	3500	3510	3520	3530	3540
Hum	:	AGTCTTCTATAACGGGACCTGGGGCAGCGTCGGCAGGAGGAACATCACCACAGCCATAGCAGGCATTGTG						
		...	.....	.....	.....	.....	.....	.....
WCI	:	GGTTTTCTACAACGGGACCTGGGGCAGTGTCTGCCGCAGCCCCATGGAAGATATCACTGTGTCCGTGATC						
		3150	3160	3170	3180	3190	3200	3210
		3550	3560	3570	3580	3590	3600	
Hum	:	TGCAGGCAGCTGGGCTGTGGGGAGAATGGAGTTGTGAGCCTCGCCCTTTA--TCT-AAGACAGGCTCTG						
		.....	.....	.....	.....	.....	.....	.....
WCI	:	TGCAGACAGCTTGGATGTGGGGACAGTGA--AGTCT-CAACACCTCTGTTGGTCTCAGGGAAGGTTCTA						
		3220	3230	3240	3250	3260	3270	3280
		3610	3620	3630	3640	3650	3660	3670
Hum	:	GTTTCATGTGGGTGGATGACATTCACTGTCTTAAACGCATATCTCCATATGGCAGTGCCTGTCTGCCCC						
		..	.....	.....	.....	.....	.....	.....
WCI	:	GACCCCGGTGGGTAGATTTAATTCACTGTGCGGAAAATGGATACCTCTCTCTGGCAGTGTCTTCTGGCCC						
		3290	3300	3310	3320	3330	3340	3350

Fig. 20

Fig. 2P

Fig. 2Q

GTGACCCACGCGTCCGGTCTGTGGCTGAGC																			M	A	L	P	A	L	G	L	D	P	W	S	
																			ATG	GCC	CTC	CCA	GCC	CTG	GGC	CTG	GAC	CCC	TGG	AGC	
																			12												
L	L	G	L	F	L	F	Q	L	L	Q	L	L	L	P	T	T	T	A	G												
CTC	CTG	GGC	CTT	TTC	CTC	TTC	CAA	CTG	CTT	CAG	CTG	CTG	CTG	CCG	ACG	ACG	ACC	GCG	GGG												
																			32												
G	G	G	Q	G	P	M	P	R	V	R	Y	Y	A	G	D	E	R	R	A												
GGA	GGC	GGG	CAG	GGG	CCC	ATG	CCC	AGG	GTC	AGA	TAC	TAT	GCA	GGG	GAT	GAA	CGT	AGG	GCA												
																			52												
L	S	F	F	H	Q	K	G	L	Q	D	F	D	T	L	L	L	S	G	D												
CTT	AGC	TTC	TTC	CAC	CAG	AAG	GGC	CTC	CAG	GAT	TTT	GAC	ACT	CTG	CTC	CTG	AGT	GGT	GAT												
																			72												
G	N	T	L	Y	V	G	A	R	E	A	I	L	A	L	D	I	Q	D	P												
GGA	AAT	ACT	CTC	TAC	GTG	GGG	GCT	CGA	GAA	GCC	ATT	CTG	GCC	TTG	GAT	ATC	CAG	GAT	CCA												
																			92												
G	V	P	R	L	K	N	M	I	P	W	P	A	S	D	R	K	K	S	E												
GGG	GTC	CCC	AGG	CTA	AAG	AAC	ATG	ATA	CCG	TGG	CCA	GCC	AGT	GAC	AGA	AAA	AAG	AGT	GAA												
																			112												
C	A	F	K	K	K	S	N	E	T	Q	C	F	N	F	I	R	V	L	V												
TGT	GCC	TTT	AAG	AAG	AAG	AGC	AAT	GAG	ACA	CAG	TGT	TTC	AAC	TTC	ATC	CGT	GTC	CTG	GTT												
																			132												
S	Y	N	V	T	H	L	Y	T	C	G	T	F	A	F	S	P	A	C	T												
TCT	TAC	AAT	GTC	ACC	CAT	CTC	TAC	ACC	TGC	GGC	ACC	TTC	GCC	TTC	AGC	CCT	GCT	TGT	ACC												
																			152												
R	I	E	L	Q	D	S	Y	L	L	P	I	S	E	D	K	V	M	E	G												
TTC	ATT	GAA	CTT	CAA	GAT	TCC	TAC	CTG	TTG	CCC	ATC	TCG	GAG	GAC	AAG	GTC	ATG	GAG	GGA												
																			172												
K	G	Q	S	P	F	D	P	A	H	K	H	T	A	V	L	V	D	G	M												
AAA	GGC	CAA	AGC	CCC	TTT	GAC	CCC	GCT	CAC	AAG	CAT	ACG	GCT	GTC	TTG	GTG	GAT	GGG	ATG												
																			192												
L	Y	S	G	T	M	N	N	F	L	G	S	E	P	I	L	M	R	T	L												
CTC	TAT	TCT	GGT	ACT	ATG	AAC	AAC	TTC	CTG	GGC	AGT	GAG	CCC	ATC	CTG	ATG	CGC	ACA	CTG												
																			212												
G	S	Q	P	V	L	K	T	D	N	F	L	R	W	L	H	H	D	A	S												
GGG	TCC	CAG	CCT	GTC	CTC	AAG	ACC	GAC	AAC	TTC	CTC	CGC	TGG	CTG	CAT	CAT	GAC	GCC	TCC												
																			232												
F	V	A	A	I	P	S	T	Q	V	V	Y	F	F	F	E	E	T	A	S												
TTT	GTG	GCA	GCC	ATC	CCT	TCG	ACC	CAG	GTC	GTC	TAC	TTC	TTC	TTC	GAG	GAG	ACA	GCC	AGC												
																			252												
E	F	D	F	F	E	R	L	H	T	S	R	V	A	R	V	C	K	N	D												
GAG	TTT	GAC	TTC	TTT	GAG	AGG	CTC	CAC	ACA	TCG	CGG	GTG	GCT	AGA	GTC	TGC	AAG	AAT	GAC												
																			272												
V	G	G	E	K	L	L	Q	K	K	W	T	T	F	L	K	A	Q	L	L												
GTG	GGC	GGC	GAA	AAG	CTG	CTG	CAG	AAG	AAG	TGG	ACC	ACC	TTC	CTG	AAG	GCC	CAG	CTG	CTC												
																			292												
C	T	Q	P	G	Q	L	P	F	N	V	I	R	H	A	V	L	L	P	A												
TGC	ACC	CAG	CCG	GGG	CAG	CTG	CCC	TTC	AAC	GTC	ATC	CGC	CAC	GCG	GTC	CTG	CTC	CCC	GCC												
																			312												
D	S	P	T	A	P	H	I	Y	A	V	F	T	S	Q	W	Q	V	G	G												
GAT	TCT	CCC	ACA	GCT	CCC	CAC	ATC	TAC	GCA	GTC	TTC	ACC	TCC	CAG	TGG	CAG	GTT	GGC	GGG												
																			332												
T	R	S	S	A	V	C	A	F	S	L	L	D	I	E	R	V	F	K	G												
ACC	AGG	AGC	TCT	GCG	GTT	TGT	GCC	TTC	TCT	CTC	TTG	GAC	ATT	GAA	CGT	GTC	TTT	AAG	GGG												
																			352												
																			1087												

K	Y	K	E	L	N	K	E	T	S	R	W	T	T	Y	R	G	P	E	T	372
AAA	TAC	AAA	GAG	TTG	AAC	AAA	GAA	ACT	TCA	CGC	TGG	ACT	ACT	TAT	AGG	GGC	CCT	GAG	ACC	1147
N	P	R	P	G	S	C	S	V	G	P	S	S	D	K	A	L	T	F	M	392
AAC	CCC	CGG	CCA	GGC	AGT	TGC	TCA	GTG	GGC	CCC	TCC	TCT	GAT	AAG	GCC	CTG	ACC	TTC	ATG	1207
K	D	H	F	L	M	D	E	Q	V	V	G	T	P	L	L	V	K	S	G	412
AAG	GAC	CAT	TTC	CTG	ATG	GAT	GAG	CAA	GTG	GTG	GGG	ACG	CCC	CTG	CTG	GTG	AAA	TCT	GGC	1267
V	E	Y	T	R	L	A	V	E	T	A	Q	G	L	D	G	H	S	H	L	432
GTG	GAG	TAT	ACA	CGG	CTT	GCA	GTG	GAG	ACA	GCC	CAG	GGC	CTT	GAT	GGG	CAC	AGC	CAT	CTT	1327
V	M	Y	L	G	T	T	T	G	S	L	H	K	A	V	V	S	G	D	S	452
GTC	ATG	TAC	CTG	GGA	ACC	ACC	ACA	GGG	TCG	CTC	CAC	AAG	GCT	GTG	GTA	AGT	GGG	GAC	AGC	1387
S	A	H	L	V	E	E	I	Q	L	F	P	D	P	E	P	V	R	N	L	472
AGT	GCT	CAT	CTG	GTG	GAA	GAG	ATT	CAG	CTG	TTC	CCT	GAC	CCT	GAA	CCT	GTT	CGC	AAC	CTG	1447
Q	L	A	P	T	Q	G	A	V	F	V	G	F	S	G	G	V	W	R	V	492
CAG	CTG	GCC	CCC	ACC	CAG	GGT	GCA	GTG	TTT	GTA	GGC	TTC	TCA	GGA	GGT	GTC	TGG	AGG	GTG	1507
P	R	A	N	C	S	V	Y	E	S	C	V	D	C	V	L	A	R	D	P	512
CCC	CGA	GCC	AAC	TGT	AGT	GTC	TAT	GAG	AGC	TGT	GTG	GAC	TGT	GTC	CTT	GCC	CGG	GAC	CCC	1567
H	C	A	W	D	P	E	S	R	T	C	C	L	L	S	A	P	N	L	N	532
CAC	TGT	GCC	TGG	GAC	CCT	GAG	TCC	CGA	ACC	TGT	TGC	CTC	CTG	TCT	GCC	CCC	AAC	CTG	AAC	1627
S	W	K	Q	D	M	E	R	G	N	P	E	W	A	C	A	S	G	P	M	552
TCC	TGG	AAG	CAG	GAC	ATG	GAG	CGG	GGG	AAC	CCA	GAG	TGG	GCA	TGT	GCC	AGT	GGC	CCC	ATG	1687
S	R	S	L	R	P	Q	S	R	P	Q	I	I	K	E	V	L	A	V	P	572
AGE	AGG	AGC	CTT	CGG	CCT	CAG	AGC	CGC	CCG	CAA	ATC	ATT	AAA	GAA	GTC	CTG	GCT	GTC	CCC	1747
N	S	I	L	E	L	P	C	P	H	L	S	A	L	A	S	Y	Y	W	S	592
AAC	TCC	ATC	CTG	GAG	CTC	CCC	TGC	CCC	CAC	CTG	TCA	GCC	TTG	GCC	TCT	TAT	TAT	TGG	AGT	1807
H	G	P	A	A	V	P	E	A	S	S	T	V	Y	N	G	S	L	L	L	612
CAT	GGC	CCA	GCA	GCA	GTC	CCA	GAA	GCC	TCT	TCC	ACT	GTC	TAC	AAT	GGC	TCC	CTC	TTG	CTG	1867
F	V	Q	D	G	V	G	G	L	Y	Q	C	W	A	T	E	N	G	F	S	632
ATA	GTG	CAG	GAT	GGA	GTT	GGG	GGT	CTC	TAC	CAG	TGC	TGG	GCA	ACT	GAG	AAT	GGC	TTT	TCA	1927
Y	P	V	I	S	Y	W	V	D	S	Q	D	Q	T	L	A	L	D	P	E	652
TAC	CCT	GTG	ATC	TCC	TAC	TGG	GTG	GAC	AGC	CAG	GAC	CAG	ACC	CTG	GCC	CTG	GAT	CCT	GAA	1987
L	A	G	I	P	R	E	H	V	K	V	P	L	T	R	V	S	G	G	A	672
CTG	GCA	GGC	ATC	CCC	CGG	GAG	CAT	GTG	AAG	GTC	CCG	TTG	ACC	AGG	GTC	AGT	GGT	GGG	GCC	2047
A	L	A	A	Q	Q	S	Y	W	P	H	F	V	T	V	T	V	L	F	A	692
GCC	CTG	GCT	GCC	CAG	CAG	TCC	TAC	TGG	CCC	CAC	TTT	GTC	ACT	GTC	ACT	GTC	CTC	TTT	GCC	2107
L	V	L	S	G	A	L	I	I	L	V	A	S	P	L	R	A	L	R	A	712
TTA	GTG	CTT	TCA	GGA	GCC	CTC	ATC	ATC	CTC	GTG	GCC	TCC	CCA	TTG	AGA	GCA	CTC	CGG	GCT	2167
R	G	K	V	Q	G	C	E	T	L	R	P	G	E	K	A	P	L	S	R	732
CGG	GGC	AAG	GTT	CAG	GGC	TGT	GAG	ACC	CTG	CGC	CCT	GGG	GAG	AAG	GCC	CCG	TTA	AGC	AGA	2227

27/53

E	Q	H	L	Q	S	P	K	E	C	R	T	S	A	S	D	V	D	A	D		752
GAG	CAA	CAC	CTC	CAG	TCT	CCC	AAG	GAA	TGC	AGG	ACC	TCT	GCC	AGT	GAT	GTG	GAC	GCT	GAC		2287
N	N	C	L	G	T	E	V	A	*												762
AAC	AAC	TGC	CTA	GGC	ACT	GAG	GTA	GCT	TAA												2317
ACTCTAGGCACAGGCCCGGGGCTGCGGTGCAGGCACCTGGCCATGCTGGCTGGGCGGCCCAAGCACAGCCCTGACTAGGA																				2396	
TGACAGCAGCACAAAAGACCACCTTTCTCCCCCTGAGAGGAGCTTCTGCTACTCTGCATCACTGATGACACTCAGCAGGG																				2475	
TGATGCACAGCAGTCTGCCTCCCCCTATGGGACTCCCTTCTACCAAGCACATGAGCTCTCTAACAGGGTGGGGGCTACCC																				2554	
CCAGACCTGCTCCTACACTGATATTGAAGAACCTGGAGAGGATCCTTCAGTTCTGGCCATTCCAGGGACCCCTCCAGAAA																				2633	
CACAGTGTTTCAAGAGATCCTAAAAAACCTGCCTGTCCCAGGACCCCTATGGTAAATGAACACCAAACATCTAAACAATC																				2712	
ATATGCTAACATGCCACTCCTGGAAACTCCACTCTGAAGCTGCCGCTTTGGACACCAACACTCCCTTCTCCCAGGGTCA																				2791	
TGCAGGGATCTGCTCCCTCCTGCTTCCCTTACCAGTCGTGCACCGCTGACTCCCAGGAAGTCTTTCTCTGAAGTCTGACC																				2870	
ACCTTTCTTCTTGCTTCAGTTGGGGCAGACTCTGATCCCTTCTGCCCTGGCAGAATGGCAGGGGTAATCTGAGCCTTCT																				2949	
TCCTCCTTTACCCTAGCTGACCCCTTCACCTCTCCCCCTCCCTTTTCCTTTGTTTTGGGATTTCAGAAAACCTGCTTGTC																				3028	
AGAGACTGTTTATTTTTTTATTAATAAGGCTTAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCCGC																				3104	

Fig. 3D

	10	20	30	40	50	60	70
Hum.	MALPALGLDPWSLLGLFLFQLLQLLP	TTTAGGGGQGPMPRVRYAGDERRALS	SFFHQKGLQDFD	TLLLS			
Mur.	MALPSLGQDSWSLLRVFFFLQLLPSL	PPASGTGGQGPMPRVKYHAGDGHRLS	SFFQKGLRDFD	TLLLS			
	10	20	30	40	50	60	70
	80	90	100	110	120	130	140
Hum.	GDGNTLYVGAREAILALDIQDPGV	PRLKNMIPWPASDRKKSECAFKKKS	NETQCFNFIRVLVSYNV	THLY			
Mur.	DDGNTLYVGARETVLALNIQNP	GIPRLKNMIPWPASERKKTECAFKKKS	NETQCFNFIRVLVSYN	NATHLY			
	80	90	100	110	120	130	140
	150	160	170	180	190	200	210
Hum.	TCGTFAFSPACTFIELQDSYLLP	ISEDKVMGKGQSPFDP	PAKHHTAVLVDGMLY	SGTMNNFLGSEP	ILMR		
Mur.	ACGTFAFSPACTFIELQDSLLP	LILDKVMDGKGQSP	TLTFTSTQAVLVDGMLY	SGTMNNFLGSEP	ILMR		
	150	160	170	180	190	200	210
	220	230	240	250	260	270	280
Hum.	TLGSQPVLKTDN	FLRWLHHDASFVAAIP	STQVVYFFFEETASE	FDFFERLHTSRVARV	CKNDVGGEKLLQ		
Mur.	TLGSHPVLKTDI	FLRWLHADASFVAAIP	STQVVYFFFEETASE	FDFFEELYISRVAQV	CKNDVGGEKLLQ		
	220	230	240	250	260	270	280
	290	300	310	320	330	340	350
Hum.	KKWTTFLKAQLL	CTQPGQLPFNVIRH	AVLLPADSPTAPHI	YAVFTSQWQVGG	TRSSAVCAFS	LLDIERVF	
Mur.	KKWTTFLKAQLL	CAQPGQLPFNIIRH	AVLLPADSPSVS	RIYAVFTSQWQVGG	TRSSAVCAFS	SLDIERVF	
	290	300	310	320	330	340	350
	360	370	380	390	400	410	420
Hum.	KGKYKELNKET	SRWTTYRGPE	TNPRPGSCSVGP	SSDKALTFMKDH	FLMDEQVVG	TPLLVKSGVEY	TRLAV
Mur.	KGKYKELNKET	SRWTTYRGSE	VSPRPGSCSMGP	SSDKALTFMKDH	FLMDEHVVG	TPLLVKSGVEY	TRLAV
	360	370	380	390	400	410	420
	430	440	450	460	470	480	490
Hum.	ETAQGLDGSH	LVMYLGT	TGSLHKAVVSGD	SSAHLVEEIQL	FPDPEPVRNL	QLAPTQGA	VFGVSGGVW
Mur.	ESARGLDGSS	HVMYLGT	STGPLHKAVVP	QDSSAYLVEET	QLSPDSE	PVRNLQLA	PAQGA
	430	440	450	460	470	480	490
	500	510	520	530	540	550	560
Hum.	RVPRANCSVY	ESCVDCLARDP	HCARDPES	RTCCLLSAPNL	NSWKQDMERGN	PEWACASG	PMSRSLRPQS
Mur.	RVPRANCSVY	ESCVDCLARDP	HCARDPES	RCLCSLLSG	ST-KPWQDMERGN	PEWVCTR	GPMPARRQS
	500	510	520	530	540	550	
	570	580	590	600	610	620	630
Hum.	RPQIIKEVL	AVPNSILEL	PCPHLSALASY	WWSHGPA	AVPEASSTV	YNGSLLLVQ	DGVGGLYQC
Mur.	PPQLIKEVL	TVPN	SILELRCPHLS	ALASYHWSH	GRAKISEAS	ATVYNGS	LLLLPQDGVGGLYQC
	560	570	580	590	600	610	620
	640	650	660	670	680	690	700
Hum.	FSYPVISY	WVDSQDQTL	ALDPELAGI	PREHVKVPL	TRVSGGA	ALAAQSYW	PHFVTVTVL
Mur.	YSYPVVS	YWVDSQDQ	PLALDPELAG	VPREVRQV	PLTRVGG	ASMAAQR	SYWPHFLIVTVLL
	630	640	650	660	670	680	690
	710	720	730	740	750	760	
Hum.	ILVASPLRAL	RARGKVQGC	ETLRPG	EKAPLSRE	QHLQSPKE	CRTSASD	VDADNNCLGTEVA
Mur.	LLLASPLGAL	RARGKVQGC	GMLPPRE	KAPLSRD	QHLQSPK	DHRTSAS	VDADNNHLGAEVA
	700	710	720	730	740	750	760

Fig. 3E

Hum. GTCG-AC-CC-----10-----ACG-----CGTCCGGT-----20-----CTGTGGCTGAGCATGGC  
 Mur. CTCGGACGCCTGGGTTAGGGGTCTGTACTGCTGGGGAACCATCTGGTGACCATCTCAGGCTGACCATGGC  
 10 20 30 40 50 60 70

Hum. CCTCCCAGCCCTGGGCTGGACCCCTGGAGCCTCCTGGGCCTTTTCTCTTCCAACTGCTTC-AGCTGCT  
 Mur. CCTACCATCCCTGGGCCAGGACTCATGGAGTCTCCTGCGTGTCTTTTCTTCCAACT-CTTCTGCTGCC  
 40 50 60 70 80 90 100 110 120 130

Hum. GCTGCCGACGACGACCGCGGGGGAGGCGGGCAGGGGCCCATGCCAGGGTCACTACTATGCAGGGGAT  
 Mur. ATCACTGCCACCTGCTTCTGGGACTGGTGGTCAAGGGGCCCATGCCAGAGTCAAATACCATGCTGGAGAC  
 110 120 130 140 150 160 170 180 190 200

Hum. GAACGTAGGGCACTTAGCTTCTTCCACCAGAAGGGCCTCCAGGATTTTGACACTCTGCTCCTGAGTGGTG  
 Mur. GGGCACAGGGCCCTCAGCTTCTTCCAAACAAAAGGCCCTCCGAGACTTTGACACGCTGCTCCTGAGTGACG  
 180 190 200 210 220 230 240 250 260 270

Hum. ATGGAAATACTCTCTACGTGGGGGCTCGAGAAGCCATTCTGGCCCTTGGATATCCAGGATCCAGGGGTCCC  
 Mur. ATGGCAACACTCTCTATGTGGGGGCTCGAGAGACCGTCTGGCCCTGAATATCCAGAACCAGGAATCCC  
 250 260 270 280 290 300 310 320 330 340

Hum. CAGGCTAAAGAACATGATACCGTGGCCAGCCAGTGACAGAAAAAGAGTGAATGTGCCTTTAAGAAGAAG  
 Mur. AAGGCTAAAGAACATGATACCGTGGCCAGCCAGTGAGAGAAAAAGACCGAATGTGCCTTTAAGAAGAAG  
 320 330 340 350 360 370 380 390 400 410

Hum. AGCAATGAGACACAGTGTTCAACTTCATCCGTGTCTGGTTCCTTACAATGTACCCATCTCTACACCT  
 Mur. AGCAATGAGACACAGTGTTCAACTTCATTGAGTCTGGTCTCTTACAATGTACTCACCTCTATGCCT  
 390 400 410 420 430 440 450 460 470 480

Hum. GCGGCACCTTCGCTTCAGCCCTGCTTGTACCTTCATTGAACTTCAAGATTCTTACCTGTTGCCCATCTC  
 Mur. GTGGGACCTTTGCCTTCAGCCCTGCTTGTACCTTCATTGAACTTCAAGATTCTTACCTGTTGCCCATCTT  
 460 470 480 490 500 510 520 530 540 550

Hum. GGAGGACAAGGTCATGGAGGGAAGGCCAAAGCCCTTTGACCCCGCTCACAAGCATACG-GCTGTCTT  
 Mur. GATAGACAAGGTCATGGACGGGAAGGGCCAAAGCCC-TTGACCCCTGTTTACAAGCACACAAGCTGTCTT  
 530 540 550 560 570 580 590 600 610 620

Hum. GGTGGATGGGATGCTCTATTCTGGTACTATGAACAACTTCCTGGGCAGTGAGCCCATCCTGATGCGCACA  
 Mur. GGTGATGGGATGCTTTATTCGGGACCATGAACAACTTCCTGGGCAGTGAGCCCATCCTGATGCGGACA  
 600 610 620 630 640 650 660 670 680 690

Hum. CTGGGATCCCAGCCTGTCTCAAGACCGACAACCTTCCTCCGCTGGCTGCATCATGACGCCTCCTTTGTGG  
 Mur. CTGGGATCCCATCCTGTCTCAAGACTGACATCTTCTTACGCTGGCTGCACGCGGATGCCTCCTTCGTGG  
 670 680 690 700 710 720 730 740 750 760

[illegible]

Hum. 1580 1590 1600 1610 1620 1630 1640  
 CCTGGGACCCTGAGTCCCGAACCTGTTGCCTCCTGTCTGCCCCAACCTGAACCTCTGGAAGCAGGACAT  
 Mur. 1610 1620 1630 1640 1650 1660 1670  
 CCTGGGACCCTGAATCAAGACTCTGCAGCCTTCTGTCTGGCTC-TACCAAGCCT--TGAAGCAGGACAT

Hum. 1650 1660 1670 1680 1690 1700 1710  
 GGAGCGGGGAACCCAGAGTGGGCATGTGCCAGTGGCCCCATGAGCAGGAGCCTTCGGCCTCAGAGCCGC  
 Mur. 1680 1690 1700 1710 1720 1730 1740  
 GGAACGCGGCAACCCGAGTGGGTATGACCCGTGGCCCCATGGCCAGGAGCCCCGGCGTCAGAGCCCC

Hum. 1720 1730 1740 1750 1760 1770 1780  
 CCGCAAATCATTAAGAAGTCTGGCTGTCCCAACTCCATCCTGGAGCTCCCTGCCCCACCTGTCTAG  
 Mur. 1750 1760 1770 1780 1790 1800 1810  
 CCTCAACTAATTAAGAAGTCTGACAGTCCCAACTCCATCCTGGAGCTGCCTGCCCCACCTGTCTAG

Hum. 1790 1800 1810 1820 1830 1840 1850  
 CCTTGGCCTCTTATTATTGGAGTCATGGCCAGCAGCAGTCCCAGAAGCCTCTTCCACTGTCTACAATGG  
 Mur. 1820 1830 1840 1850 1860 1870 1880  
 CACTGGCCTCTTACCAGTGGAGTCATGGCCGAGCCAAAATCTCAGAAGCCTCTGCTACCGTCTACAATGG-

Hum. 1860 1870 1880 1890 1900 1910 1920  
 CTCCCTCTTGCTGATAGTGCAGGATGGAGTTGGGGGTCTCTACCAGTGTGGGCAACTGAGAATGGCTTT  
 Mur. 1890 1900 1910 1920 1930 1940 1950  
 CTCCCTCTTGCTGCTGCCGAGGATGGTGTGGGGGCTCTACCAGTGTGTGGCGACTGAGAACGGCTAC

Hum. 1930 1940 1950 1960 1970 1980 1990  
 TCATACCCTGTGATCTCCTACTGGGTGGACAGCCAGGACCAGACCTGGCCCTGGATCCTGAACCTGGCAG  
 Mur. 1960 1970 1980 1990 2000 2010 2020  
 TCATACCCTGTGGTCTCCTATTGGGTAGACAGCCAGGACCAGCCCTGGCGCTGGACCTGAGCTGGCGG

Hum. 2000 2010 2020 2030 2040 2050 2060  
 GCATCCCCCGGGAGCATGTGAAGGTCCCGTTGACCAGGGTCAGTGGTGGGGCCGCCCTGGCTGCCCAGCA  
 Mur. 2030 2040 2050 2060 2070 2080 2090  
 GCGTTCCCGTGGAGCGTGTGCAGGTCCCGCTGACCAGGGTCGGAGGCGGAGCTTCCATGGCTGCCCAGCG

Hum. 2070 2080 2090 2100 2110 2120 2130  
 GTCCTACTGGCCCCACTTTGTCACTGTCACTGTCTCTTTGCTTAGTGCTTTCAGGAGCCCTCATCATC  
 Mur. 2100 2110 2120 2130 2140 2150 2160  
 GTCCTACTGGCCCCATTTTCTCATCGTTACCGTCTCTGGCCATCGTGTCTCTGGGAGTGCTCACTCTC

Hum. 2140 2150 2160 2170 2180 2190 2200  
 CTCGTGGCCTCCCCATTGAGAGCACTCCGGGCTCGGGGCAAGGTTAGGGCTGTGAGACCCTGCGCCCTG  
 Mur. 2170 2180 2190 2200 2210 2220 2230  
 CTCCTCGCTTCCCCACTGGGGGCGCTGCGGGCTCGGGGTAAGGTTAGGGCTGTGGGATGCTGCCCCCA

Hum. 2210 2220 2230 2240 2250 2260 2270  
 GGGAGAAGGCCCGTTAAGCAGAGAGCAACACCTCCAGTCTCCCAAGGAATGCAGGACCTCTGCCAGTGA  
 Mur. 2240 2250 2260 2270 2280 2290 2300  
 GGGAAAAGGCTCCACTGAGCAGGACCAGCACCTCCAGCCCTCCAAGGACCACAGGACCTCTGCCAGTGA

Hum. 2280 2290 2300 2310 2320 2330 2340  
 TGTGGACGCTGACAACAACCTAGGCAGTGGGTAGCTTAACTCTAGGCACAGG-CCGGGGCTG--C  
 Mur. 2310 2320 2330 2340 2350 2360 2370  
 CGTAGATGCCGACAACAACCATCTGGGCGCCGAAGTGGCTTAAACA-GGGACACAGATCCGCAGCTGAGC

Hum. 2350 2360 2370 2380 2390 2400 2410  
 GGTGCAGGCACCTGGCCATGCTGGCTGGGCGGCCCAAGCACAGCCCTGACTAGGATGACAGCAGCAAAA  
 Mur. 2380 2390 2400 2410 2420  
 AGAGCAAGCCACTGGCCTTGTGGCTATGC---CAGGCACAG-----TGCCACTCT--

2420 2430 2440 2450 2460 2470 2480

Hum. AGACCACCTTTCTCCCCTGAGAGGAGCTTCTGCTACTCTGCATCACTGATGACACTCAGCAGGGTGATGC  
Mur. -GACCA-----GGGTAGGAG--GCT-CT-C-CTGCTA-ACGTGTGTAC-CTACAG-----C

2490 2500 2510 2520 2530 2540 2550

Hum. ACAGCAGTCTG-CCTCCCCATGCGGACTCCCTTCTACCAAGCACATGAGCTCTCTAACAGGGTGGGGGCT  
Mur. ACC-CAGTAGGTCTCTCCCCTGTGGGACTCTCTTCTGC-AAGCACATT-----GGGCT

2560 2570 2580 2590 2600 2610

Hum. ACCCCCAGACCTGCTCCTACACTGATA-TTGAAGAACCTGGAGAGGATCCTTCAGTTCTGGCCATTCCAG  
Mur. GTCTCCATACCTGTACTTGTGCTGTGACAGGAAGAGCCAGAC-AGGTTTCTTTGATTTTGATTGACCCAA

2620 2630 2640 2650 2660 2670 2680

Hum. GGACCCT-CCAGAAACACA-GTGT'TTCAAGAGATCCTAAAAAACCTGCCTGTCCCAGGACCCTATGGTA  
Mur. GAGCCCTGCCTGTAACAAACGTGCTCCAGGAGA-CCATGAAAGTGTGGCTGTCT-GGGATTCTGTGGTG-

2690 2700 2710 2720 2730 2740 2750

Hum. ATGAACACCAAACATCTAAACAATCATATGCTAA-CATGC---CAC--TCCTGGAAACT-CCACTCTGAA  
Mur. ACAAAC-CTAAGCATCCGAGCAAGCTGGGGCTATTCTCTGCAAACCTCCATCCTGAACGCTGTCACTCTAGA

2760 2770 2780 2790 2800 2810

Hum. ---GCTGCCGCTTTGGACACCAACACTCCCTTCT-CCCAGG-GTCATGCAGGGATCTGCTCCCTCCTGC  
Mur. AGCAGCTGCTGCTTTGAACACCAGCCACCCTCCTTCCCAAGAGTCTCTATGGAGTTGGC-CCCTTGTGT

2820 2830 2840 2850 2860 2870 2880

Hum. TTCCCTTACCAGTCGTGCACCGCTGACTCCCAGGAAGTCTTCTCTGAAGTCTGACCACCTTTCTTCTTGC  
Mur. TTCCTTTACCAGTCGGGCCATACTGTTT---GGGAAGTCATCTCTGAAGTCTAACCACCTTCCTTCTTGG

2890 2900 2910 2920 2930 2940 2950

Hum. TTCAGTTGGGGCAGACTCTGATCCCT--TCTGCCCTGGCAGAATGGCAGGGGTAATCTGAGCCTTCTTC  
Mur. TTCAGTTTGGACAGATTGTTATTATTGTCTCTGCCCTGGCTAGAATGGGGGCATAATCTGAGCCTTGTTC

2960 2970 2980 2990 3000 3010

Hum. ACTCCTTTACCC---TAGCTGACCCCTTCACCTCTCCC--CCTCCCTTTTCCTTTGTTTGGGATTTCAGA  
Mur. ---CCTTGTCCAGTGTGGCTGACCC-TTGACCTCTTCCTTCTCTCC---TCCCTTTGTTTGGGATTTCAGA

3020 3030 3040 3050 3060 3070 3080

Hum. AAAGTCTTGTCTCAGAGACTGTTTATTTTTTATTAATAAAGGCTTAAAAAAAAAAAAAAAAAAAAAAAAA  
Mur. AAAGTCTTGTCTCAGAGACAATTTATTTTTTATTAATAA-----AGATATAA

3090 3100

Hum. AAAAAAAGGGCGGCCCGC  
Mur. GCTTTAAAG-----

3140

CHITRO-6572220

Fig. 3I




Fig. 4A

GTCGACCCACGCGTCCGCGGACGCGTGGGGACGGCTCCCGGCTGCAGTCTGCCCCCCCCCGCGCGGGGGCCGAGTC 79

GCGAAGCGCGCCTGCGACCCGGCGTCCGGGCGCGCTGGAGAGGACGCGAGGAGCC	M	R	R	Q	P	A	6
ATG AGG CGC CAG CCT GCG							152
K V A A L L L G L L L E C T E A K K H C							26
AAG GTG GCG GCG CTG CTG CTC GGG CTG CTC TTG GAG TGC ACA GAA GCC AAA AAG CAT TGC							212
W Y F E G L Y P T Y Y I C R S Y E D C C							46
TGG TAT TTC GAA GGA CTC TAT CCA ACC TAT TAT ATA TGC CGC TCC TAC GAG GAC TGC TGT							272
G S R C C V R A L S I Q R L W Y F W F L							66
GGC TCC AGG TGC TGT GTG CGG GCC CTC TCC ATA CAG AGG CTG TGG TAC TTC TGG TTC CTT							332
L M M G V L F C C G A G F F I R R R M Y							86
CTG ATG ATG GGC GTG CTT TTC TGC TGC GGA GCC GGC TTC TTC ATC CGG AGG CGC ATG TAC							392
P P P L I E E P A F N V S Y T R Q P P N							106
CCC CCG CCG CTG ATC GAG GAG CCA GCC TTC AAT GTG TCC TAC ACC AGG CAG CCC CCA AAT							452
P G P G A Q Q P G P P Y Y T D P G G P G							126
CCC GGC CCA GGA GCC CAG CAG CCG GGG CCG CCC TAT TAC ACT GAC CCA GGA GGA CCG GGG							512
M N P V G N S M A M A F Q V P P N S P Q							146
ATG AAC CCT GTC GGG AAT TCC ATG GCA ATG GCT TTC CAG GTC CCA CCC AAC TCA CCC CAG							572
G S V A C P P P P A Y C N T P P P P Y E							166
GGG AGT GTG GCC TGC CCG CCC CCT CCA GCC TAC TGC AAC ACG CCT CCG CCC CCG TAC GAA							632
Q V V K A K *							173
CAG GTA GTG AAG GCC AAG TAG							653

TGGGGTGGCCACGTGCAAGAGGAGAGACAGGAGAGGGCCTTTCCCTGGCCTTTCTGTCTTCGTTGATGTTCACTTCCAG	732
GAACGGTCTCGTGGGCTGCTAAGGGCAGTTCCCTCTGATATCCTCACAGCAAGCACAGCTCTCTTTCAGGCTTTCCATGG	811
AGTACAATATATGAACTCACACTTTGTCTCCTCTGTTGCTTCTGTTTCTGACGCAGTCTGTGCTCTCACATGGTAGTGT	890
GGTGACAGTCCCCGAGGGCTGACGTCCTTACGGTGGCGTGACCAGATCTACAGGAGAGAGACTGAGAGGAAGAAGGCAG	969
TGCTGGAGGTGCAGGTGGCATGTAGAGGGGGCCAGGCCGAGCATCCCAGGCAAGCATCCTTCTGCCCEGGGTATTAATAGG	1048
AAGCCCCATGCCGGGCGGCTCAGCCGATGAAGCAGCAGCCGACTGAGCTGAGCCCAGCAGGTCATCTGCTCCAGCCTGT	1127
CCTCTCGTCAGCCTTCCCTCTTCCAGAAGCTGTTGGAGAGACATTCAGGAGAGAGCAAGCCCCCTTGTTCATGTTTCTGTCT	1206
CTGTTTCATATCCTAAAGATAGACTTCTCCTGCACCGCCAGGGAAGGGTAGCACGTGCAGCTCTCACCGCAGGATGGGGC	1285
CTAGAATCAGGCTTGCCTTGGAGGCCTGACAGTGATCTGACATCCACTAAGCAAATTTATTTAAATTCATGGGAAATCA	1364
CTTCCCTGCCCCAAACTGAGACATTGCATTTTGTGAGCTCTTGGTCTGATTTGGAGAAAGGACTGTTACCCATTTTCTTG	1443
GTGTGTTTATGGAAGTGCATGTAGAGCGTCTGCCCCCTTGAAATCAGACTGGGTGTGTGCTTCCCTGGACATCACTGC	1522
CTCTCCAGGGCATTCTCAGGCCCGGGGTCTCCTTCCCTCAGGCAGCTCCAGTGGTGGGTCTGAAGGGTGCTTTCAAA	1601
ACGGGGCACATCTGGCTGGGAAGTCACATGGACTCTTCCAGGGAGAGAGACCAGCTGAGGCGTCTCTCTCTGAGGTGT	1680

GTTGGGTCTAAGCGGGTGTGTGCTGGGCTCCAAGGAGGAGGAGCTTGCTGGGAAAAGACAGGAGAAGTACTGACTCAAC 1759  
 TGCCTGACCATGTTGTCATAATTAGAATAAAGAAGAAGTGGTCGGAAATGCACATTCCCTGGATAGGAATCACAGCTCA 1838  
 CCCCAGGATCTCACAGGTAGTCTCCTGAGTAGTTGACGGCTAGCGGGGAGCTAGTTCCGCGCATAGTTATAGTGTTGA 1917  
 TGTGTGAACGCTGACCTGTCTGTGTGCTAAGAGCTATGCAGCTTAGCTGAGGCGCCTAGATTACTAGATGTGCTGTAT 1996  
 CACGGGGAATGAGGTGGGGGTGCTTATTTTTTTAATGAACATAATCAGAGCCTCTTGAGAAATTGTTACTCATTGAACTGG 2075  
 AGCATCAAGACATCTCATGGAAGTGGATACGGAGTGATTTGGTGTCATGCTTTTCACTCTGAGGACATTTAATCGGAG 2154  
 AACCTCCTGGGGAATTTTGTGGGAGACACTTGGGAACAAAAACAGACACCCTGGGAATGCAGTTGCAAGCACAGATGCTG 2233  
 CCACCAGTGCTCTGACCACCCTGGTGTGACTGCTGACTGCCAGCGTGGTACCTCCCATGCTGCAGGCCTCCATCTAAA 2312  
 TGAGACAACAAAGCACAAATGTTCACTGTTTACAACCAAGACAACATGCGTGGGTCCAAACACTCCTCTTCTCCAGGTCA 2391  
 TTTGTTTTGCATTTTTTAATGTCTTTATTTTTTTGTAATGAAAAAGCACACTAAGCTGCCCCCTGGAATCGGGTGCAGCTGA 2470  
 ATAGGCACCCAAAAGTCCGTGACTAAATTTTCGTTTGTCTTTTTTGATAGCAAATTATGTTAAGAGACAGTGATGGCTAGG 2549  
 GCTCAACAATTTTGTATTCCCATGTTTGTGTGAGACAGAGTTGTTTTCCCTTGAACCTTGGTTAGAATTGTGCTACTGT 2628  
 GAACGCTGATCCTGCATATGGAAGTCCCACTTTGGTGACATTTCCCTGGCCATTCTTGTTTCCATTGTGTGGATGGTGGG 2707  
 TTGTGCCCCACTTCCTGGAGTGAGACAGCTCCTGGTGTGTAGAATTCCCGGAGCGTCCGTGGTTCAGAGTAAACTTGAAG 2786  
 CAGATCTGTGCATGCTTTTCTCTGCAACAATTGGCTCGTTTCTCTTTTTTGTCTCTTTTGATAGGATCCTGTTTCTCT 2865  
 ATGTGTGCAAAATAAAAATAAATTTGGGCAA 2944  
 AAAAAAAAAAGGGCGGCCGC 2964

Fig. 4B

Fig. 4C

GTCGACCCACGCGTCCGGCCGCGCGTCCTTCTGCCGGCTTCAGCTCGTATCCCCGGAGTCCACCCGCCCCGTCCCGGGGT 79  
 GCGGACTGGCCCTGAGCTGGCCGTACAGCCCCGGCTTCGGACGGTCTCTCGCTGGAGCC ATG GGC CGC CGG CTC 5  
 151  
 G R V A A L L L G L L V E C T E A K K H 25  
 GGC AGG GTG GCG GCG CTG CTG CTC GGG CTG CTA GTG GAG TGC ACT GAG GCC AAA AAA CAT 211  
 C W Y F E G L Y P T Y Y I C R S Y E D C 45  
 TGC TGG TAT TTT GAA GGA CTC TAT CCC ACA TAC TAT ATA TGC CGT TCC TAT GAA GAC TGC 271  
 C G S R C C V R A L S I Q R L W Y F W F 65  
 TGT GGC TCC AGG TGC TGT GTG AGG GCC CTT TCC ATA CAG AGG CTG TGG TAT TTT TGG TTC 331  
 L L M M G V L F C C G A G F F I R R R M 85  
 CTG CTG ATG ATG GGT GTG CTG TTC TGC TGT GGT GCC GGT TTC TTC ATT CGC CGG CGC ATG 391  
 Y P P P L I E E P T F N V S Y T R Q P P 105  
 TAT CCG CCA CCA CTC ATT GAG GAG CCC ACA TTC AAT GTG TCC TAT ACC AGG CAG CCA CCA 451  
 N P A P G A Q Q M G P P Y Y T D P G G P 125  
 AAT CCT GCT CCA GGA GCA CAG CAA ATG GGA CCG CCA TAT TAC ACC GAC CCT GGA GGA CCC 511  
 G M N P V G N T M A M A F Q V Q P N S P 145  
 GGG ATG AAT CCT GTT GGC AAT ACC ATG GCT ATG GCT TTC CAG GTC CAG CCC AAT TCA CCT 571  
 H G G T T Y P P P P S Y C N T P P P P Y 165  
 CAC GGA GGC ACA ACT TAC CCA CCC CCT CCT TCC TAC TGC AAC ACG CCT CCA CCC CCC TAT 631  
 E Q V V K D K \* 173  
 GAA CAG GTG GTG AAG GAC AAG TAG 655  
 CAAGATGCTACATCAAAGGCAAAGAGGATGGACAGGCCCTTTTGTTTACCTTCCCATCCTCACCATACTTGCTGATAG 734  
 GGTGGTCCAAGGGAAACTTGGATATTCTCAAAGCAAGCCCAGCTCTCTTTCAAGTCTTTTGTGGAGGACATTTGAATC 813  
 CACACTGTCTCCTCTGTTGCTTCTGTTTCTGATGTAGTCTGTGCTCTCTGAGAGAGTGTGGCAACAGTCCCTGAGGGTT 892  
 GATATTCTAGGGTGTCCAGGGTAGATCCTCGGGAGAGAGGCTAAGGGGAAAGGAAGGCATAGCCTGTGTGTTAGGGGG 971  
 CAGATAAAGTGGTCAGGCTGAGATAAGACTCACATGATGCAGTAGTTGGCAGTGAACCTCGAAGAGACACTATCCACCA 1050  
 TCCCAGCCCATTCTCCTAATAGAAGCTGTGGGGCTGTGTTGTTGATGCTCTTTGGTCTCCACTCACATTTTGAAAATAG 1129  
 GCTTTCCTCTGCAGGAATAGGAAAGACCCAAGTACATATTTGCTTCCACTTAAAAATGAGGGTCAGAACCAGGCCCTCAG 1208  
 TTGGACATCTATAGTTAAATAAAGGCCATTAGAGAGGGGAAATCTTTAAGTTAGGGGAAATCTCTAAATGGAGACATT 1287  
 GCGTTTTATGAATCATCGTCTGGCTTTTCTTTTAGTGATGTATTGAAGTGAGGGTGTCTTTGAGATCAGATGGGGAG 1366  
 AGTGAACCTCTGCGGGGGGTGGGGTGTCTCTACTCAGAGGGCTCCAACACCCTTTTCTTAGGTAGTTCTGGTGATGGGGTT 1445  
 TTATGGGCACTATAGAGCTGAGGGGCACATTAGGCCGGGTAGTTACATTGACCCTTGGAGAGGAAGAGGACAGCCAAAG 1524

AAACTCAGCAAAGCAAGACCAGCATTGCTGAGTTAGAGCTAGGGTTGTATGTGATCCCAACAGAGATGTGCTGGCCTCA 1603

GAAGAGGGGACGTTTGTGGATAGAGCCGTGAAAACCTACTTAGTTGCACAGATGACATAATCAAAAGTAGAGAAAGAAG 1682

TGTAGTTAGAGATGCCATTTCCCAGGTGAGAATCAGAGCTCATCCATAGATTTACAAGTAGTGGCTGGAGTTAACAGTA 1761

TGGAGTTCTTTTCCCTTGCGTAGTTAGTCACGTTGATGTGTATTTAAACCCAGGTTGAGACCTTGTGTACTAAGAGCAA 1840

GGAAGTATAGCTAAGATGTCTAGATTATTTATATGTAGTATGGTGGGGAGTGGGGCTGCAAGGAAGGGGGCTGACATTG 1919

TAAATGAGAAAATCAGAGCCATTTGATAAACTGTTACTTGTGGATCAGGCATCCAAAAGTGTCTCTTGAGTGGACATT 1998

GAGTATTCTTTTACCACCTACAAGACCAGGAGGCATGGTGTCAATTCTCCATTGGGGTATTTATATGAGGTAGAGGTTTCA 2077

GAATCGACAGTAGCTGTGTGGGCTTAGTTTAAGGACTGAAAGCATAGGGACTGGTAGACAGTTTCATAGGAAACTGCGG 2156

GGAAGGAATGGATACCTTTAAAGACAGTTTGTGGATGCAGATGCTGCCACCCATCATTGAGCACCCCTTGTGTCTCTGGC 2235

TTCCTGTCACTGGATCCAGTACCCCTCCATGCTTGGGTCCCTTGTTTTACATAAGACAACAAAGCACAAATGTCTGCTGTT 2314

TACAATCAAGACGACTACATGGTCCAAACATTTCTTCTCTCTTCTATCACTTGTGGCTTTAACTTCCATTTCCCTCCGTT 2393

CCTTTTAAAAATCAAGAAGCACAGTCAGAGCTGCCCCCTGGGATTGCATCAGGGAACGGCTGATCAAGGCATTCAGTGTC 2472

CATGACTAAATCTTATCTTTTGTAGCAAATCCTTTTAAGAACTGAACAATTGCTAAGGCTCAGCAATTTTATACTC 2551

CAATGTCGTGTAAGGTAAATTTTGTGTTGCCATTGAGCCCACATTGGAATTCCTTCTGACGTCAACACTGACAATGCCT 2630

ATGGAAATTGCACTTCTGGGTATATGTCCCAGCATCCTTGTTTTCTTATGTTTGGTGAGTAAGGCTCACCCCTTCCAGC 2709

AGCTCTACTTCTGTGTGCTGAGGTCCTGTAGAGCCGGGGCTTGGGCACAGACATGAGGCAGACTTGTGCATGCTCTTTC 2788

TTGGCAACACTTGGCTCATATTTCTTGTCTCTTTTGTAGAGTCCTGTTTCTTATGTATTTAAAAAATAATAAAAGTG 2867

AAATTTAGTCAAAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCCGC 2915

Fig. 4D

	10	20	30	40	50	60	70
<i>Hum.</i>	MRRQPAKVAALLLGLLLECTEAKKHCWYFEGLYPTYIICRSYEDCCGSRCCVRALSIQRLWYFWFLLMMG						
	:	:	:	:	:	:	:
<i>Mur.</i>	MGRRLGRVAALLLGLLVECTEAKKHCWYFEGLYPTYIICRSYEDCCGSRCCVRALSIQRLWYFWFLLMMG						
	10	20	30	40	50	60	70
	80	90	100	110	120	130	140
<i>Hum.</i>	VLFCGAGFFIRRRMYPPPLIEEPAFNVSYTRQPPNPQGAQQPGPPYYTDPGGPGMNPVGNMAMAFQV						
	:	:	:	:	:	:	:
<i>Mur.</i>	VLFCGAGFFIRRRMYPPPLIEEPTFNVSYTRQPPNPAPGAQQMGPPYYTDPGGPGMNPVGNMTAMAFQV						
	80	90	100	110	120	130	140
	150	160	170				
<i>Hum.</i>	PPNSPQGSVACPPPPAYCNTFPFPPYEQVVAK						
	:	:	:	:	:	:	:
<i>Mur.</i>	QPNSPHGGTTYPPPPSYCNTFPFPPYEQVVKDK						
	150	160	170				

Fig. 4F





Fig. 4G



## Fig. 5A

GTCGACCCACGCGTCCGCAGCTTTGGACACTTCCTCTGCTTGAGGACACCTTGACTAACCTCCAAGGGCAACTAAAGGA 79

M C T K T I 6

TCAAGAAAGGCCCAGCACAGCAGAAGATCAGCTGGATCTAGCTCCTGCAGGAG ATG TGT ACA AAG ACA ATC 150

P V L W G C F L L W N L Y V S S S Q T I 26  
CCA GTC CTC TGG GGA TGT TTC CTC CTG TGG AAT CTC TAT GTC TCA TCC TCT CAG ACC ATT 210

Y P G I K A R I T Q R A L D Y G V Q A G 46  
TAC CCT GGA ATC AAG GCA AGG ATT ACT CAG AGG GCA CTT GAC TAT GGT GTT CAA GCT GGA 270

M K M I E Q M L K E K K L P D L S G S E 66  
ATG AAG ATG ATT GAG CAA ATG CTA AAA GAA AAG AAA CTC CCA GAT TTA AGC GGT TCT GAG 330

S L E F L K V D Y V N Y N F S N I K I S 86  
TCT CTT GAA TTT CTA AAA GTT GAT TAT GTA AAC TAC AAT TTT TCA AAT ATA AAA ATC AGT 390

A F S F P N T S L A F V P G V G I K A L 106  
GCC TTT TCA TTT CCA AAT ACC TCA TTG GCT TTT GTG CCT GGA GTG GGA ATC AAA GCG CTA 450

T N H G T A N I S T D W G F E S P L F V 126  
ACC AAC CAT GGC ACT GCC AAC ATC AGC ACA GAC TGG GGG TTC GAG TCT CCA CTT TTT GTT 510

L Y N S F A E P M E K P I L K N L N E M 146  
GTG TAT AAC TCC TTT GCT GAG CCC ATG GAG AAA CCC ATT TTA AAG AAC TTA AAT GAA ATG 570

L C P I I A S E V K A L N A N L S T L E 166  
CTC TGT CCC ATT ATT GCA AGT GAA GTC AAA GCG CTA AAT GCC AAC CTC AGC ACA CTG GAG 630

V L T K I D N Y T L L D Y S L I S S P E 186  
GTT TTA ACC AAG ATT GAC AAC TAC ACT CTG CTG GAT TAC TCC CTA ATC AGT TCT CCA GAA 690

I T E N Y L D L N L K G V F Y P L E N L 206  
AAT ACT GAG AAC TAC CTT GAC CTG AAC TTG AAG GGT GTA TTC TAC CCA CTG GAA AAC CTC 750

T D P P F S P V P F V L P E R S N S M L 226  
ACC GAC CCC CCC TTC TCA CCA GTT CCT TTT GTG CTC CCA GAA CGC AGC AAC TCC ATG CTC 810

Y I G I A E Y F F K S A S F A H F T A G 246  
TAC ATT GGA ATC GCC GAG TAT TTC TTT AAA TCT GCG TCC TTT GCT CAT TTC ACA GCT GGG 870

V F N L T L S T E E I S N H F V Q N S Q 266  
GTT TTC AAT CTC ACT CTC TCC ACC GAA GAG ATT TCC AAC CAT TTT GTT CAA AAC TCT CAA 930

G L G N V L S R I A E I Y I L S Q P F M 286  
GGC CTT GGC AAC GTG CTC TCC CGG ATT GCA GAG ATC TAC ATC TTG TCC CAG CCC TTC ATG 990

V R I M A T E P P I I N L Q P G N F T L 306  
GTG AGG ATC ATG GCC ACA GAG CCT CCC ATA ATC AAT CTA CAA CCA GGC AAT TTC ACC CTG 1050

D I P A S I M M L T Q P K N S T V E T I 326  
GAC ATC CCT GCC TCC ATC ATG ATG CTC ACC CAA CCC AAG AAC TCC ACA GTT GAA ACC ATC 1110

V S M D F V A S T S V G L V I L G Q R L 346

GTT TCC ATG GAC TTC GTT GCT AGT ACC AGT GTT GGC CTG GTT ATT TTG GGA CAA AGA CTG	1170
V C S L S L N R F R L A L P E S N R S N	366
GTC TGC TCC TTG TCT CTG AAC AGA TTC CGC CTT GCT TTG CCA GAG TCC AAT CGC AGC AAC	1230
I E V L R F E N I L S S I L H F G V L P	386
ATT GAG GTC TTG AGG TTT GAA AAT ATT CTA TCG TCC ATT CTT CAC TTT GGA GTC CTC CCA	1290
L A N A K L Q Q G F P L P N P H K F L F	406
CTG GCC AAT GCA AAA TTG CAG CAA GGA TTT CCT CTG CCC AAT CCA CAC AAA TTC TTA TTC	1350
V N S D I E V L E G F L L I S T D L K Y	426
GTC AAT TCA GAT ATT GAA GTT CTT GAG GGT TTC CTT TTG ATT TCC ACC GAC CTG AAG TAT	1410
E T S S K Q Q P S F H V W E G L N L I S	446
GAA ACA TCC TCA AAG CAG CAG CCA AGT TTC CAC GTA TGG GAA GGT CTG AAC CTG ATA AGC	1470
R Q W R G K S A P *	456
AGA CAG TGG AGG GGG AAG TCA GCC CCT TGA	1500
TTGCCGGTTTGC AATTCACCCCAGGAAGTAAATGGTCCTTAATCCTACA ACTACTGTAAACCCAGAAGGGAAAGACAGT	1579
AGACACTGGAATTGTAAAGCCCTTGTGAATTGCTTAGGCAGAAAGTTTCTTTCTTAAGCCTTCAGGAACCCAGAATAA	1658
GGCAGACTCTGTTAAAGGGATAAATAGAGGTGTCTGAATGTGAGTGTATGCATGCTGCGTGTGTCTGTGTTTATGTTTG	1737
TTTGTTTGTGTTGGGGCAAGAAAGATTCTAGGACAAGAGCTAGGCATGTACTTCTGACCAGGTGGGTAAGCAACTCTAAG	1816
TCTGTATTTGTATTGGTCATTCTCAGTGGAATCCCTTAGGCCCTCTAGTGTTTCCCCTACCTGCATATTGGTTTTTC	1895
ATGTTTTATATTCACTGTTACTATCTTCTGTGTTTAATTAAAATTGTTTTCTATCAAAAAAAAAAAAAAAAAAAGGGC	1974
GGCCGC	1980

Fig. 5B



```

10      20      30      40      50      60
286 MCTKT-IPVLWGCFL-LWNLYVSSSQTIYPGIKARITQRALDYGVQAGMKMIEQMLKEKKLPDLGSES
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI MARGPCNAPRWVSLMVLVAIGTAVTAAVNPGVVVRISQKGLDYASQQGTAALQKELKRIKIPDYS--DSF
      10      20      30      40      50      60

70      80      90      100      110      120      130
286 EFLKVDYVNYNFSNIKISAFSFPNTSLAFVPGVGIKALTNHGTANISTDWGFESPLFVLYNSFAEPME-
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI KIKHLGKGHYSFYSDIREFQLPSSQISMVFNVLKFSISNANIKISGKWKAAQKRFLKMSGNFDLISIEGM
70      80      90      100      110      120      130

140      150
286 -----KPI-----
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI SISADLKLGSNPTSGKPTITCSSSHINSVHVHISKSKVGLIQLFHKKIESALRNKMNSQVCEKVTNS
140      150      160      170      180      190      200

160      170      180      190      200      210      220
286 VKA-LNANLSTLEVLTKIDNYTLDDYSLISSPEITENYLDLNLKGVFYPLENLTDPFSPVPFVLPERSN
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI VSSKLQPYFQTLPMVKIDSVAGINYGLVAPPATTAETLDVOMKGEFYSENHHNPPPFAPPVMEFPAHD
210      220      230      240      250      260      270

230      240      250      260      270      280      290
286 SMLYIGIAEYFFKSASFHFTAGVFNLTSTEEISNH--FVQNSQGLGNVLSRIAETIYILSQPFMVRIMA
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI RMVYLGLSDYFFNTAGLVYQEAGVLKMTLRDDMIPKESKFRLTTKFFGTFLPEVAKKFP-NMKIQIHVSA
280      290      300      310      320      330      340

300      310      320      330      340      350      360
286 TEPPILNLPQGNFTLDIPASIMMLTQPKNSTVETIVSMDFVASTSVGLVILGQRLVCSLSLNRFRALPE
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI STPPHLSVQPTGLTFYPAVDVQAFVLPNSSLASLFLIGMHTTGSMEVSAESNRLVGELKLDRLLELKH
350      360      370      380      390      400      410

370      380      390      400      410      420      430
286 SNRSNIEVLRFENILSSILHFGVLPLANAKLQQGFPLPNPHKFLFVNSDIEVLEGFLLISTDLKYETSSK
:      :      :      :      :      :      :      :      :      :      :      :      :      :      :
BPI SNIGPFPVELLQDIMNYIVPILVLPVNEKLQKGFPLPTPARVQLYNVVLQPHQNFLLFGADVYK----
420      430      440      450      460      470      480

440      450
286 QQPSFHVWEGNLNISRQWRGKSAP -

```

Fig. 5E

**286** MCTKTIPVLWGCFLLWNLYVSSSQTI--YPGIKARITQRALDYGVOAGMKMIEQMLKEKKLPDLSGSESL  
:  
**REN** MGALARAL--PSILLALLLTSTPEALGANPGLVARITDKGLQYAAQEGLLALQSELLRITLPDFTG--DL  
10 20 30 40 50 60

286 EFLKVDYVNYNFSNIKISAFSFPNTSLAFVPGVGIKALTNHGTANISTDWGFESPLFVLYNSFAEPME--  
 . . . : : . : : . : : . : : . : : . : : . : : . : : . : : . : : . : : . : : . : : . : :  
 RNP RIPHVGRGRYEFHSLNIHEFQLPSSQISMVNPVGLKFSISNANIKISGKWAQKRFLKMSGNFDLSIEGM  
 70 80 90 100 110 120 130

286 -----KPI-----LKN-LNEMLCPIIASE  
RENPD ::::: :::  
SISADLKLGSNPTSGKPTITCSSCSSHINSVHVHISKSKVGLIQLFHKKIESALRNKMNSQVCEKVTNS  
140 150 160 170 180 190 200

286 VKA-LNANLSTLEVLTKIDNYTLLDYSLISSPEITENYLDNLKGVFYPLENLTDPFSPVPFVLPERSN  
:  
REN P VSSKLQPYFQTLPMVTKIDSVAGINYGLVAPPATTAETLDVQMGEFYSENHHNPPPFAPPVMEFPAHD  
210 220 230 240 250 260 270

286 SMLYIGIAEYFFKSASFHFTAGVFNLTLSTEEISNH--FVQNSQGLGNVLSRIAETIYILSQPFMVRIMA  
 RNP RMVYLGLSDYFFNTAGLVYQEAGVLKMTLRDDMIPKESKFRLTTKFFGTFLPEVAKKFP-NMKIQIHVSA  
 280 290 300 310 320 330 340

**Z66** TEPIINLQPGNFTLDIPASIMMLTQPKNSTVETIVSMDFVASTSVGLVILGQRLVCSSLNRFRALPE  
**REP** . :  
 STPHLSVQPTGLTFYPAVDVQALAVLPNSSLASFLIGMHTTGSMFVSANRNLVGELKLDRIILLKH  
 == 350 360 370 380 390 400 410

**REN** SNRSNIEVLR FENILSSILHFGVLPLANAKLQQGFPLPNPHKFLFVNSDIEVLEGFLLISTDLKYETSSK  
:: :: : . . . . . :  
SNIGPFPVELLQDIMNYIVPILVLPVNEKLQKGFPLPTPARVOLYNVVLQPHQNFLFGADVYK----

286 Q Q P S F H V W E G L N L I S R Q W R G K S A P -

## Fig. 6A

GTCGACCCACGCGTCCGGGGAATTGCAGCAGGAAAAATATGTGAAGAGTTTTTAAACCCACAAATTCTTCTTACTTTAGA 79  
 M L E T L S R Q 8  
 ATTAGTTGTTACATTGGCAGGAAAAAATAAATGCAGATGTTGGACC ATG TTG GAA ACC TTG TCA AGA CAG 149  
 W I V S H R M E M W L L I L V A Y M F Q 28  
 TGG ATT GTC TCA CAC AGA ATG GAA ATG TGG CTT CTG ATT CTG GTG GCG TAT ATG TTC CAG 209  
 R N V N S V H M P T K A V D P E A F M N 48  
 AGA AAT GTG AAT TCA GTA CAT ATG CCA ACT AAA GCT GTG GAC CCA GAA GCA TTC ATG AAT 269  
 I S E I I Q H Q G Y P C E E Y E V A T E 68  
 ATT AGT GAA ATC ATC CAA CAT CAA GGC TAT CCC TGT GAG GAA TAT GAA GTC GCA ACT GAA 329  
 D G Y I L S V N R I P R G L V Q P K K T 88  
 GAT GGG TAT ATC CTT TCT GTT AAC AGG ATT CCT CGA GGC CTA GTG CAA CCT AAG AAG ACA 389  
 G S R P V V L L Q H G L V G G A S N W I 108  
 GGT TCC AGG CCT GTG GTG TTA CTG CAG CAT GGC CTA GTT GGA GGT GCT AGC AAC TGG ATT 449  
 S N L P N N S L G F I L A D A G F D V W 128  
 TCC AAC CTG CCC AAC AAT AGC CTG GGC TTC ATT CTG GCA GAT GCT GGT TTT GAC GTG TGG 509  
 M G N S R G N A W S R K H K T L S I D Q 148  
 ATG GGG AAC AGC AGG GGA AAC GCC TGG TCT CGA AAA CAC AAG ACA CTC TCC ATA GAC CAA 569  
 D E F W A F S Y D E M A R F D L P A V I 168  
 GAT GAG TTC TGG GCT TTC AGT TAT GAT GAG ATG GCT AGG TTT GAC CTT CCT GCA GTG ATA 629  
 N F I L Q K T G Q E K I Y Y V G Y S Q G 188  
 AAC TTT ATT TTG CAG AAA ACG GGC CAG GAA AAG ATC TAT TAT GTC GGC TAT TCA CAG GGC 689  
 T M G F I A F S T M P E L A Q K I K M 208  
 ACC ACC ATG GGC TTT ATT GCA TTT TCC ACC ATG CCA GAG CTG GCT CAG AAA ATC AAA ATG 749  
 V F A L A P I A T V K H A K S P G T K F 228  
 TAT TTT GCT TTA GCA CCC ATA GCC ACT GTT AAG CAT GCA AAA AGC CCC GGG ACC AAA TTT 809  
 L L L P D M M I K G L F G K K E F L Y Q 248  
 TTG TTG CTG CCA GAT ATG ATG ATC AAG GGA TTG TTT GGC AAA AAA GAA TTT CTG TAT CAG 869  
 T R F L R Q L V -I Y L C G Q V I L D Q I 268  
 ACC AGA TTT CTC AGA CAA CTT GTT ATT TAC CTT TGT GGC CAG GTG ATT CTT GAT CAG ATT 929  
 C S N I M L L L G G F N T N N M N M S R 288  
 TGT AGT AAT ATC ATG TTA CTT CTG GGT GGA TTC AAC ACC AAC AAT ATG AAC ATG AGC CGA 989  
 A S V Y A A H T L A G T S V Q N I L H W 308  
 GCA AGT GTA TAT GCT GCC CAC ACT CTT GCT GGA ACA TCT GTG CAA AAT ATT CTA CAC TGG 1049  
 S Q A V N S G E L R A F D W G S E T K N 328  
 AGC CAG GCA GTG AAT TCT GGT GAA CTC CGG GCA TTT GAC TGG GGG AGT GAG ACC AAA AAT 1109  
 L E K C N Q P T P V R Y R V R D M T V P 348

CTG GAA AAA TGC AAT CAG CCA ACT CCT GTA AGG TAC AGA GTC AGA GAT ATG ACG GTC CCT	1169
T A M W T G G Q D W L S N P E D V K M L	368
ACA GCA ATG TGG ACA GGA GGT CAG GAC TGG CTT TCA AAT CCA GAA GAC GTG AAA ATG CTG	1229
L S E V T N L I Y H K N I P E W A H V D	388
CTC TCT GAG GTG ACC AAC CTC ATC TAC CAT AAG AAT ATT CCT GAA TGG GCT CAC GTG GAT	1289
F I W G L D A P H R M Y N E I I H L M Q	408
TTC ATC TGG GGT TTG GAT GCT CCT CAC CGT ATG TAC AAT GAA ATC ATC CAT CTG ATG CAG	1349
Q E E T N L S Q G R C E A V L *	424
CAG GAG GAG ACC AAC CTT TCC CAG GGA CGG TGT GAG GCC GTA TTG TGA	1397
AGCATCTGACACTGACGATCTTAGGACAACCTCCTGAGGGATGGGGCTAGGACCCATGAAGGCAGAATTACGGAGAGCA	1476
GAGACCTAGTATACATTTTTTCAGATTCCCTGCACTTGGCACTAAATCCGACACTTACATTTACATTTTTTTTCTGTAAA	1555
TTAAAGTACTTATTAGGTAAATAGAGGTTTTGTATGCTATTATATATTCTACCATCTTGAAGGGTAGGTTTTTACCTGAT	1634
AGCCAGAAAATATCTAGACATTCTCTATATCATTCAGGTAAATCTCTTTAAAACACCTATTGTTTTTTCTATAAGCCAT	1713
ATTTTTGGAGCACTAAAGTAAATGGCAAATTGGGACAGATATTGAGGTCTGGAGTCTGTGGATTATTGTTGACTTTGA	1792
CAAAATAAGCTAGACATTTTCACCTTGTTGCCACAGAGACATAACACTACCTCAGGAAGCTGAGCTGCTTTAAGGACAA	1871
CAACAACAAAATCAGTGTTACAGTATGGATGAAATCTATGTTAAGCATTCTCAGAATAAGGCCAAGTTTTATAGTTGCA	1950
TCTCAGGGAAGAAAATTTTATAGGATGTTTATGAGTTCTCCAATAAATGCATTCTGCATTACATAAAAAAAAAAAAAAA	2029
AAAAAGGGCGGCCGC	2044

Fig. 6B

Fig. 6C

294 10 20 30 40 50 60 70  
 MLETL SRQWIVSHR MEMWLLILVAYMFQRNVNSVHMPTKAVDPEAFMNISEIIQHOGYPCEEYEVEDG  
 HLP M-----WLL---LTMASLISVLGTTGHLFGKLH---PGSPEVTMNISQMITYWGYPNEEYEVVTEDEG  
 10 20 30 40 50

294 80 90 100 110 120 130 140  
 YILSVNRIPRGLVQPKKTGSRPVVLLQHGLVGGASNWISNLPNNSLGFILADAGFDVWGMNSRGNAWSRK  
 HLP YILEVNRIPYGGKNSGNTGQRPVVFLQHGLLASATNWNISNLPNNSLAFILADAGYDVWLGNSRGNTWARR  
 60 70 80 90 100 110 120

294 150 160 170 180 190 200 210  
 HKTL SIDQDEFWAFSYDEMARFDLPAVINFILQKTGQEKIYYVGYSQGT TMGFIAFSTMP ELAQKIKMYF  
 HLP NLYYSPDSVEFWAFSFD EMAKYDLPATIDFIVKKTGQKQLHYVGHSQGT TIGFIAFSTNPSLAKRIKTFY  
 130 140 150 160 170 180 190


294 220 230 240 250 260 270  
 ALAPIATVKHAKSPGTFLLLPDMMIKGLFGKKEFLYQTRFLRQ-LVIYLCGQVILDQICSNIMLLGGF  
 HLP ALAPVATVKYTKSLINKLRFPVQSLFKFIFGDKIF-YPHNFFDQFLATEVCSREMLNLLCSNALFIICGF  
 200 210 220 230 240 250 260

294 280 290 300 310 320 330 340  
 NTNMMNMSRASVYAAHTLAGTSVQNILHWSQAVNSGELRAFDWGSETKNLEKCNQPTPVRYRVRDMTVPT  
 HLP DSKNFNTSRLDVYL SHNPAGTSVQNMFWHTQAVKSGKFQAYDWGSPVQNRMHYDQSQPPYYNVTAMNVP  
 270 280 290 300 310 320 330

294 350 360 370 380 390 400 410  
 AMWTGGQDWLSNPEDVKMLLSEVTNLIYHKNIPEWAHVDFIWGLDAPHRMYNEIIHLMQOEETNLSQGRG  
 HLP AVWNGGKDLLADPDVGLLLPKLPNLIYHKEIPFYNHLDFIWAMDAPQEVYNDIVSMISEDKK-----  
 340 350 360 370 380 390

420  
 294 EAVL  
 HLP ----

Fig. 6D



001630-6572260

[illegible]


Fig. 7A

GTCGACCCACGCGTCCACGGCGAGGGCTCCCGGGGCGCAGCATTGCCCCCCTGCACCACCTCACCAAG																			M A	2
ATG GCT																				75
T	L	G	H	T	F	P	F	Y	A	G	P	K	P	T	F	P	M	D	T	22
ACT	TTG	GGA	CAC	ACA	TTC	CCC	TTC	TAT	GCT	GGC	CCC	AAG	CCA	ACC	TTC	CCG	ATG	GAC	ACC	135
T	L	A	S	I	I	M	I	F	L	T	A	L	A	T	F	I	V	I	L	42
ACT	TTG	GCC	AGC	ATC	ATC	ATG	ATC	TTT	CTG	ACT	GCA	CTG	GCC	ACG	TTC	ATC	GTC	ATC	CTG	195
P	G	I	R	G	K	T	R	L	F	W	L	L	R	V	V	T	S	L	F	62
CCT	GGC	ATT	CGG	GGA	AAG	ACG	AGG	CTG	TTC	TGG	CTG	CTT	CGG	GTG	GTG	ACC	AGC	TTA	TTC	255
I	G	A	A	I	L	A	V	N	F	S	S	E	W	S	V	G	Q	V	S	82
ATC	GGG	GCT	GCA	ATC	CTG	GCT	GTG	AAT	TTC	AGT	TCT	GAG	TGG	TCT	GTG	GGC	CAG	GTC	AGC	315
T	N	T	S	Y	K	A	F	S	S	E	W	I	S	A	D	I	G	L	Q	102
ACC	AAC	ACA	TCA	TAC	AAG	GCC	TTC	AGT	TCT	GAG	TGG	ATC	AGC	GCT	GAT	ATT	GGG	CTG	CAG	375
V	G	L	G	G	V	N	I	T	L	T	G	T	P	V	Q	Q	L	N	E	122
GTC	GGG	CTG	GGT	GGA	GTC	AAC	ATC	ACA	CTC	ACA	GGG	ACC	CCC	GTG	CAG	CAG	CTG	AAT	GAG	435
T	I	N	Y	N	E	E	F	T	W	R	L	G	E	N	Y	A	E	E	C	142
ACC	ATC	AAT	TAC	AAC	GAG	GAG	TTC	ACC	TGG	CGC	CTG	GGT	GAG	AAC	TAT	GCT	GAG	GAG	TGT	495
A	K	A	L	E	K	G	L	P	D	P	V	L	Y	L	A	E	K	F	T	162
GCA	AAG	GCT	CTG	GAG	AAG	GGG	CTG	CCA	GAC	CCT	GTG	TTG	TAC	CTA	GCT	GAG	AAG	TTC	ACT	555
P	R	S	P	C	G	L	Y	R	Q	Y	R	L	A	G	H	Y	T	S	A	182
CCA	AGA	AGC	CCA	TGT	GGC	CTA	TAC	CGC	CAG	TAC	CGC	CTG	GCG	GGA	CAC	TAC	ACC	TCA	GCC	615
M	L	W	V	A	F	L	C	W	L	L	A	N	V	M	L	S	M	P	V	202
ATG	CTA	TGG	GTG	GCA	TTC	CTC	TGC	TGG	CTG	CTG	GCC	AAT	GTG	ATG	CTC	TCC	ATG	CCT	GTG	675
L	V	Y	G	G	Y	M	L	L	A	T	G	I	F	Q	L	L	A	L	L	222
CTG	GTA	TAT	GGT	GGC	TAC	ATG	CTA	TTG	GCC	ACG	GGC	ATC	TTC	CAG	CTG	TTG	GCT	CTG	CTC	735
F	F	S	M	A	T	S	L	T	S	P	C	P	L	H	L	G	A	S	V	242
TTC	TTC	TCC	ATG	GCC	ACA	TCA	CTC	ACC	TCA	CCC	TGT	CCC	CTG	CAC	CTG	GGC	GCT	TCT	GTG	795
L	H	T	H	H	G	P	A	F	W	I	T	L	T	T	G	L	L	C	V	262
CTG	CAT	ACT	CAC	CAT	GGG	CCT	GCC	TTC	TGG	ATC	ACA	TTG	ACC	ACA	GGA	CTG	CTG	TGT	GTG	855
L	L	G	L	A	M	A	V	A	H	R	M	Q	P	H	R	L	K	A	F	282
CTG	CTG	GGC	CTG	GCT	ATG	GCG	GTG	GCC	CAC	AGG	ATG	CAG	CCT	CAC	AGG	CTG	AAG	GCT	TTC	915
F	N	Q	S	V	D	E	D	P	M	L	E	W	S	P	E	E	G	G	L	302
TTC	AAC	CAG	AGT	GTG	GAT	GAA	GAC	CCC	ATG	CTG	GAG	TGG	AGT	CCT	GAG	GAA	GGT	GGA	CTC	975
L	S	P	R	Y	R	S	M	A	D	S	P	K	S	Q	D	I	P	L	S	322
CTG	AGC	CCC	CGC	TAC	CGG	TCC	ATG	GCT	GAC	AGT	CCC	AAG	TCC	CAG	GAC	ATT	CCC	CTG	TCA	1035
E	A	S	S	T	K	A	Y	C	K	E	A	H	P	K	D	P	D	C	A	342
GAG	GCT	TCC	TCC	ACC	AAG	GCA	TAC	TGT	AAG	GAG	GCA	CAC	CCC	AAA	GAT	CCT	GAT	TGT	GCT	1095

L	*		
TTA	TAA		344
			1101
CATTCTCCCCGTGGAGGCCACCTGGACTTCCAGTCTGGCTCCAAACCTCATTGGCGCCCCATAAAACCAGCAGAACTG		1180	
CCCTCAGGGTGGCTGTTACCAGACACCCAGCACCAATCTACAGACGGAGTAGAAAAAGGAGGCTCTATATACTGATGTT		1259	
AAAAAACAAAAACAAAAACAAAAAGCCCTAAGGGACTGAAGAGATGCTGGGCCTGTCCATAAAGCCTGTTGCCATGATAAG		1338	
GCCAAGCAGGGGCTAGCTTATCTGCACAGCAACCCAGCCTTTCCGTGCTGCCTTGCCCTCTTCAAGATGCTATTCACTGA		1417	
AACCTAACTTCACCCCCATAACACCAGCAGGGTGGGGGTTACATATGATTCTCCTATGGTTTCCTCTCATCCCTCGGCA		1496	
CCTCTTGTTTTCTCTTTTCCTGGGTTCTTTTGTCTTCCTTTACTTCTCCAGCTTGTGTGGCCTTTTGGTACAATGAA		1575	
AGACAGCACTGGAAAGGAGGGGAAACCAAACCTTCTCATCCTAGGTCTAACATTAACCAACTATGCCACATTCTCTTTGA		1654	
GCTTCAGTTFCCCAAATTTGCTACATAAGATTGCAAGACTTGCCAAGAATCTTGGGATTTATCTTTCTATGCCTTGCTGA		1733	
CACCTACCTTGGCCCTCAAACACCACCTCACAAGAAGCCAGGTGGGAAGTTAGGGAATCAACTCCAAAACGCTATTCTT		1812	
TCCCACCCCACTCAGCTGGGCTAGCTGAGTGGCATCCAGGACGGGGGAGTGGGTGACCTGCCTCATCACTGCCACCTAA		1891	
CGTCCCCCTGGGGTGGTTCAGAAAGATGCTAGCTCTGGTAGGGTCCCTCCGGCCTCACTAGAGGGCGCCCTATTACTC		1970	
TGGAGTCGACGCAGAGAATCAGGTTTCACAGCACTGCGGAGAGTGTACTAGGCTGTCTCCAGCCCAGCGAAGCTCATGA		2049	
GGACGTGCGACCCCGGCGGAGAAGCCATGAAAATTAATGGGAAAAACAGTTTTTAAAAAAAAAAAAAAAAAAGGGCG		2128	
GGCGC		2133	

Fig. 7B

Fig. 7C



```

296      10      20      30      40      50      60      70
CRP MATLGHTFFPYAGPKPTFFPMDTTLASIIIMIFLTALATFIVILPGIRGKTRLFWLLRVVTSLSFIGAAILAV
:  . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . :
M-RIAH-----ASSRGN-----SIFSVFLIPLIAYILILPGVR-RKRVVTTVTYVLMMLAVGGALIAS
              10              20              30              40              50

      80      90      100      110      120      130
296 NFSSEWSVGQVSTNTSYKAFSSEWISADIGLQVGLGGVNITL-----TGTPVQQLNETIN--YNEEFTW
:  . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . :
CRP LIYPCWASGSQMIYTQFRGHSNERILAKIGVEIGLQKVNVTLKFERLLSSNDVLPGSMDTELYYNEGFDI
      60      70      80      90      100      110      120

      140      150      160      170      180      190      200
296 RLGENYAECEKALEKGLPDPVLYLAEKFT-PRSPCGLYRQYRLAGHYTSAMLWVAFLCWLLANV-MLSM
:  . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . :
CRP SGISSMAEALHHGLENGLPYPMLSVLEYFSLNQDSFDWGRHYRVAGHYTHAAIWFAFACWCLSVVLMFL
      130      140      150      160      170      180      190

      210      220      230      240      250      260
296 PVLVYGGYMLLATGIFQLLALLFFSMATSLTSPCPLHL---GASVLHTHHGPAF---WITLTTGLLCVL
:  . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . :
CRP PHNAYKS--ILATGISCLIACLVL---LLSPCELRIAFGTGENFERVDLTATFSFCFYLFIFAIGILCVL
      200      210      220      230      240      250      260

      270      280      290      300      310      320
296 LGLAMAVAHRMQPHRLKAFFNQSVDEDPMLEW-----SPEEGLLSPRY--RSMADSPKSDIPLSEAS
:  . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . :
CRP CGLGLGICEHWRIYTLSTFLDASLDEHVGPWKKLPTGGPALQGVQIGAYGTNTTNSSRDKNDISSDKTA
      270      280      290      300      310      320      330

      330      340
296 STKAY-----CK-----EAHPKDPD-----CA---L
:  . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . : . . . :
CRP GSSGFQSRSTSTCQSSASSASLSRQSSIETVHDEAELERTHVHFLQEPCCSSST
      340      350      360      370      380

```

**DECLARATION AND POWER OF ATTORNEY**

(Original Application)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**NOVEL GENES ENCODING PROTEINS HAVING DIAGNOSTIC,  
PREVENTIVE, THERAPEUTIC, AND OTHER USES**

the specification of which is attached hereto and/or was filed on June 14, 1999 as Application No. \_\_\_\_\_.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to herein.

I acknowledge the duty to disclose information which is material to patentability in accordance with Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d), of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

**FOREIGN PRIORITY APPLICATION(S)**

**Priority Claimed**

☐ Yes ☐ No

\_\_\_\_\_  
(Number) (Country) (Day/month/year filed)

☐ Yes ☐ No

\_\_\_\_\_  
(Number) (Country) (Day/month/year filed)

I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional patent application(s) listed below and have also identified below any United States provisional patent application(s) having a filing date before that of the application on which priority is claimed:

**PROVISIONAL PRIORITY PATENT APPLICATION(S)**

**Priority Claimed**

☐ Yes ☐ No

\_\_\_\_\_  
(Application No.) (Filing Date)

☐ Yes ☐ No

\_\_\_\_\_  
(Application No.) (Filing Date)

And I hereby appoint the registered attorneys and agents associated with **Panitch Schwarze Jacobs & Nadel, P.C., Customer No. 000570**, as my attorneys or agents with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

Address all correspondence to **Customer No. 000570, namely, PANITCH SCHWARZE JACOBS & NADEL, P.C.**, One Commerce Square, 2005 Market Street, 22nd Floor, Philadelphia, Pennsylvania 19103-7086. Please direct all communications and telephone calls to **Gary D. Colby, Ph.D., J.D.** at 215-965-1285.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full name of sole  
or first inventor Sean A. McCarthy

Inventor's Signature \_\_\_\_\_

Date \_\_\_\_\_

Residence Boston, Massachusetts

Citizenship United Kingdom

Post Office Address 62 Commonwealth Avenue, #4

Boston, MA 02116

Full name of second joint  
inventor, if any Christopher C. Fraser

Inventor's Signature \_\_\_\_\_

Date \_\_\_\_\_

Residence Lexington, Massachusetts

Citizenship United States of America

Post Office Address 52 Grassland Street

Lexington, MA 02421

Full name of third joint  
inventor, if any John D. Sharp

Inventor's Signature \_\_\_\_\_

Date \_\_\_\_\_

Residence Arlington, Massachusetts

Citizenship United States of America

Post Office Address 245 Park Avenue

Arlington, MA 02476

Full name of forth joint  
inventor, if any Thomas S. Barnes

Inventor's Signature \_\_\_\_\_

Date \_\_\_\_\_

Residence Boston, Massachusetts

Citizenship Australia

Post Office Address 22 Hanson Street #2

Boston, MA 02118

PSJN2/236803.1

# SEQUENCE LISTING

<110> McCarthy, Sean A  
Fraser, Christopher C  
Sharp, John D  
Barnes, Thomas M

<120> NOVEL GENES ENCODING PROTEINS HAVING DIAGNOSTIC,  
PREVENTIVE, THERAPEUTIC, AND OTHER USES

<130> 10147-6

<140> Not Yet Assigned

<141> 1999-06-14

<160> 79

<170> PatentIn Ver. 2.0

<210> 1

<211> 1656

<212> DNA

<213> Homo sapiens

<400> 1

```
gtcgacccac gcggtccgcc acggtccgg cccatggcgc cgcgcgcgc cgcctcgcc 60
ctgctctccg ccgcggcgct cacgtggcg gcccggcccg cgcctagccc cggcctcgcc 120
cccgaccccg agtgtttcac agccaatggt gcggattata ggggaacaca gaactggaca 180
gcactacaag gcgggaagcc atgtctgttt tggaacgaga ctttcagca tccatacaac 240
actctgaaat accccaacgg ggaggggggc ctgggtgagc acaactattg cagaaatcca 300
gatggagacg tgagcccctg gtgctatgtg gcagagcagc aggatgggtg ctactggaag 360
tactgtgaga tacctgcttg ccagatgcct ggaaaccttg gctgctacaa ggatcatgga 420
aaccacctc ctctaactgg caccagtaaa acgtccaaca aactcaccat acaaacttgc 480
atcagttttt gtcggagtca gaggttcaag tttgctggga tggagtcagg ctatgcttgc 540
ttctgtggaa acaatcctga ttactggaag tacggggagg cagccagtac cgaatgcaac 600
agcgtctgct tcggggatca caccacaacc tgtggtggcg atggcaggat catcctcttt 660
gatactctcg tgggcgcctg cgttggaac tactcagoca tgtcttctgt ggtctattcc 720
cctgacttcc ccgacaccta tgcacgggg aggtctgct actggaccat ccgggttccg 780
ggggcctccc acatccactt cagcttcccc ctatttgaca tcagggactc ggcgacatg 840
gtggagcttc tggatggcta caccacccgt gtccatagccc gcttcacagg gaggagccgc 900
ccacctctgt ccttcaacgt ctctctggac ttogtoatct tgtatttctt ctctgatcgc 960
atcaatcagg cccagggatt tgctgtttta taccaagccg tcaaggaaga actgccacag 1020
gagaggcccg ctgtcaacca gacggtggcc gaggtgatca cggagcaggc caacctcagt 1080
gtcagcgctg cccggtcctc eaaagtcctc tatgtcatca ccaccagccc cagccacca 1140
cctcagactg tcccaggtag caattcctgg gcgccacca tgggggctgg aagccacaga 1200
gttgaaggat ggacagtcta tgggtctggca actctctca tctcacagt cacagccatt 1260
gtagcaaaga tacttctgca cgtcacattc aaatcccatc gtgttctgc ttcaggggac 1320
cttagggatt gtcatcaacc agggacttgc ggggaaatct ggagcatttt ttacaagcct 1380
```

tccacttcaa tttccatctt taagaagaaa ctcaagggtc agagtcaaca agatgaccgc 1440  
aatccccttg tgagtgacta aaaacccccc tgtgcctagg acttgaggtc cctctttgag 1500  
ctcaaggctg ccgtggtcaa cctctcctgt ggttcttctc tgacagactc ttccctcctc 1560  
tcctcttgcc tcggcctctt cggggaaacc ctccctcctac agactaggaa gaggcacctg 1620  
ctgccagggc aggagagcc tggattcctc ctgctt 1656

<210> 2

<211> 1425

<212> DNA

<213> Homo sapiens

<400> 2

atggcgccgc ccgcccgcgc cctcgccctg ctctccgcgc cggcgctcac gctggcggcc 60  
cgccccgcgc ctagccccgc cctcgccccc ggaccgcagt gtttcacagc caatggtgcg 120  
gattataggg gaacacagaa ctggacagca ctacaaggcg ggaagccatg tctgttttg 180  
aacgagactt tccagcatcc atacaacact ctgaaatacc ccaacgggga ggggggcctg 240  
ggtgagcaca actattgcag aaatccagat ggagacgtga gcccctggtg ctatgtggca 300  
gagcacgagg atggtgtcta ctggaagtac tgtgagatac ctgcttgcca gatgcctgga 360  
aaccttggct gctacaagga tcatggaaac ccacctcctc taactggcac cagtaaaacg 420  
tccaacaaac tcaccataca aacttgcata agtttttgc ggagtcagag gttcaagttt 480  
gctgggatgg agtcaggcta tgcttgcttc tgtggaaaca atcctgatta ctggaagtac 540  
ggggaggcag ccagtaccga atgcaacagc gtctgcttcg gggatcacac ccaaccctgt 600  
ggtggcgatg gcaggatcat cctctttgat actctcgctg gcgcctgcgc tgggaactac 660  
tcagccatgt cttctgtggt ctattcccct gacttccccg acacctatgc cacggggagg 720  
gtctgtact ggaccatccg ggttccgggg gcctcccaca tccacttcag cttcccccta 780  
tttgacatca gggactcggc ggacatggtg gagcttcttg atggtacac ccaccgtgtc 840  
ctagcccgc tccacgggag gagccgcca cctctgtcct tcaacgtctc tctggacttc 900  
gtcatcttgt atttcttctc tgatgcatac aatcaggccc agggatttgc tgttttatac 960  
caagccgtca aggaagaact gccacaggag aggccgcgtg tcaaccagac ggtggccgag 1020  
gtgatcacgg agcaggccaa cctcagtgtc agcgcgtgcc ggtcctccaa agtcctctat 1080  
gtcatcacca ccagccccag ccaccacact cagactgtcc caggtagcaa ttcctgggcg 1140  
ccaccatgg gggctggaag ccacagagtt gaaggatgga cagtctatgg tctggcaact 1200  
ctcctcatcc tcacagtcac agccattgta gcaaagatac ttctgcacgt cacattcaaa 1260  
tcccatcgtg ttctgcttc aggggacctt agggattgtc atcaaccagg gacttcgggg 1320  
gaaatctgga gcatttttta caagccttcc acttcaattt ccatctttaa gaagaaactc 1380  
aagggtcaga gtcaacaaga tgaccgcaat ccccttgtga gtgac 1425

<210> 3

<211> 475

<212> PRT

<213> Homo sapiens

<400> 3

Met Ala Pro Pro Ala Ala Arg Leu Ala Leu Leu Ser Ala Ala Ala Leu  
1 5 10 15

Thr Leu Ala Ala Arg Pro Ala Pro Ser Pro Gly Leu Gly Pro Gly Pro  
20 25 30

Glu Cys Phe Thr Ala Asn Gly Ala Asp Tyr Arg Gly Thr Gln Asn Trp  
 35 40 45  
 Thr Ala Leu Gln Gly Gly Lys Pro Cys Leu Phe Trp Asn Glu Thr Phe  
 50 55 60  
 Gln His Pro Tyr Asn Thr Leu Lys Tyr Pro Asn Gly Glu Gly Gly Leu  
 65 70 75 80  
 Gly Glu His Asn Tyr Cys Arg Asn Pro Asp Gly Asp Val Ser Pro Trp  
 85 90 95  
 Cys Tyr Val Ala Glu His Glu Asp Gly Val Tyr Trp Lys Tyr Cys Glu  
 100 105 110  
 Ile Pro Ala Cys Gln Met Pro Gly Asn Leu Gly Cys Tyr Lys Asp His  
 115 120 125  
 Gly Asn Pro Pro Pro Leu Thr Gly Thr Ser Lys Thr Ser Asn Lys Leu  
 130 135 140  
 Thr Ile Gln Thr Cys Ile Ser Phe Cys Arg Ser Gln Arg Phe Lys Phe  
 145 150 155 160  
 Ala Gly Met Glu Ser Gly Tyr Ala Cys Phe Cys Gly Asn Asn Pro Asp  
 165 170 175  
 Tyr Trp Lys Tyr Gly Glu Ala Ala Ser Thr Glu Cys Asn Ser Val Cys  
 180 185 190  
 Phe Gly Asp His Thr Gln Pro Cys Gly Gly Asp Gly Arg Ile Ile Leu  
 195 200 205  
 Phe Asp Thr Leu Val Gly Ala Cys Gly Gly Asn Tyr Ser Ala Met Ser  
 210 215 220  
 Ser Val Val Tyr Ser Pro Asp Phe Pro Asp Thr Tyr Ala Thr Gly Arg  
 225 230 235 240  
 Val Cys Tyr Trp Thr Ile Arg Val Pro Gly Ala Ser His Ile His Phe  
 245 250 255  
 Ser Phe Pro Leu Phe Asp Ile Arg Asp Ser Ala Asp Met Val Glu Leu  
 260 265 270  
 Leu Asp Gly Tyr Thr His Arg Val Leu Ala Arg Phe His Gly Arg Ser  
 275 280 285

Arg Pro Pro Leu Ser Phe Asn Val Ser Leu Asp Phe Val Ile Leu Tyr  
290 295 300

Phe Phe Ser Asp Arg Ile Asn Gln Ala Gln Gly Phe Ala Val Leu Tyr  
305 310 315 320

Gln Ala Val Lys Glu Glu Leu Pro Gln Glu Arg Pro Ala Val Asn Gln  
325 330 335

Thr Val Ala Glu Val Ile Thr Glu Gln Ala Asn Leu Ser Val Ser Ala  
340 345 350

Ala Arg Ser Ser Lys Val Leu Tyr Val Ile Thr Thr Ser Pro Ser His  
355 360 365

Pro Pro Gln Thr Val Pro Gly Ser Asn Ser Trp Ala Pro Pro Met Gly  
370 375 380

Ala Gly Ser His Arg Val Glu Gly Trp Thr Val Tyr Gly Leu Ala Thr  
385 390 395 400

Leu Leu Ile Leu Thr Val Thr Ala Ile Val Ala Lys Ile Leu Leu His  
405 410 415

Val Thr Phe Lys Ser His Arg Val Pro Ala Ser Gly Asp Leu Arg Asp  
420 425 430

Cys His Gln Pro Gly Thr Ser Gly Glu Ile Trp Ser Ile Phe Tyr Lys  
435 440 445

Pro Ser Thr Ser Ile Ser Ile Phe Lys Lys Lys Leu Lys Gly Gln Ser  
450 455 460

Gln Gln Asp Asp Arg Asn Pro Leu Val Ser Asp  
465 470 475

<210> 4

<211> 19

<212> PRT

<213> Homo sapiens

<400> 4

Met Ala Pro Pro Ala Ala Arg Leu Ala Leu Leu Ser Ala Ala Ala Leu  
1 5 10 15

Thr Leu Ala

<210> 5  
 <211> 456  
 <212> PRT  
 <213> Homo sapiens

<400> 5

Ala	Arg	Pro	Ala	Pro	Ser	Pro	Gly	Leu	Gly	Pro	Gly	Pro	Glu	Cys	Phe
1				5					10					15	
Thr	Ala	Asn	Gly	Ala	Asp	Tyr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Ala	Leu
		20						25					30		
Gln	Gly	Gly	Lys	Pro	Cys	Leu	Phe	Trp	Asn	Glu	Thr	Phe	Gln	His	Pro
	35						40					45			
Tyr	Asn	Thr	Leu	Lys	Tyr	Pro	Asn	Gly	Glu	Gly	Gly	Leu	Gly	Glu	His
	50					55						60			
Asn	Tyr	Cys	Arg	Asn	Pro	Asp	Gly	Asp	Val	Ser	Pro	Trp	Cys	Tyr	Val
65				70						75					80
Ala	Glu	His	Glu	Asp	Gly	Val	Tyr	Trp	Lys	Tyr	Cys	Glu	Ile	Pro	Ala
			85						90					95	
Cys	Gln	Met	Pro	Gly	Asn	Leu	Gly	Cys	Tyr	Lys	Asp	His	Gly	Asn	Pro
		100						105						110	
Pro	Pro	Leu	Thr	Gly	Thr	Ser	Lys	Thr	Ser	Asn	Lys	Leu	Thr	Ile	Gln
		115					120						125		
Thr	Cys	Ile	Ser	Phe	Cys	Arg	Ser	Gln	Arg	Phe	Lys	Phe	Ala	Gly	Met
	130					135					140				
Glu	Ser	Gly	Tyr	Ala	Cys	Phe	Cys	Gly	Asn	Asn	Pro	Asp	Tyr	Trp	Lys
145					150					155					160
Tyr	Gly	Glu	Ala	Ala	Ser	Thr	Glu	Cys	Asn	Ser	Val	Cys	Phe	Gly	Asp
			165						170					175	
His	Thr	Gln	Pro	Cys	Gly	Gly	Asp	Gly	Arg	Ile	Ile	Leu	Phe	Asp	Thr
		180						185						190	
Leu	Val	Gly	Ala	Cys	Gly	Gly	Asn	Tyr	Ser	Ala	Met	Ser	Ser	Val	Val
	195						200					205			

Tyr Ser Pro Asp Phe Pro Asp Thr Tyr Ala Thr Gly Arg Val Cys Tyr  
 210 215 220

Trp Thr Ile Arg Val Pro Gly Ala Ser His Ile His Phe Ser Phe Pro  
 225 230 235 240

Leu Phe Asp Ile Arg Asp Ser Ala Asp Met Val Glu Leu Leu Asp Gly  
 245 250 255

Tyr Thr His Arg Val Leu Ala Arg Phe His Gly Arg Ser Arg Pro Pro  
 260 265 270

Leu Ser Phe Asn Val Ser Leu Asp Phe Val Ile Leu Tyr Phe Phe Ser  
 275 280 285

Asp Arg Ile Asn Gln Ala Gln Gly Phe Ala Val Leu Tyr Gln Ala Val  
 290 295 300

Lys Glu Glu Leu Pro Gln Glu Arg Pro Ala Val Asn Gln Thr Val Ala  
 305 310 315 320

Glu Val Ile Thr Glu Gln Ala Asn Leu Ser Val Ser Ala Ala Arg Ser  
 325 330 335

Ser Lys Val Leu Tyr Val Ile Thr Thr Ser Pro Ser His Pro Pro Gln  
 340 345 350

Thr Val Pro Gly Ser Asn Ser Trp Ala Pro Pro Met Gly Ala Gly Ser  
 355 360 365

His Arg Val Glu Gly Trp Thr Val Tyr Gly Leu Ala Thr Leu Leu Ile  
 370 375 380

Leu Thr Val Thr Ala Ile Val Ala Lys Ile Leu Leu His Val Thr Phe  
 385 390 395 400

Lys Ser His Arg Val Pro Ala Ser Gly Asp Leu Arg Asp Cys His Gln  
 405 410 415

Pro Gly Thr Ser Gly Glu Ile Trp Ser Ile Phe Tyr Lys Pro Ser Thr  
 420 425 430

Ser Ile Ser Ile Phe Lys Lys Lys Leu Lys Gly Gln Ser Gln Gln Asp  
 435 440 445

Asp Arg Asn Pro Leu Val Ser Asp  
 450 455

<210> 6  
 <211> 373  
 <212> PRT  
 <213> Homo sapiens

<400> 6

Ala Arg Pro Ala Pro Ser Pro Gly Leu Gly Pro Gly Pro Glu Cys Phe  
 1 5 10 15

Thr Ala Asn Gly Ala Asp Tyr Arg Gly Thr Gln Asn Trp Thr Ala Leu  
 20 25 30

Gln Gly Gly Lys Pro Cys Leu Phe Trp Asn Glu Thr Phe Gln His Pro  
 35 40 45

Tyr Asn Thr Leu Lys Tyr Pro Asn Gly Glu Gly Gly Leu Gly Glu His  
 50 55 60

Asn Tyr Cys Arg Asn Pro Asp Gly Asp Val Ser Pro Trp Cys Tyr Val  
 65 70 75 80

Ala Glu His Glu Asp Gly Val Tyr Trp Lys Tyr Cys Glu Ile Pro Ala  
 85 90 95

Cys Gln Met Pro Gly Asn Leu Gly Cys Tyr Lys Asp His Gly Asn Pro  
 100 105 110

Pro Pro Leu Thr Gly Thr Ser Lys Thr Ser Asn Lys Leu Thr Ile Gln  
 115 120 125

Thr Cys Ile Ser Phe Cys Arg Ser Gln Arg Phe Lys Phe Ala Gly Met  
 130 135 140

Glu Ser Gly Tyr Ala Cys Phe Cys Gly Asn Asn Pro Asp Tyr Trp Lys  
 145 150 155 160

Tyr Gly Glu Ala Ala Ser Thr Glu Cys Asn Ser Val Cys Phe Gly Asp  
 165 170 175

His Thr Gln Pro Cys Gly Gly Asp Gly Arg Ile Ile Leu Phe Asp Thr  
 180 185 190

Leu Val Gly Ala Cys Gly Gly Asn Tyr Ser Ala Met Ser Ser Val Val  
 195 200 205

Tyr Ser Pro Asp Phe Pro Asp Thr Tyr Ala Thr Gly Arg Val Cys Tyr  
 210 215 220

Trp Thr Ile Arg Val Pro Gly Ala Ser His Ile His Phe Ser Phe Pro  
225 230 235 240

Leu Phe Asp Ile Arg Asp Ser Ala Asp Met Val Glu Leu Leu Asp Gly  
245 250 255

Tyr Thr His Arg Val Leu Ala Arg Phe His Gly Arg Ser Arg Pro Pro  
260 265 270

Leu Ser Phe Asn Val Ser Leu Asp Phe Val Ile Leu Tyr Phe Phe Ser  
275 280 285

Asp Arg Ile Asn Gln Ala Gln Gly Phe Ala Val Leu Tyr Gln Ala Val  
290 295 300

Lys Glu Glu Leu Pro Gln Glu Arg Pro Ala Val Asn Gln Thr Val Ala  
305 310 315 320

Glu Val Ile Thr Glu Gln Ala Asn Leu Ser Val Ser Ala Ala Arg Ser  
325 330 335

Ser Lys Val Leu Tyr Val Ile Thr Thr Ser Pro Ser His Pro Pro Gln  
340 345 350

Thr Val Pro Gly Ser Asn Ser Trp Ala Pro Pro Met Gly Ala Gly Ser  
355 360 365

His Arg Val Glu Gly  
370

<210> 7

<211> 23

<212> PRT

<213> Homo sapiens

<400> 7

Trp Thr Val Tyr Gly Leu Ala Thr Leu Leu Ile Leu Thr Val Thr Ala  
1 5 10 15

Ile Val Ala Lys Ile Leu Leu  
20

<210> 8

<211> 60

<212> PRT

<213> Homo sapiens

<400> 8

His Val Thr Phe Lys Ser His Arg Val Pro Ala Ser Gly Asp Leu Arg  
1 5 10 15

Asp Cys His Gln Pro Gly Thr Ser Gly Glu Ile Trp Ser Ile Phe Tyr  
20 25 30

Lys Pro Ser Thr Ser Ile Ser Ile Phe Lys Lys Lys Leu Lys Gly Gln  
35 40 45

Ser Gln Gln Asp Asp Arg Asn Pro Leu Val Ser Asp  
50 55 60

<210> 9

<211> 4628

<212> DNA

<213> Homo sapiens

<400> 9

gcggccgctc gcgatctaga actagtaatg atgctgcctc aaaactcgtg gcatattgat 60  
tttggaagat gctgctgtca tcagaacctt ttctctgctg tggtaacttg catcctgctc 120  
ctgaattcct gctttctcat cagcagtttt aatggaacag atttggagtt gaggctgggc 180  
aatggagacg gtccctgctc tgggacagtg gaggtgaaat tccagggaca gtgggggact 240  
gtgtgtgatg atgggtggaa cactactgcc tcaactgtcg tgtgcaaaca gcttggatgt 300  
ccattttctt tcgccatggt tcgttttgga caagccgtga ctagacatgg aaaaatttgg 360  
cttgatgatg tttcctgtta tggaaatgag tcagctctct gggaatgtca acaccgggaa 420  
tggggaagcc ataactgtta tcatggagaa gatgttggtg tgaactgtta tgggtgaagcc 480  
aatctggggt tggagctagt ggatggaaac aactcctgtt caggagagat ggaggtgaaa 540  
ttccaagaaa ggtgggggac tataatgtgat gatgggtgga acttgaatac tgctgccgtg 600  
gtgtgcaggc aactagatg tccatcttct tttatttctt ctggagttgt taatagccct 660  
gctgtattgc gccccatttg gctggatgac attttatgcc aggggaatga gttggcactc 720  
tggaattgca gacatcgtgg atggggaaat catgactgca gtcacaatga ggatgtcaca 780  
ttaacttggt atgatagtag tgatcttgaa ctaaggcttg taggtggaac taaccgctgt 840  
atggggagag tagagctgaa aatccaagga aggtggggga ccgatgccca ccataagtgg 900  
aacaatgctg cagctgatgt cgtatgcaag cagttgggat gtggaaccgc acttcacttc 960  
gctggcttgc ctcatcttgc gtcagggtct gatgttgat ggcttgatgg tgtctcctgc 1020  
tccggtaatg aatcttttct ttgggactgc agacattccg gaaccgtcaa ttttgactgt 1080  
cttcatcaaa acgatgtgtc tgtgatctgc tcagatggag cagatttga actgcgacta 1140  
gcagatggaa gtaacaattg ttcagggaga gtagaggtga gaattcatga acagtgggtg 1200  
acaatatgtg accagaactg gaagaatgaa caagcccttg tggtttgtaa gcagctagga 1260  
tgtccgttca gcgtcttttg cagtcgtcgt gctaaacctg gtaatgaagc tagagacatt 1320  
tggtataaca gcatatcttg cactgggaat gagtcagctc tctgggactg cacatatgat 1380  
ggaaaagcaa agcgaacatg cttccgaaga tcagatgctg gagtaatttg ttctgataag 1440  
gcagatctgg acctaaggct tgtcggggct catagcccct gttatgggag attggaggtg 1500  
aaataccaag gagagtgggg gactgtgtgt catgacagat ggagcacaag gaatgcagct 1560

gttgtgtgta	aacaattggg	atgtggaaag	cctatgcatg	tgttttggtat	gacctatttt	1620
aaagaagcat	caggacctat	ttggctggat	gacgtttctt	gcattggaaa	tgagtcaa	1680
atctgggact	gtgaacacag	tggatgggga	aagcataatt	gtgtacacag	agaggatgtg	1740
attgtaacct	gtcaggtga	tgcaacatgg	ggcctgaggc	tggtagggcg	cagcaaccgc	1800
tgctcgggaa	gactggaggt	gtactttcaa	ggacgggtgg	gcacagtgtg	tgatgacggc	1860
tggaaacagta	aagctgcagc	tgtggtgtgt	agccagctgg	actgcccatc	ttctatcatt	1920
ggcatgggtc	tgggaaacgc	ttctacagga	tatggaaaaa	tttggctcga	tgatgtttcc	1980
tgtgatggag	atgagtcaga	tctctgggtca	tgcaggaaaca	gtgggtgggg	aaataatgac	2040
tgcagtcaca	gtgaagatgt	tggagtgatc	tgttctgatg	catcgatat	ggagctgagg	2100
cttgtgggtg	gaagcagcag	gtgtgctgga	aaagttgagg	tgaatgtcca	gggtgccgtg	2160
ggaattctgt	gtgctaattg	ctggggaatg	aacattgctg	aagttgtttg	caggcaactt	2220
gaatgtgggt	ctgcaatcag	ggtctccaga	gagcctcatt	tcacagaaag	aacattacac	2280
atcttaattgt	cgaattctgg	ctgcaactgga	ggggaagcct	ctctctggga	ttgtatacga	2340
tgggagtggga	aacagactgc	gtgtcattta	aatatggaag	caagtttgat	ctgctcagcc	2400
cacaggcagc	ccaggctggt	tggagctgat	atgccctgct	ctggacgtgt	tgaagtga	2460
catgcagaca	catggcgctc	tgtctgtgat	tctgatttct	ctcttcacgc	tgccaatgtg	2520
ctgtgcagag	aattaaattg	tggagatgcc	atatctcttt	ctgtgggaga	tcacttttga	2580
aaagggaatg	gtctaacttg	ggccgaaaag	ttccagtgtg	aagggagtga	aactcacctt	2640
gcattatgcc	ccattgttca	acatccggaa	gacacttgta	tcacagcag	agaagttgga	2700
gttgtctgtt	cccgatatac	agatgtccga	cttgtgaatg	gcaaatacca	gtgtgacggg	2760
caagtggaga	tcaacgtgct	tggacactgg	ggctcactgt	gtgacacca	ctgggacca	2820
gaagatgccc	gtgttctatg	cagacagctc	agctgtggga	ctgctctctc	aaccacagga	2880
ggaaaatata	ttggagaaag	aagtgttcgt	gtgtggggac	acaggtttca	ttgcttaggg	2940
aatgagtcac	ttctggataa	ctgtcaaattg	acagttcttg	gagcacctcc	ctgtatccat	3000
ggaaatactg	tctctgtgat	ctgcacagga	agcctgaccc	agccactgtt	tccatgcctc	3060
gcaaattgtat	ctgacccata	tttgtctgca	gttccagagg	gcagtgtctt	gatctgtctt	3120
gaggacaaaac	ggctccgcct	agtggatggg	gacagccgct	gtgccgggag	agtagagatc	3180
tatcacgacg	gcttctgggg	caccatctgt	gatgacggct	gggacctgag	cgatgccac	3240
gtggtgtgtc	aaaagctggg	ctgtggagtg	gccttcaatg	ccacggtctc	tgctcacttt	3300
ggggaggggt	cagggcccat	ctggctggat	gacctgaact	gcacaggaac	ggagtccac	3360
ttgtggcagt	gcccttcccg	cggctggggg	cagcacgact	gcaggcaca	ggaggacgca	3420
ggggtcatct	gctcagaatt	cacagccttg	aggctctaca	gtgaaactga	aacagagagc	3480
tgtgctggga	gattggaagt	cttctataac	gggacctggg	gcagcgtcgg	caggaggaac	3540
atcaccacag	ccatagcagg	cattgtgtgc	aggcagctgg	gctgtgggga	gaatggagtt	3600
gtcagcctcg	cccctttatc	taagacaggc	tctggtttca	tgtgggtgga	tgacattcag	3660
tgtcctaaaa	cgcatatctc	catatggcag	tgctgtctg	ccccatggga	gcgaagaatc	3720
tcagcccag	cagaagagac	ctggatcaca	tgtgaagata	gaataagagt	gcgtggagga	3780
gacaccgagt	gctctgggag	agtggagatc	tggcacgcag	gctcctgggg	cacagtgtgt	3840
gatgactcct	gggacctggc	cgaggcgga	gtggtgtgtc	agcagctggg	ctgtggctct	3900
gctctggctg	ccctgaggga	cgcttcgttt	ggccaggga	ctggaaccat	ctggttggat	3960
gacatgcggt	gcaaaggaaa	tgagtcat	ctatgggaact	gtcacgcaa	accctgggga	4020
cagagtgact	gtggacacaa	ggaagatgct	ggcgtgaggt	gctctggaca	gtcgctgaaa	4080
tcactgaatg	cctcctcagg	tcatttagca	cttattttat	ccagtatctt	tgggctcctt	4140
ctcctgggtc	tgttttattct	atttctcacg	tgggtccgag	ttcagaaaca	aaaacatctg	4200
cccctcagag	tttcaaccag	aaggaggggt	tctctcgagg	agaatttatt	ccatgagatg	4260
gagacctgcc	tcaagagaga	ggacccacat	gggacaagaa	cctcagatga	cacccccaac	4320
catggttgtg	aagatgctag	cgacacatcg	ctgttgggag	ttcttctctg	ctctgaagcc	4380
acaaaatgac	tttagacttc	cagggtcac	cagatcaacc	tctaaatata	tttgaaggag	4440

acaacaactt	ttaaataaat	aaagaggaag	tcaagttgcc	ctatggaaaa	cttgtccaaa	4500
taacatttct	tgaacaatag	gagaacagct	aaattgataa	agactgggtga	taataaaaaat	4560
tgaattatgt	atatcactgt	taaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	acggacgcgt	4620
gggtcgac						4628

<210> 10  
 <211> 4359  
 <212> DNA  
 <213> Homo sapiens

<400> 10						
atgatgctgc	ctcaaaactc	gtggcatatt	gatttttgaa	gatgctgctg	tcatcagaac	60
cttttctctg	ctgtggtaac	ttgcatcctg	ctcctgaatt	cctgctttct	catcagcagt	120
tttaaatgga	cagatttgga	gttgaggctg	gtcaatggag	acggtccttg	ctctgggaca	180
gtggaggtga	aattccaggg	acagtggggg	actgtgtgtg	atgatgggtg	gaacactact	240
gcctcaactg	tcgtgtgcaa	acagcttgga	tgtccatttt	ctttcgccat	gtttcgtttt	300
ggacaagccg	tgactagaca	tggaaaaatt	tggcttgatg	atgtttcctg	ttatggaaat	360
gagtcagctc	tctgggaatg	tcaacaccgg	gaatggggaa	gccataactg	ttatcatgga	420
gaagatgttg	gtgtgaactg	ttatggtgaa	gccaatctgg	gtttgaggct	agtggatgga	480
aacaactcct	gttcaggagg	agtggagggt	aaattccaag	aaagggtggg	gactatatgt	540
gatgatgggt	ggaacttgaa	tactgctgcc	gtggtgtgca	ggcaactagg	atgtccatct	600
tcttttattt	cttctggagt	tgtaataagc	cctgctgtat	tgcgccccat	ttggctggat	660
gacattttat	gccaggggaa	tgagttagga	ctctggaatt	gcagacatcg	tggatgggga	720
aatcatgact	gcagtcacaa	tgaggatgtc	acattaactt	gttatgatag	tagtgatctt	780
gaactaaggc	ttgtagggtg	aactaaccgc	tgtatgggga	gagtagagct	gaaaatccaa	840
ggaagggtgg	ggaccgtatg	ccaccataag	tggaaacaatg	ctgcagctga	tgtcgtatgc	900
aagcagttgg	gatgtggaac	cgcacttcac	ttcgctggct	tgcctcattt	gcagtcaggg	960
tctgatgttg	tatggcttga	tggtgtctcc	tgtccggta	atgaatcttt	tctttgggac	1020
tgcagacatt	ccggaaccgt	caattttgac	tgtcttcac	aaaacgatgt	gtctgtgatc	1080
tgctcagatg	gagcagattt	ggaactgcga	ctagcagatg	gaagtaacaa	ttgttcaggg	1140
agagtagagg	tgagaattca	tgaacagtgg	tggacaatat	gtgaccagaa	ctggaagaat	1200
gaacaagccc	ttgtggtttg	taagcagcta	ggatgtccgt	tcagcgtctt	tggcagtcgt	1260
cgtgctaaac	ctagtaatga	agctagagac	atttggataa	acagcatatc	ttgcactggg	1320
aatgagtcag	ctctctggga	ctgcacatat	gatggaaaag	caaagcgaac	atgcttccga	1380
agatcagatg	ctggagtaat	ttgttctgat	aaggcagatc	tggacctaa	gcttgtcggg	1440
gctcatagcc	cctgttatgg	gagattggag	gtgaaatacc	aaggagagt	ggggactgtg	1500
tgtcatgaca	gatggagcac	aaggaatgca	gctgttgtgt	gtaaacaatt	gggatgtgga	1560
aagcctatgc	atgtgttttg	tatgacctat	tttaaagaag	catcaggacc	tatttggctg	1620
gatgacgttt	cttgcatggg	aaatgagtca	aatatctggg	actgtgaaca	cagtggatgg	1680
ggaaagcata	attgtgtaca	cagagaggat	gtgattgtaa	cctgctcagg	tgatgcaaca	1740
tggggcctga	ggctgggtgg	cggcagcaac	cgctgctcgg	gaagactgga	ggtgtacttt	1800
caaggacggg	ggggcacagt	gtgtgatgac	ggctggaaca	gtaaagctgc	agctgtgggtg	1860
tgtagccagc	tggactgccc	atcttctatc	attggcatgg	gtctgggaaa	cgcttctaca	1920
ggatatggaa	aaatttggct	cgatgatgtt	tcctgtgatg	gagatgagtc	agatctcttg	1980
tcatgcagga	acagtgggtg	gggaaataat	gactgcagtc	acagtgaaga	tgttggagt	2040
atctgttctg	atgcatcgga	tatggagctg	aggcttgtgg	gtggaagcag	caggtgtgct	2100
ggaaaagtgg	aggtgaatgt	ccagggtgcc	gtgggaattc	tgtgtgctaa	tggctgggga	2160
atgaacattg	ctgaagttgt	ttgcaggcaa	cttgaatgtg	ggtctgcaat	caggggtctcc	2220

agagagcctc atttcacaga aagaacatta cacatcttaa tgtcgaattc tggctgcact 2280  
ggaggggaag cctctctctg ggattgtata cgatgggagt ggaaacagac tgcgtgtcat 2340  
ttaaataatgg aagcaagttt gatctgctca gccacagggc agcccaggct ggttggagct 2400  
gatatgcctt gctctggacg tgttgaagtg aaacatgcag acacatggcg ctctgtctgt 2460  
gattctgatt tctctcttca tgctgccaat gtgctgtgca gagaattaaa ttgtggagat 2520  
gccatatctc tttctgtggg agatcacttt ggaaaaggga atggtctaac ttgggccgaa 2580  
aagttccagt gtgaaggag tgaaactcac cttgcattat gcccattgt tcaacatccg 2640  
gaagacactt gtatccacag cagagaagtt ggagttgtct gttcccgata tacagatgtc 2700  
cgacttgtga atggcaaata ccagtgtgac gggcaagtgg agatcaacgt gcttggacac 2760  
tggggctcac tgtgtgacac ccactgggac ccagaagatg cccgtgttct atgcagacag 2820  
ctcagctgtg ggactgctct ctcaaccaca ggaggaaaat atattggaga aagaagtgtt 2880  
cgtgtgtggg gacacagggt tcattgctta gggaatgagt cacttctgga taactgtcaa 2940  
atgacagttc ttggagcacc tccctgtatc catggaaata ctgtctctgt gatctgcaca 3000  
ggaagcctga cccagccact gtttccatgc ctgcgaaatg tatctgacct atatttgtct 3060  
gcagttccag agggcagtg tttgatctgc ttagaggaca aacggctccg cctagtggat 3120  
ggggacagcc gctgtgccg gagagtagag atctatcacg acggcttctg gggcaccatc 3180  
tgtgatgacg gctgggacct gagcgtgcc cacgtggtgt gtcaaaagct gggctgtgga 3240  
gtggccttca atgccacggt ctctgctcac tttggggagg ggtcagggcc catctggctg 3300  
gatgacctga actgcacagg aacggagtcc cacttgtggc agtgcccttc ccgcggtgg 3360  
gggcagcacg actgcaggca caaggaggac gcaggggtca tctgctcaga attcacagcc 3420  
ttgaggctct acagtgaac tgaaacagag agctgtgctg ggagattgga agtcttctat 3480  
aacgggacct ggggcagcgt cggcaggagg aacatcacca cagccatagc aggcattgtg 3540  
tgcaggcagc tgggctgtgg ggagaatgga gttgtcagcc tcgccccctt atctaagaca 3600  
ggctctggtt tcatgtgggt ggatgacatt cagtgtccta aaacgcatat ctccatattg 3660  
cagtgcctgt ctgccccatg ggagcgaaga atctccagcc cagcagaaga gacctggatc 3720  
acatgtgaag atagaataag agtgctgga ggagacaccg agtgctctgg gagagtggag 3780  
atctggcacg caggctcctg gggcacagtg tgtgatgact cctgggacct ggccgaggcg 3840  
gaagtgggtg gtcagcagct gggctgtggc tctgctctgg ctgccctgag ggacgcttcg 3900  
tttgccagc gaactggaac catctgggtg gatgacatgc ggtgcaaagg aaatgagtca 3960  
tttctatggg actgtcacgc caaacctgg ggacagagt actgtggaca caaggaagat 4020  
gctggcgtga ggtgctctgg acagtgcgtg aaatcactga atgcctoctc aggtcattta 4080  
gcacttattt tatccagtat ctttgggctc cttctcctgg ttctgtttat tctatttctc 4140  
acgtggtgcc gagttcagaa acaaaaacat ctgccctca gagtttcaac cagaaggagg 4200  
ggttctctcg aggagaattt attccatgag atggagacct gcctcaagag agaggacca 4260  
catgggacaa gaacctcaga tgacaccccc aacctggtt gtgaagatgc tagcgacaca 4320  
tcgctgttgg gagttcttcc tgcctctgaa gccacaaaa 4359

<210> 11

<211> 1453

<212> PRT

<213> Homo sapiens

<400> 11

Met Met Leu Pro Gln Asn Ser Trp His Ile Asp Phe Gly Arg Cys Cys  
1 5 10 15

Cys His Gln Asn Leu Phe Ser Ala Val Val Thr Cys Ile Leu Leu Leu  
20 25 30



His Lys Trp Asn Asn Ala Ala Ala Asp Val Val Cys Lys Gln Leu Gly  
 290 295 300

Cys Gly Thr Ala Leu His Phe Ala Gly Leu Pro His Leu Gln Ser Gly  
 305 310 315 320

Ser Asp Val Val Trp Leu Asp Gly Val Ser Cys Ser Gly Asn Glu Ser  
 325 330 335

Phe Leu Trp Asp Cys Arg His Ser Gly Thr Val Asn Phe Asp Cys Leu  
 340 345 350

His Gln Asn Asp Val Ser Val Ile Cys Ser Asp Gly Ala Asp Leu Glu  
 355 360 365

Leu Arg Leu Ala Asp Gly Ser Asn Asn Cys Ser Gly Arg Val Glu Val  
 370 375 380

Arg Ile His Glu Gln Trp Trp Thr Ile Cys Asp Gln Asn Trp Lys Asn  
 385 390 395 400

Glu Gln Ala Leu Val Val Cys Lys Gln Leu Gly Cys Pro Phe Ser Val  
 405 410 415

Phe Gly Ser Arg Arg Ala Lys Pro Ser Asn Glu Ala Arg Asp Ile Trp  
 420 425 430

Ile Asn Ser Ile Ser Cys Thr Gly Asn Glu Ser Ala Leu Trp Asp Cys  
 435 440 445

Thr Tyr Asp Gly Lys Ala Lys Arg Thr Cys Phe Arg Arg Ser Asp Ala  
 450 455 460

Gly Val Ile Cys Ser Asp Lys Ala Asp Leu Asp Leu Arg Leu Val Gly  
 465 470 475 480

Ala His Ser Pro Cys Tyr Gly Arg Leu Glu Val Lys Tyr Gln Gly Glu  
 485 490 495

Trp Gly Thr Val Cys His Asp Arg Trp Ser Thr Arg Asn Ala Ala Val  
 500 505 510

Val Cys Lys Gln Leu Gly Cys Gly Lys Pro Met His Val Phe Gly Met  
 515 520 525

Thr Tyr Phe Lys Glu Ala Ser Gly Pro Ile Trp Leu Asp Asp Val Ser  
 530 535 540

Cys Ile Gly Asn Glu Ser Asn Ile Trp Asp Cys Glu His Ser Gly Trp  
545 550 555 560

Gly Lys His Asn Cys Val His Arg Glu Asp Val Ile Val Thr Cys Ser  
565 570 575

Gly Asp Ala Thr Trp Gly Leu Arg Leu Val Gly Gly Ser Asn Arg Cys  
580 585 590

Ser Gly Arg Leu Glu Val Tyr Phe Gln Gly Arg Trp Gly Thr Val Cys  
595 600 605

Asp Asp Gly Trp Asn Ser Lys Ala Ala Ala Val Val Cys Ser Gln Leu  
610 615 620

Asp Cys Pro Ser Ser Ile Ile Gly Met Gly Leu Gly Asn Ala Ser Thr  
625 630 635 640

Gly Tyr Gly Lys Ile Trp Leu Asp Asp Val Ser Cys Asp Gly Asp Glu  
645 650 655

Ser Asp Leu Trp Ser Cys Arg Asn Ser Gly Trp Gly Asn Asn Asp Cys  
660 665 670

Ser His Ser Glu Asp Val Gly Val Ile Cys Ser Asp Ala Ser Asp Met  
675 680 685

Glu Leu Arg Leu Val Gly Gly Ser Ser Arg Cys Ala Gly Lys Val Glu  
690 695 700

Val Asn Val Gln Gly Ala Val Gly Ile Leu Cys Ala Asn Gly Trp Gly  
705 710 715 720

Met Asn Ile Ala Glu Val Val Cys Arg Gln Leu Glu Cys Gly Ser Ala  
725 730 735

Ile Arg Val Ser Arg Glu Pro His Phe Thr Glu Arg Thr Leu His Ile  
740 745 750

Leu Met Ser Asn Ser Gly Cys Thr Gly Gly Glu Ala Ser Leu Trp Asp  
755 760 765

Cys Ile Arg Trp Glu Trp Lys Gln Thr Ala Cys His Leu Asn Met Glu  
770 775 780

Ala Ser Leu Ile Cys Ser Ala His Arg Gln Pro Arg Leu Val Gly Ala  
785 790 795 800

Asp Met Pro Cys Ser Gly Arg Val Glu Val Lys His Ala Asp Thr Trp  
805 810 815

Arg Ser Val Cys Asp Ser Asp Phe Ser Leu His Ala Ala Asn Val Leu  
820 825 830

Cys Arg Glu Leu Asn Cys Gly Asp Ala Ile Ser Leu Ser Val Gly Asp  
835 840 845

His Phe Gly Lys Gly Asn Gly Leu Thr Trp Ala Glu Lys Phe Gln Cys  
850 855 860

Glu Gly Ser Glu Thr His Leu Ala Leu Cys Pro Ile Val Gln His Pro  
865 870 875 880

Glu Asp Thr Cys Ile His Ser Arg Glu Val Gly Val Val Cys Ser Arg  
885 890 895

Tyr Thr Asp Val Arg Leu Val Asn Gly Lys Ser Gln Cys Asp Gly Gln  
900 905 910

Val Glu Ile Asn Val Leu Gly His Trp Gly Ser Leu Cys Asp Thr His  
915 920 925

Trp Asp Pro Glu Asp Ala Arg Val Leu Cys Arg Gln Leu Ser Cys Gly  
930 935 940

Thr Ala Leu Ser Thr Thr Gly Gly Lys Tyr Ile Gly Glu Arg Ser Val  
945 950 955 960

Arg Val Trp Gly His Arg Phe His Cys Leu Gly Asn Glu Ser Leu Leu  
965 970 975

Asp Asn Cys Gln Met Thr Val Leu Gly Ala Pro Pro Cys Ile His Gly  
980 985 990

Asn Thr Val Ser Val Ile Cys Thr Gly Ser Leu Thr Gln Pro Leu Phe  
995 1000 1005

Pro Cys Leu Ala Asn Val Ser Asp Pro Tyr Leu Ser Ala Val Pro Glu  
1010 1015 1020

Gly Ser Ala Leu Ile Cys Leu Glu Asp Lys Arg Leu Arg Leu Val Asp  
1025 1030 1035 1040

Gly Asp Ser Arg Cys Ala Gly Arg Val Glu Ile Tyr His Asp Gly Phe  
1045 1050 1055

Trp Gly Thr Ile Cys Asp Asp Gly Trp Asp Leu Ser Asp Ala His Val  
 1060 1065 1070

Val Cys Gln Lys Leu Gly Cys Gly Val Ala Phe Asn Ala Thr Val Ser  
 1075 1080 1085

Ala His Phe Gly Glu Gly Ser Gly Pro Ile Trp Leu Asp Asp Leu Asn  
 1090 1095 1100

Cys Thr Gly Thr Glu Ser His Leu Trp Gln Cys Pro Ser Arg Gly Trp  
 1105 1110 1115 1120

Gly Gln His Asp Cys Arg His Lys Glu Asp Ala Gly Val Ile Cys Ser  
 1125 1130 1135

Glu Phe Thr Ala Leu Arg Leu Tyr Ser Glu Thr Glu Thr Glu Ser Cys  
 1140 1145 1150

Ala Gly Arg Leu Glu Val Phe Tyr Asn Gly Thr Trp Gly Ser Val Gly  
 1155 1160 1165

Arg Arg Asn Ile Thr Thr Ala Ile Ala Gly Ile Val Cys Arg Gln Leu  
 1170 1175 1180

Gly Cys Gly Glu Asn Gly Val Val Ser Leu Ala Pro Leu Ser Lys Thr  
 1185 1190 1195 1200

Gly Ser Gly Phe Met Trp Val Asp Asp Ile Gln Cys Pro Lys Thr His  
 1205 1210 1215

Ile Ser Ile Trp Gln Cys Leu Ser Ala Pro Trp Glu Arg Arg Ile Ser  
 1220 1225 1230

Ser Pro Ala Glu Glu Thr Trp Ile Thr Cys Glu Asp Arg Ile Arg Val  
 1235 1240 1245

Arg Gly Gly Asp Thr Glu Cys Ser Gly Arg Val Glu Ile Trp His Ala  
 1250 1255 1260

Gly Ser Trp Gly Thr Val Cys Asp Asp Ser Trp Asp Leu Ala Glu Ala  
 1265 1270 1275 1280

Glu Val Val Cys Gln Gln Leu Gly Cys Gly Ser Ala Leu Ala Ala Leu  
 1285 1290 1295

Arg Asp Ala Ser Phe Gly Gln Gly Thr Gly Thr Ile Trp Leu Asp Asp  
 1300 1305 1310

Met Arg Cys Lys Gly Asn Glu Ser Phe Leu Trp Asp Cys His Ala Lys  
 1315 1320 1325

Pro Trp Gly Gln Ser Asp Cys Gly His Lys Glu Asp Ala Gly Val Arg  
 1330 1335 1340

Cys Ser Gly Gln Ser Leu Lys Ser Leu Asn Ala Ser Ser Gly His Leu  
 1345 1350 1355 1360

Ala Leu Ile Leu Ser Ser Ile Phe Gly Leu Leu Leu Leu Val Leu Phe  
 1365 1370 1375

Ile Leu Phe Leu Thr Trp Cys Arg Val Gln Lys Gln Lys His Leu Pro  
 1380 1385 1390

Leu Arg Val Ser Thr Arg Arg Arg Gly Ser Leu Glu Glu Asn Leu Phe  
 1395 1400 1405

His Glu Met Glu Thr Cys Leu Lys Arg Glu Asp Pro His Gly Thr Arg  
 1410 1415 1420

Thr Ser Asp Asp Thr Pro Asn His Gly Cys Glu Asp Ala Ser Asp Thr  
 1425 1430 1435 1440

Ser Leu Leu Gly Val Leu Pro Ala Ser Glu Ala Thr Lys  
 1445 1450

<210> 12  
 <211> 40  
 <212> PRT  
 <213> Homo sapiens

<400> 12  
 Met Met Leu Pro Gln Asn Ser Trp His Ile Asp Phe Gly Arg Cys Cys  
 1 5 10 15

Cys His Gln Asn Leu Phe Ser Ala Val Val Thr Cys Ile Leu Leu Leu  
 20 25 30

Asn Ser Cys Phe Leu Ile Ser Ser  
 35 40

<210> 13  
 <211> 1413  
 <212> PRT

<213> Homo sapiens

<400> 13

Phe Asn Gly Thr Asp Leu Glu Leu Arg Leu Val Asn Gly Asp Gly Pro  
1 5 10 15

Cys Ser Gly Thr Val Glu Val Lys Phe Gln Gly Gln Trp Gly Thr Val  
20 25 30

Cys Asp Asp Gly Trp Asn Thr Thr Ala Ser Thr Val Val Cys Lys Gln  
35 40 45

Leu Gly Cys Pro Phe Ser Phe Ala Met Phe Arg Phe Gly Gln Ala Val  
50 55 60

Thr Arg His Gly Lys Ile Trp Leu Asp Asp Val Ser Cys Tyr Gly Asn  
65 70 75 80

Glu Ser Ala Leu Trp Glu Cys Gln His Arg Glu Trp Gly Ser His Asn  
85 90 95

Cys Tyr His Gly Glu Asp Val Gly Val Asn Cys Tyr Gly Glu Ala Asn  
100 105 110

Leu Gly Leu Arg Leu Val Asp Gly Asn Asn Ser Cys Ser Gly Arg Val  
115 120 125

Glu Val Lys Phe Gln Glu Arg Trp Gly Thr Ile Cys Asp Asp Gly Trp  
130 135 140

Asn Leu Asn Thr Ala Ala Val Val Cys Arg Gln Leu Gly Cys Pro Ser  
145 150 155 160

Ser Phe Ile Ser Ser Gly Val Val Asn Ser Pro Ala Val Leu Arg Pro  
165 170 175

Ile Trp Leu Asp Asp Ile Leu Cys Gln Gly Asn Glu Leu Ala Leu Trp  
180 185 190

Asn Cys Arg His Arg Gly Trp Gly Asn His Asp Cys Ser His Asn Glu  
195 200 205

Asp Val Thr Leu Thr Cys Tyr Asp Ser Ser Asp Leu Glu Leu Arg Leu  
210 215 220

Val Gly Gly Thr Asn Arg Cys Met Gly Arg Val Glu Leu Lys Ile Gln  
225 230 235 240



Pro Ile Trp Leu Asp Asp Val Ser Cys Ile Gly Asn Glu Ser Asn Ile  
 500 505 510

Trp Asp Cys Glu His Ser Gly Trp Gly Lys His Asn Cys Val His Arg  
 515 520 525

Glu Asp Val Ile Val Thr Cys Ser Gly Asp Ala Thr Trp Gly Leu Arg  
 530 535 540

Leu Val Gly Gly Ser Asn Arg Cys Ser Gly Arg Leu Glu Val Tyr Phe  
 545 550 555 560

Gln Gly Arg Trp Gly Thr Val Cys Asp Asp Gly Trp Asn Ser Lys Ala  
 565 570 575

Ala Ala Val Val Cys Ser Gln Leu Asp Cys Pro Ser Ser Ile Ile Gly  
 580 585 590

Met Gly Leu Gly Asn Ala Ser Thr Gly Tyr Gly Lys Ile Trp Leu Asp  
 595 600 605

Asp Val Ser Cys Asp Gly Asp Glu Ser Asp Leu Trp Ser Cys Arg Asn  
 610 615 620

Ser Gly Trp Gly Asn Asn Asp Cys Ser His Ser Glu Asp Val Gly Val  
 625 630 635 640

Ile Cys Ser Asp Ala Ser Asp Met Glu Leu Arg Leu Val Gly Gly Ser  
 645 650 655

Ser Arg Cys Ala Gly Lys Val Glu Val Asn Val Gln Gly Ala Val Gly  
 660 665 670

Ile Leu Cys Ala Asn Gly Trp Gly Met Asn Ile Ala Glu Val Val Cys  
 675 680 685

Arg Gln Leu Glu Cys Gly Ser Ala Ile Arg Val Ser Arg Glu Pro His  
 690 695 700

Phe Thr Glu Arg Thr Leu His Ile Leu Met Ser Asn Ser Gly Cys Thr  
 705 710 715 720

Gly Gly Glu Ala Ser Leu Trp Asp Cys Ile Arg Trp Glu Trp Lys Gln  
 725 730 735

Thr Ala Cys His Leu Asn Met Glu Ala Ser Leu Ile Cys Ser Ala His  
 740 745 750

Arg Gln Pro Arg Leu Val Gly Ala Asp Met Pro Cys Ser Gly Arg Val  
755 760 765

Glu Val Lys His Ala Asp Thr Trp Arg Ser Val Cys Asp Ser Asp Phe  
770 775 780

Ser Leu His Ala Ala Asn Val Leu Cys Arg Glu Leu Asn Cys Gly Asp  
785 790 795 800

Ala Ile Ser Leu Ser Val Gly Asp His Phe Gly Lys Gly Asn Gly Leu  
805 810 815

Thr Trp Ala Glu Lys Phe Gln Cys Glu Gly Ser Glu Thr His Leu Ala  
820 825 830

Leu Cys Pro Ile Val Gln His Pro Glu Asp Thr Cys Ile His Ser Arg  
835 840 845

Glu Val Gly Val Val Cys Ser Arg Tyr Thr Asp Val Arg Leu Val Asn  
850 855 860

Gly Lys Ser Gln Cys Asp Gly Gln Val Glu Ile Asn Val Leu Gly His  
865 870 875 880

Trp Gly Ser Leu Cys Asp Thr His Trp Asp Pro Glu Asp Ala Arg Val  
885 890 895

Leu Cys Arg Gln Leu Ser Cys Gly Thr Ala Leu Ser Thr Thr Gly Gly  
900 905 910

Lys Tyr Ile Gly Glu Arg Ser Val Arg Val Trp Gly His Arg Phe His  
915 920 925

Cys Leu Gly Asn Glu Ser Leu Leu Asp Asn Cys Gln Met Thr Val Leu  
930 935 940

Gly Ala Pro Pro Cys Ile His Gly Asn Thr Val Ser Val Ile Cys Thr  
945 950 955 960

Gly Ser Leu Thr Gln Pro Leu Phe Pro Cys Leu Ala Asn Val Ser Asp  
965 970 975

Pro Tyr Leu Ser Ala Val Pro Glu Gly Ser Ala Leu Ile Cys Leu Glu  
980 985 990

Asp Lys Arg Leu Arg Leu Val Asp Gly Asp Ser Arg Cys Ala Gly Arg  
995 1000 1005

Val Glu Ile Tyr His Asp Gly Phe Trp Gly Thr Ile Cys Asp Asp Gly  
 1010 1015 1020  
 Trp Asp Leu Ser Asp Ala His Val Val Cys Gln Lys Leu Gly Cys Gly  
 1025 1030 1035 1040  
 Val Ala Phe Asn Ala Thr Val Ser Ala His Phe Gly Glu Gly Ser Gly  
 1045 1050 1055  
 Pro Ile Trp Leu Asp Asp Leu Asn Cys Thr Gly Thr Glu Ser His Leu  
 1060 1065 1070  
 Trp Gln Cys Pro Ser Arg Gly Trp Gly Gln His Asp Cys Arg His Lys  
 1075 1080 1085  
 Glu Asp Ala Gly Val Ile Cys Ser Glu Phe Thr Ala Leu Arg Leu Tyr  
 1090 1095 1100  
 Ser Glu Thr Glu Thr Glu Ser Cys Ala Gly Arg Leu Glu Val Phe Tyr  
 1105 1110 1115 1120  
 Asn Gly Thr Trp Gly Ser Val Gly Arg Arg Asn Ile Thr Thr Ala Ile  
 1125 1130 1135  
 Ala Gly Ile Val Cys Arg Gln Leu Gly Cys Gly Glu Asn Gly Val Val  
 1140 1145 1150  
 Ser Leu Ala Pro Leu Ser Lys Thr Gly Ser Gly Phe Met Trp Val Asp  
 1155 1160 1165  
 Asp Ile Gln Cys Pro Lys Thr His Ile Ser Ile Trp Gln Cys Leu Ser  
 1170 1175 1180  
 Ala Pro Trp Glu Arg Arg Ile Ser Ser Pro Ala Glu Glu Thr Trp Ile  
 1185 1190 1195 1200  
 Thr Cys Glu Asp Arg Ile Arg Val Arg Gly Gly Asp Thr Glu Cys Ser  
 1205 1210 1215  
 Gly Arg Val Glu Ile Trp His Ala Gly Ser Trp Gly Thr Val Cys Asp  
 1220 1225 1230  
 Asp Ser Trp Asp Leu Ala Glu Ala Glu Val Val Cys Gln Gln Leu Gly  
 1235 1240 1245  
 Cys Gly Ser Ala Leu Ala Ala Leu Arg Asp Ala Ser Phe Gly Gln Gly  
 1250 1255 1260

Thr Gly Thr Ile Trp Leu Asp Asp Met Arg Cys Lys Gly Asn Glu Ser  
1265 1270 1275 1280

Phe Leu Trp Asp Cys His Ala Lys Pro Trp Gly Gln Ser Asp Cys Gly  
1285 1290 1295

His Lys Glu Asp Ala Gly Val Arg Cys Ser Gly Gln Ser Leu Lys Ser  
1300 1305 1310

Leu Asn Ala Ser Ser Gly His Leu Ala Leu Ile Leu Ser Ser Ile Phe  
1315 1320 1325

Gly Leu Leu Leu Leu Val Leu Phe Ile Leu Phe Leu Thr Trp Cys Arg  
1330 1335 1340

Val Gln Lys Gln Lys His Leu Pro Leu Arg Val Ser Thr Arg Arg Arg  
1345 1350 1355 1360

Gly Ser Leu Glu Glu Asn Leu Phe His Glu Met Glu Thr Cys Leu Lys  
1365 1370 1375

Arg Glu Asp Pro His Gly Thr Arg Thr Ser Asp Asp Thr Pro Asn His  
1380 1385 1390

Gly Cys Glu Asp Ala Ser Asp Thr Ser Leu Leu Gly Val Leu Pro Ala  
1395 1400 1405

Ser Glu Ala Thr Lys  
1410

<210> 14

<211> 1319

<212> PRT

<213> Homo sapiens

<400> 14

Phe Asn Gly Thr Asp Leu Glu Leu Arg Leu Val Asn Gly Asp Gly Pro  
1 5 10 15

Cys Ser Gly Thr Val Glu Val Lys Phe Gln Gly Gln Trp Gly Thr Val  
20 25 30

Cys Asp Asp Gly Trp Asn Thr Thr Ala Ser Thr Val Val Cys Lys Gln  
35 40 45

Leu Gly Cys Pro Phe Ser Phe Ala Met Phe Arg Phe Gly Gln Ala Val  
50 55 60

Thr Arg His Gly Lys Ile Trp Leu Asp Asp Val Ser Cys Tyr Gly Asn  
65 70 75 80

Glu Ser Ala Leu Trp Glu Cys Gln His Arg Glu Trp Gly Ser His Asn  
85 90 95

Cys Tyr His Gly Glu Asp Val Gly Val Asn Cys Tyr Gly Glu Ala Asn  
100 105 110

Leu Gly Leu Arg Leu Val Asp Gly Asn Asn Ser Cys Ser Gly Arg Val  
115 120 125

Glu Val Lys Phe Gln Glu Arg Trp Gly Thr Ile Cys Asp Asp Gly Trp  
130 135 140

Asn Leu Asn Thr Ala Ala Val Val Cys Arg Gln Leu Gly Cys Pro Ser  
145 150 155 160

Ser Phe Ile Ser Ser Gly Val Val Asn Ser Pro Ala Val Leu Arg Pro  
165 170 175

Ile Trp Leu Asp Asp Ile Leu Cys Gln Gly Asn Glu Leu Ala Leu Trp  
180 185 190

Asn Cys Arg His Arg Gly Trp Gly Asn His Asp Cys Ser His Asn Glu  
195 200 205

Asp Val Thr Leu Thr Cys Tyr Asp Ser Ser Asp Leu Glu Leu Arg Leu  
210 215 220

Val Gly Gly Thr Asn Arg Cys Met Gly Arg Val Glu Leu Lys Ile Gln  
225 230 235 240

Gly Arg Trp Gly Thr Val Cys His His Lys Trp Asn Asn Ala Ala Ala  
245 250 255

Asp Val Val Cys Lys Gln Leu Gly Cys Gly Thr Ala Leu His Phe Ala  
260 265 270

Gly Leu Pro His Leu Gln Ser Gly Ser Asp Val Val Trp Leu Asp Gly  
275 280 285

Val Ser Cys Ser Gly Asn Glu Ser Phe Leu Trp Asp Cys Arg His Ser  
290 295 300

Gly Thr Val Asn Phe Asp Cys Leu His Gln Asn Asp Val Ser Val Ile  
305 310 315 320

Cys Ser Asp Gly Ala Asp Leu Glu Leu Arg Leu Ala Asp Gly Ser Asn  
 325 330 335

Asn Cys Ser Gly Arg Val Glu Val Arg Ile His Glu Gln Trp Trp Thr  
 340 345 350

Ile Cys Asp Gln Asn Trp Lys Asn Glu Gln Ala Leu Val Val Cys Lys  
 355 360 365

Gln Leu Gly Cys Pro Phe Ser Val Phe Gly Ser Arg Arg Ala Lys Pro  
 370 375 380

Ser Asn Glu Ala Arg Asp Ile Trp Ile Asn Ser Ile Ser Cys Thr Gly  
 385 390 395 400

Asn Glu Ser Ala Leu Trp Asp Cys Thr Tyr Asp Gly Lys Ala Lys Arg  
 405 410 415

Thr Cys Phe Arg Arg Ser Asp Ala Gly Val Ile Cys Ser Asp Lys Ala  
 420 425 430

Asp Leu Asp Leu Arg Leu Val Gly Ala His Ser Pro Cys Tyr Gly Arg  
 435 440 445

Leu Glu Val Lys Tyr Gln Gly Glu Trp Gly Thr Val Cys His Asp Arg  
 450 455 460

Trp Ser Thr Arg Asn Ala Ala Val Val Cys Lys Gln Leu Gly Cys Gly  
 465 470 475 480

Lys Pro Met His Val Phe Gly Met Thr Tyr Phe Lys Glu Ala Ser Gly  
 485 490 495

Pro Ile Trp Leu Asp Asp Val Ser Cys Ile Gly Asn Glu Ser Asn Ile  
 500 505 510

Trp Asp Cys Glu His Ser Gly Trp Gly Lys His Asn Cys Val His Arg  
 515 520 525

Glu Asp Val Ile Val Thr Cys Ser Gly Asp Ala Thr Trp Gly Leu Arg  
 530 535 540

Leu Val Gly Gly Ser Asn Arg Cys Ser Gly Arg Leu Glu Val Tyr Phe  
 545 550 555 560

Gln Gly Arg Trp Gly Thr Val Cys Asp Asp Gly Trp Asn Ser Lys Ala  
 565 570 575

Ala Ala Val Val Cys Ser Gln Leu Asp Cys Pro Ser Ser Ile Ile Gly  
580 585 590

Met Gly Leu Gly Asn Ala Ser Thr Gly Tyr Gly Lys Ile Trp Leu Asp  
595 600 605

Asp Val Ser Cys Asp Gly Asp Glu Ser Asp Leu Trp Ser Cys Arg Asn  
610 615 620

Ser Gly Trp Gly Asn Asn Asp Cys Ser His Ser Glu Asp Val Gly Val  
625 630 635 640

Ile Cys Ser Asp Ala Ser Asp Met Glu Leu Arg Leu Val Gly Gly Ser  
645 650 655

Ser Arg Cys Ala Gly Lys Val Glu Val Asn Val Gln Gly Ala Val Gly  
660 665 670

Ile Leu Cys Ala Asn Gly Trp Gly Met Asn Ile Ala Glu Val Val Cys  
675 680 685

Arg Gln Leu Glu Cys Gly Ser Ala Ile Arg Val Ser Arg Glu Pro His  
690 695 700

Phe Thr Glu Arg Thr Leu His Ile Leu Met Ser Asn Ser Gly Cys Thr  
705 710 715 720

Gly Gly Glu Ala Ser Leu Trp Asp Cys Ile Arg Trp Glu Trp Lys Gln  
725 730 735

Thr Ala Cys His Leu Asn Met Glu Ala Ser Leu Ile Cys Ser Ala His  
740 745 750

Arg Gln Pro Arg Leu Val Gly Ala Asp Met Pro Cys Ser Gly Arg Val  
755 760 765

Glu Val Lys His Ala Asp Thr Trp Arg Ser Val Cys Asp Ser Asp Phe  
770 775 780

Ser Leu His Ala Ala Asn Val Leu Cys Arg Glu Leu Asn Cys Gly Asp  
785 790 795 800

Ala Ile Ser Leu Ser Val Gly Asp His Phe Gly Lys Gly Asn Gly Leu  
805 810 815

Thr Trp Ala Glu Lys Phe Gln Cys Glu Gly Ser Glu Thr His Leu Ala  
820 825 830

Leu Cys Pro Ile Val Gln His Pro Glu Asp Thr Cys Ile His Ser Arg  
835 840 845

Glu Val Gly Val Val Cys Ser Arg Tyr Thr Asp Val Arg Leu Val Asn  
850 855 860

Gly Lys Ser Gln Cys Asp Gly Gln Val Glu Ile Asn Val Leu Gly His  
865 870 875 880

Trp Gly Ser Leu Cys Asp Thr His Trp Asp Pro Glu Asp Ala Arg Val  
885 890 895

Leu Cys Arg Gln Leu Ser Cys Gly Thr Ala Leu Ser Thr Thr Gly Gly  
900 905 910

Lys Tyr Ile Gly Glu Arg Ser Val Arg Val Trp Gly His Arg Phe His  
915 920 925

Cys Leu Gly Asn Glu Ser Leu Leu Asp Asn Cys Gln Met Thr Val Leu  
930 935 940

Gly Ala Pro Pro Cys Ile His Gly Asn Thr Val Ser Val Ile Cys Thr  
945 950 955 960

Gly Ser Leu Thr Gln Pro Leu Phe Pro Cys Leu Ala Asn Val Ser Asp  
965 970 975

Pro Tyr Leu Ser Ala Val Pro Glu Gly Ser Ala Leu Ile Cys Leu Glu  
980 985 990

Asp Lys Arg Leu Arg Leu Val Asp Gly Asp Ser Arg Cys Ala Gly Arg  
995 1000 1005

Val Glu Ile Tyr His Asp Gly Phe Trp Gly Thr Ile Cys Asp Asp Gly  
1010 1015 1020

Trp Asp Leu Ser Asp Ala His Val Val Cys Gln Lys Leu Gly Cys Gly  
1025 1030 1035 1040

Val Ala Phe Asn Ala Thr Val Ser Ala His Phe Gly Glu Gly Ser Gly  
1045 1050 1055

Pro Ile Trp Leu Asp Asp Leu Asn Cys Thr Gly Thr Glu Ser His Leu  
1060 1065 1070

Trp Gln Cys Pro Ser Arg Gly Trp Gly Gln His Asp Cys Arg His Lys  
1075 1080 1085

Glu Asp Ala Gly Val Ile Cys Ser Glu Phe Thr Ala Leu Arg Leu Tyr  
 1090 1095 1100

Ser Glu Thr Glu Thr Glu Ser Cys Ala Gly Arg Leu Glu Val Phe Tyr  
 1105 1110 1115 1120

Asn Gly Thr Trp Gly Ser Val Gly Arg Arg Asn Ile Thr Thr Ala Ile  
 1125 1130 1135

Ala Gly Ile Val Cys Arg Gln Leu Gly Cys Gly Glu Asn Gly Val Val  
 1140 1145 1150

Ser Leu Ala Pro Leu Ser Lys Thr Gly Ser Gly Phe Met Trp Val Asp  
 1155 1160 1165

Asp Ile Gln Cys Pro Lys Thr His Ile Ser Ile Trp Gln Cys Leu Ser  
 1170 1175 1180

Ala Pro Trp Glu Arg Arg Ile Ser Ser Pro Ala Glu Glu Thr Trp Ile  
 1185 1190 1195 1200

Thr Cys Glu Asp Arg Ile Arg Val Arg Gly Gly Asp Thr Glu Cys Ser  
 1205 1210 1215

Gly Arg Val Glu Ile Trp His Ala Gly Ser Trp Gly Thr Val Cys Asp  
 1220 1225 1230

Asp Ser Trp Asp Leu Ala Glu Ala Glu Val Val Cys Gln Gln Leu Gly  
 1235 1240 1245

Cys Gly Ser Ala Leu Ala Ala Leu Arg Asp Ala Ser Phe Gly Gln Gly  
 1250 1255 1260

Thr Gly Thr Ile Trp Leu Asp Asp Met Arg Cys Lys Gly Asn Glu Ser  
 1265 1270 1275 1280

Phe Leu Trp Asp Cys His Ala Lys Pro Trp Gly Gln Ser Asp Cys Gly  
 1285 1290 1295

His Lys Glu Asp Ala Gly Val Arg Cys Ser Gly Gln Ser Leu Lys Ser  
 1300 1305 1310

Leu Asn Ala Ser Ser Gly His  
 1315

<210> 15

<211> 24  
 <212> PRT  
 <213> Homo sapiens

<400> 15  
 Leu Ala Leu Ile Leu Ser Ser Ile Phe Gly Leu Leu Leu Leu Val Leu  
 1 5 10 15

Phe Ile Leu Phe Leu Thr Trp Cys  
 20

<210> 16  
 <211> 70  
 <212> PRT  
 <213> Homo sapiens

<400> 16  
 Arg Val Gln Lys Gln Lys His Leu Pro Leu Arg Val Ser Thr Arg Arg  
 1 5 10 15

Arg Gly Ser Leu Glu Glu Asn Leu Phe His Glu Met Glu Thr Cys Leu  
 20 25 30

Lys Arg Glu Asp Pro His Gly Thr Arg Thr Ser Asp Asp Thr Pro Asn  
 35 40 45

His Gly Cys Glu Asp Ala Ser Asp Thr Ser Leu Leu Gly Val Leu Pro  
 50 55 60

Ala Ser Glu Ala Thr Lys  
 65 70

<210> 17  
 <211> 3104  
 <212> DNA  
 <213> Homo sapiens

<400> 17  
 gtcgacccac gcgtcoggtc tgtggctgag catggccctc ccagccctgg gcctggaccc 60  
 ctggagcctc ctgggccttt tcctcttcca actgcttcag ctgctgctgc cgacgacgac 120  
 cgcgggggga ggcgggcagg ggcccatgcc cagggtcaga tactatgcag gggatgaacg 180  
 tagggcactt agcttcttcc accagaaggg cctccaggat tttgacactc tgctcctgag 240  
 tggatgatgga aatactctct acgtgggggc tcgagaagcc attctggcct tggatatcca 300  
 ggatccaggg gtccccaggc taaagaacat gataccgtgg ccagccagtg acagaaaaaa 360  
 gagtgaatgt gcctttaaga agaagagcaa tgagacacag tgtttcaact tcatccgtgt 420  
 cctggtttct tacaatgtca cccatctcta cacctgcggc accttcgcct tcagccctgc 480

```

ttgtaccttc attgaacttc aagattccta cctggtgcc atctcggagg acaaggatcat 540
ggaggggaaaa ggccaaagcc cctttgaccc cgctcacaag catacggctg tcttggtgga 600
tgggatgctc tattctggta ctatgaacaa ctccctgggc agtgagccca tcctgatgcg 660
cacactggga tcccagcctg tcctcaagac cgacaacttc ctccgctggc tgcacatga 720
cgcctccttt gtggcagcca tcccttcgac ccaggtcgtc tactttctct tcgaggagac 780
agccagcgag tttgacttct ttgagaggct ccacacatcg cgggtggcta gagtctgcaa 840
gaatgacgtg ggccggcgaaa agctgctgca gaagaagtgg accaccttcc tgaaggccca 900
gctgctctgc acccagccgg ggagctgcc cttcaacgtc atccgccacg cggctcctgct 960
ccccgcgat tctccacag ctcccccacat ctacgcagtc ttcacctccc agtggcaggt 1020
tggcgggacc aggagctctg cggtttgtgc cttctctctc ttggacattg aacgtgtctt 1080
taaggggaaa taaaaagagt tgaacaaaga aacttcacgc tggactactt ataggggccc 1140
tgagaccaac ccccgccag gcagttgctc agtgggcccc tcctctgata aggccctgac 1200
cttcatgaag gaccatttcc tgatggatga gcaagtgggt gggacgcccc tgctggtgaa 1260
atctggcgtg gagtatacac ggcttgcat ggagacagcc cagggccttg atgggcacag 1320
ccatcttgct atgtacctgg gaaccaccac agggctcgtc cacaaggctg tggtaagtgg 1380
ggacagcagt gctcatctgg tggaaagat tcagctgttc cctgacctg aacctgttcg 1440
caacctgcag ctggccccc cccagggtgc agtgtttgta ggcttctcag gaggtgtctg 1500
gaggggtgcc cgagccaact gtagtgtcta tgagagctgt gtggactgtg tccttgcccg 1560
ggacccccac tgtgcctggg acctgagtc ccgaacctgt tgcctcctgt ctgcccccaa 1620
cctgaactcc tggaaagcagg acatggagcg ggggaaccca gagtgggcat gtgccagtgg 1680
ccccatgagc aggagccttc ggctcagag ccgcccga atcattaaag aagtccctggc 1740
tgtcccaac tccatcctgg agctcccctg cccccacctg tcagccttg cctcttatta 1800
ttggagtc at ggcccagcag cagtcccaga agcctcttcc actgtctaca atggctccct 1860
cttgctgata gtgcaggatg gagttggggg tctctaccag tgctgggcaa ctgagaatgg 1920
cttttcatac cctgtgatct cctactgggt ggacagccag gaccagacc tgccctgga 1980
tcctgaactg gcaggcatcc cccgggagca tgtgaaggct ccgttgacca gggtcagtgg 2040
tggggccgcc ctggctgccc agcagtccta ctggccccac tttgtcactg tcaactgtcct 2100
ctttgcctta gtgctttcag gagccctcat catcctcgtg gcctcccat tgagagcact 2160
ccgggctcgg ggcaagggttc agggctgtga gaccctgcgc cctggggaga agggcccggt 2220
aagcagagag caacacctcc agtctcccaa ggaatgcagg acctctgcca gtgatgtgga 2280
cgctgacaac aactgcctag gcactgaggt agcttaaaact ctaggcacag gccggggctg 2340
cgggtgcaggc acctggccat gctggctggg cggcccaagc acagccctga ctaggatgac 2400
agcagcaca aagaccacct ttctccctg agaggagctt ctgctactct gcatcactga 2460
tgacactcag cagggtgatg cacagcagtc tgcctccct atgggactcc cttctaccaa 2520
gcacatgagc tctctaacag ggtgggggct acccccagac ctgctcctac actgatattg 2580
aagaacctgg agaggatcct tcagttctgg ccattccagg gaccctccag aaacacagt 2640
tttcaagaga tcctaaaaaa acctgcctgt ccaggaccc tatggtaatg aacaccaaac 2700
atctaaacaa tcatatgcta acatgccact cctggaaaact ccaacttgaa gctgcgctt 2760
tggacaccaa cactcccttc tcccagggtc atgcagggat ctgctccctc ctgcttccct 2820
taccagtctg gcaccgctga ctcccaggaa gtctttcctg aagtctgacc acctttcttc 2880
ttgcttcagt tggggcagac tctgatccct tctgccctgg cagaatggca ggggtaatct 2940
gagccttctt cactccttta cctagctga ccccttcacc tctccccctc ccttttctt 3000
tgttttggga ttcagaaaac tgctgtcag agactgttta ttttttatta aaaatataag 3060
gcttaaaaaa aaaaaaaaaa aaaaaaaaaa aaaagggcgg ccgc 3104

```

<210> 18  
 <211> 2283  
 <212> DNA

<213> Homo sapiens

<400> 18

atggccctcc cagccctggg cctggacccc tggagcctcc tgggcctttt cctcttccaa 60  
ctgcttcagc tgctgctgcc gacgacgacc gcggggggag gcgggcaggg gcccatgccc 120  
agggtcagat actatgcagg ggatgaacgt agggcactta gcttcttcca ccagaagggc 180  
ctccaggatt ttgacactct gctcctgagt ggtgatggaa atactctcta cgtgggggct 240  
cgagaagcca ttctggcctt ggatatccag gatccagggg tccccaggct aaagaacatg 300  
ataccgtggc cagccagtga cagaaaaaag agtgaatgtg cctttaagaa gaagagcaat 360  
gagacacagt gtttcaactt catccgtgtc ctggtttctt acaatgtcac ccatctctac 420  
acctgcggca ccttcgcctt cagccctgct tgtaccttca ttgaacttca agattcctac 480  
ctggttgcca tctcgaggga caaggtcatg gagggaaaag gccaaagccc ctttgacccc 540  
gctcacaagc atacggctgt cttggtggat gggatgctct attctggtac tatgaacaac 600  
ttcctgggca gtgagcccat cctgatgctc aactgggat ccagcctgt cctcaagacc 660  
gacaacttcc tccgctggct gcatcatgac gcctcctttg tggcagccat cccttcgacc 720  
caggtcgtct acttcttctt cgaggagaca gccagcgagt ttgacttctt tgagaggctc 780  
cacacatcgc ggggtggctag agtctgcaag aatgacgtgg gcggcgaaaa gctgctgcag 840  
aagaagtgga ccaccttctt gaaggcccag ctgctctgca ccagccggg gcagctgccc 900  
ttcaacgtca tccgccacgc ggtcctgctc ccgcgcgatt ctcccacagc tcccacatc 960  
tacgcagtct tcacctcca gtggcagggt ggcgggacca ggagctctgc ggtttgtgcc 1020  
ttctctctct tggacattga acgtgtcttt aaggggaaat acaaagagtt gaacaaagaa 1080  
acttcacgct ggactactta taggggccct gagaccaacc ccggccagg cagttgctca 1140  
gtgggccccct cctctgataa ggccctgacc ttcataagag accatttctt gatggatgag 1200  
caagtgggtg ggacgccccct gctggtgaaa tctggcgtgg agtatacacg gcttgacagt 1260  
gagacagccc agggccttga tgggcacagc catcttgctc tgtacctggg aaccaccaca 1320  
gggtcgctcc acaaggctgt ggtaagtggg gacagcagt ctcatctggt ggaagagatt 1380  
cagctgttcc ctgacctga acctgttcgc aacctgcagc tggccccac ccagggtgca 1440  
gtgtttgtag gcttctcagg aggtgtctgg agggtgcccc gagccaaact tagtgtctat 1500  
gagagctgtg tggactgtgt ccttgccccg gacccccact gtgcctggga ccctgagtc 1560  
cgaacctgtt gcctcctgtc tgcccccaac ctgaactcct ggaagcagga catggagcgg 1620  
gggaacccag agtgggcatg tgccagtggc cccatgagca ggagccttcg gcctcagagc 1680  
cgcccgcaaa tcattaaaga agtctgtggt gtccccaact ccatcctgga gctcccctgc 1740  
ccccacctgt cagccttggc ctcttattat tggagtcatg gcccagcagc agtcccagaa 1800  
gcctcttcca ctgtctacaa tggctccctc ttgctgatag tgcaggatgg agttgggggt 1860  
ctctaccagt gctgggcaac tgagaatggc ttttcatacc ctgtgatctc ctactgggtg 1920  
gacagccagg accagaccct ggccctggat cctgaactgg caggcatccc ccgggagcat 1980  
gtgaaggctc cgttgaccag ggtcagtggg ggggcccc tggctgccc gcagtcctac 2040  
tgccccact ttgtcactgt cactgtcctc tttgccttag tgctttcagg agccctcatc 2100  
atcctcgtgg cctccccatt gagagcactc cgggctcggg gcaaggttca gggctgtgag 2160  
accctgcgcc ctggggagaa ggccccgtta agcagagagc aacacctcca gtctcccaag 2220  
gaatgcagga cctctgccag tgatgtggac gctgacaaca actgcctagg cactgaggta 2280  
gct 2283

<210> 19

<211> 761

<212> PRT

<213> Homo sapiens

<400> 19

Met Ala Leu Pro Ala Leu Gly Leu Asp Pro Trp Ser Leu Leu Gly Leu  
1 5 10 15

Phe Leu Phe Gln Leu Leu Gln Leu Leu Leu Pro Thr Thr Thr Ala Gly  
20 25 30

Gly Gly Gly Gln Gly Pro Met Pro Arg Val Arg Tyr Tyr Ala Gly Asp  
35 40 45

Glu Arg Arg Ala Leu Ser Phe Phe His Gln Lys Gly Leu Gln Asp Phe  
50 55 60

Asp Thr Leu Leu Leu Ser Gly Asp Gly Asn Thr Leu Tyr Val Gly Ala  
65 70 75 80

Arg Glu Ala Ile Leu Ala Leu Asp Ile Gln Asp Pro Gly Val Pro Arg  
85 90 95

Leu Lys Asn Met Ile Pro Trp Pro Ala Ser Asp Arg Lys Lys Ser Glu  
100 105 110

Cys Ala Phe Lys Lys Lys Ser Asn Glu Thr Gln Cys Phe Asn Phe Ile  
115 120 125

Arg Val Leu Val Ser Tyr Asn Val Thr His Leu Tyr Thr Cys Gly Thr  
130 135 140

Phe Ala Phe Ser Pro Ala Cys Thr Phe Ile Glu Leu Gln Asp Ser Tyr  
145 150 155 160

Leu Leu Pro Ile Ser Glu Asp Lys Val Met Glu Gly Lys Gly Gln Ser  
165 170 175

Pro Phe Asp Pro Ala His Lys His Thr Ala Val Leu Val Asp Gly Met  
180 185 190

Leu Tyr Ser Gly Thr Met Asn Asn Phe Leu Gly Ser Glu Pro Ile Leu  
195 200 205

Met Arg Thr Leu Gly Ser Gln Pro Val Leu Lys Thr Asp Asn Phe Leu  
210 215 220

Arg Trp Leu His His Asp Ala Ser Phe Val Ala Ala Ile Pro Ser Thr  
225 230 235 240

Gln Val Val Tyr Phe Phe Phe Glu Glu Thr Ala Ser Glu Phe Asp Phe  
245 250 255

Phe Glu Arg Leu His Thr Ser Arg Val Ala Arg Val Cys Lys Asn Asp  
 260 265 270

Val Gly Gly Glu Lys Leu Leu Gln Lys Lys Trp Thr Thr Phe Leu Lys  
 275 280 285

Ala Gln Leu Leu Cys Thr Gln Pro Gly Gln Leu Pro Phe Asn Val Ile  
 290 295 300

Arg His Ala Val Leu Leu Pro Ala Asp Ser Pro Thr Ala Pro His Ile  
 305 310 315 320

Tyr Ala Val Phe Thr Ser Gln Trp Gln Val Gly Gly Thr Arg Ser Ser  
 325 330 335

Ala Val Cys Ala Phe Ser Leu Leu Asp Ile Glu Arg Val Phe Lys Gly  
 340 345 350

Lys Tyr Lys Glu Leu Asn Lys Glu Thr Ser Arg Trp Thr Thr Tyr Arg  
 355 360 365

Gly Pro Glu Thr Asn Pro Arg Pro Gly Ser Cys Ser Val Gly Pro Ser  
 370 375 380

Ser Asp Lys Ala Leu Thr Phe Met Lys Asp His Phe Leu Met Asp Glu  
 385 390 395 400

Gln Val Val Gly Thr Pro Leu Leu Val Lys Ser Gly Val Glu Tyr Thr  
 405 410 415

Arg Leu Ala Val Glu Thr Ala Gln Gly Leu Asp Gly His Ser His Leu  
 420 425 430

Val Met Tyr Leu Gly Thr Thr Thr Gly Ser Leu His Lys Ala Val Val  
 435 440 445

Ser Gly Asp Ser Ser Ala His Leu Val Glu Glu Ile Gln Leu Phe Pro  
 450 455 460

Asp Pro Glu Pro Val Arg Asn Leu Gln Leu Ala Pro Thr Gln Gly Ala  
 465 470 475 480

Val Phe Val Gly Phe Ser Gly Gly Val Trp Arg Val Pro Arg Ala Asn  
 485 490 495

Cys Ser Val Tyr Glu Ser Cys Val Asp Cys Val Leu Ala Arg Asp Pro  
 500 505 510

His Cys Ala Trp Asp Pro Glu Ser Arg Thr Cys Cys Leu Leu Ser Ala  
515 520 525

Pro Asn Leu Asn Ser Trp Lys Gln Asp Met Glu Arg Gly Asn Pro Glu  
530 535 540

Trp Ala Cys Ala Ser Gly Pro Met Ser Arg Ser Leu Arg Pro Gln Ser  
545 550 555 560

Arg Pro Gln Ile Ile Lys Glu Val Leu Ala Val Pro Asn Ser Ile Leu  
565 570 575

Glu Leu Pro Cys Pro His Leu Ser Ala Leu Ala Ser Tyr Tyr Trp Ser  
580 585 590

His Gly Pro Ala Ala Val Pro Glu Ala Ser Ser Thr Val Tyr Asn Gly  
595 600 605

Ser Leu Leu Leu Ile Val Gln Asp Gly Val Gly Gly Leu Tyr Gln Cys  
610 615 620

Trp Ala Thr Glu Asn Gly Phe Ser Tyr Pro Val Ile Ser Tyr Trp Val  
625 630 635 640

Asp Ser Gln Asp Gln Thr Leu Ala Leu Asp Pro Glu Leu Ala Gly Ile  
645 650 655

Pro Arg Glu His Val Lys Val Pro Leu Thr Arg Val Ser Gly Gly Ala  
660 665 670

Ala Leu Ala Ala Gln Gln Ser Tyr Trp Pro His Phe Val Thr Val Thr  
675 680 685

Val Leu Phe Ala Leu Val Leu Ser Gly Ala Leu Ile Ile Leu Val Ala  
690 695 700

Ser Pro Leu Arg Ala Leu Arg Ala Arg Gly Lys Val Gln Gly Cys Glu  
705 710 715 720

Thr Leu Arg Pro Gly Glu Lys Ala Pro Leu Ser Arg Glu Gln His Leu  
725 730 735

Gln Ser Pro Lys Glu Cys Arg Thr Ser Ala Ser Asp Val Asp Ala Asp  
740 745 750

Asn Asn Cys Leu Gly Thr Glu Val Ala  
755 760

<210> 20  
 <211> 31  
 <212> PRT  
 <213> Homo sapiens

<400> 20  
 Met Ala Leu Pro Ala Leu Gly Leu Asp Pro Trp Ser Leu Leu Gly Leu  
           1                  5                  10                  15

Phe Leu Phe Gln Leu Leu Gln Leu Leu Leu Pro Thr Thr Thr Ala  
                   20                  25                  30

<210> 21  
 <211> 730  
 <212> PRT  
 <213> Homo sapiens

<400> 21  
 Gly Gly Gly Gly Gln Gly Pro Met Pro Arg Val Arg Tyr Tyr Ala Gly  
           1                  5                  10                  15

Asp Glu Arg Arg Ala Leu Ser Phe Phe His Gln Lys Gly Leu Gln Asp  
                   20                  25                  30

Phe Asp Thr Leu Leu Leu Ser Gly Asp Gly Asn Thr Leu Tyr Val Gly  
                   35                  40                  45

Ala Arg Glu Ala Ile Leu Ala Leu Asp Ile Gln Asp Pro Gly Val Pro  
           50                  55                  60

Arg Leu Lys Asn Met Ile Pro Trp Pro Ala Ser Asp Arg Lys Lys Ser  
           65                  70                  75                  80

Glu Cys Ala Phe Lys Lys Lys Ser Asn Glu Thr Gln Cys Phe Asn Phe  
                   85                  90                  95

Ile Arg Val Leu Val Ser Tyr Asn Val Thr His Leu Tyr Thr Cys Gly  
           100                  105                  110

Thr Phe Ala Phe Ser Pro Ala Cys Thr Phe Ile Glu Leu Gln Asp Ser  
           115                  120                  125

Tyr Leu Leu Pro Ile Ser Glu Asp Lys Val Met Glu Gly Lys Gly Gln  
           130                  135                  140

Ser Pro Phe Asp Pro Ala His Lys His Thr Ala Val Leu Val Asp Gly  
 145 150 155 160  
 Met Leu Tyr Ser Gly Thr Met Asn Asn Phe Leu Gly Ser Glu Pro Ile  
 165 170 175  
 Leu Met Arg Thr Leu Gly Ser Gln Pro Val Leu Lys Thr Asp Asn Phe  
 180 185 190  
 Leu Arg Trp Leu His His Asp Ala Ser Phe Val Ala Ala Ile Pro Ser  
 195 200 205  
 Thr Gln Val Val Tyr Phe Phe Phe Glu Glu Thr Ala Ser Glu Phe Asp  
 210 215 220  
 Phe Phe Glu Arg Leu His Thr Ser Arg Val Ala Arg Val Cys Lys Asn  
 225 230 235 240  
 Asp Val Gly Gly Glu Lys Leu Leu Gln Lys Lys Trp Thr Thr Phe Leu  
 245 250 255  
 Lys Ala Gln Leu Leu Cys Thr Gln Pro Gly Gln Leu Pro Phe Asn Val  
 260 265 270  
 Ile Arg His Ala Val Leu Leu Pro Ala Asp Ser Pro Thr Ala Pro His  
 275 280 285  
 Ile Tyr Ala Val Phe Thr Ser Gln Trp Gln Val Gly Gly Thr Arg Ser  
 290 295 300  
 Ser Ala Val Cys Ala Phe Ser Leu Leu Asp Ile Glu Arg Val Phe Lys  
 305 310 315 320  
 Gly Lys Tyr Lys Glu Leu Asn Lys Glu Thr Ser Arg Trp Thr Thr Tyr  
 325 330 335  
 Arg Gly Pro Glu Thr Asn Pro Arg Pro Gly Ser Cys Ser Val Gly Pro  
 340 345 350  
 Ser Ser Asp Lys Ala Leu Thr Phe Met Lys Asp His Phe Leu Met Asp  
 355 360 365  
 Glu Gln Val Val Gly Thr Pro Leu Leu Val Lys Ser Gly Val Glu Tyr  
 370 375 380  
 Thr Arg Leu Ala Val Glu Thr Ala Gln Gly Leu Asp Gly His Ser His  
 385 390 395 400

Leu Val Met Tyr Leu Gly Thr Thr Thr Gly Ser Leu His Lys Ala Val  
 405 410 415

Val Ser Gly Asp Ser Ser Ala His Leu Val Glu Glu Ile Gln Leu Phe  
 420 425 430

Pro Asp Pro Glu Pro Val Arg Asn Leu Gln Leu Ala Pro Thr Gln Gly  
 435 440 445

Ala Val Phe Val Gly Phe Ser Gly Gly Val Trp Arg Val Pro Arg Ala  
 450 455 460

Asn Cys Ser Val Tyr Glu Ser Cys Val Asp Cys Val Leu Ala Arg Asp  
 465 470 475 480

Pro His Cys Ala Trp Asp Pro Glu Ser Arg Thr Cys Cys Leu Leu Ser  
 485 490 495

Ala Pro Asn Leu Asn Ser Trp Lys Gln Asp Met Glu Arg Gly Asn Pro  
 500 505 510

Glu Trp Ala Cys Ala Ser Gly Pro Met Ser Arg Ser Leu Arg Pro Gln  
 515 520 525

Ser Arg Pro Gln Ile Ile Lys Glu Val Leu Ala Val Pro Asn Ser Ile  
 530 535 540

Leu Glu Leu Pro Cys Pro His Leu Ser Ala Leu Ala Ser Tyr Tyr Trp  
 545 550 555 560

Ser His Gly Pro Ala Ala Val Pro Glu Ala Ser Ser Thr Val Tyr Asn  
 565 570 575

Gly Ser Leu Leu Leu Ile Val Gln Asp Gly Val Gly Gly Leu Tyr Gln  
 580 585 590

Cys Trp Ala Thr Glu Asn Gly Phe Ser Tyr Pro Val Ile Ser Tyr Trp  
 595 600 605

Val Asp Ser Gln Asp Gln Thr Leu Ala Leu Asp Pro Glu Leu Ala Gly  
 610 615 620

Ile Pro Arg Glu His Val Lys Val Pro Leu Thr Arg Val Ser Gly Gly  
 625 630 635 640

Ala Ala Leu Ala Ala Gln Gln Ser Tyr Trp Pro His Phe Val Thr Val  
 645 650 655

Thr Val Leu Phe Ala Leu Val Leu Ser Gly Ala Leu Ile Ile Leu Val  
660 665 670

Ala Ser Pro Leu Arg Ala Leu Arg Ala Arg Gly Lys Val Gln Gly Cys  
675 680 685

Glu Thr Leu Arg Pro Gly Glu Lys Ala Pro Leu Ser Arg Glu Gln His  
690 695 700

Leu Gln Ser Pro Lys Glu Cys Arg Thr Ser Ala Ser Asp Val Asp Ala  
705 710 715 720

Asp Asn Asn Cys Leu Gly Thr Glu Val Ala  
725 730

<210> 22  
<211> 652  
<212> PRT  
<213> Homo sapiens

<400> 22  
Gly Gly Gly Gly Gln Gly Pro Met Pro Arg Val Arg Tyr Tyr Ala Gly  
1 5 10 15

Asp Glu Arg Arg Ala Leu Ser Phe Phe His Gln Lys Gly Leu Gln Asp  
20 25 30

Phe Asp Thr Leu Leu Leu Ser Gly Asp Gly Asn Thr Leu Tyr Val Gly  
35 40 45

Ala Arg Glu Ala Ile Leu Ala Leu Asp Ile Gln Asp Pro Gly Val Pro  
50 55 60

Arg Leu Lys Asn Met Ile Pro Trp Pro Ala Ser Asp Arg Lys Lys Ser  
65 70 75 80

Glu Cys Ala Phe Lys Lys Lys Ser Asn Glu Thr Gln Cys Phe Asn Phe  
85 90 95

Ile Arg Val Leu Val Ser Tyr Asn Val Thr His Leu Tyr Thr Cys Gly  
100 105 110

Thr Phe Ala Phe Ser Pro Ala Cys Thr Phe Ile Glu Leu Gln Asp Ser  
115 120 125

Tyr Leu Leu Pro Ile Ser Glu Asp Lys Val Met Glu Gly Lys Gly Gln  
130 135 140

Ser Pro Phe Asp Pro Ala His Lys His Thr Ala Val Leu Val Asp Gly  
 145 150 155 160

Met Leu Tyr Ser Gly Thr Met Asn Asn Phe Leu Gly Ser Glu Pro Ile  
 165 170 175

Leu Met Arg Thr Leu Gly Ser Gln Pro Val Leu Lys Thr Asp Asn Phe  
 180 185 190

Leu Arg Trp Leu His His Asp Ala Ser Phe Val Ala Ala Ile Pro Ser  
 195 200 205

Thr Gln Val Val Tyr Phe Phe Phe Glu Glu Thr Ala Ser Glu Phe Asp  
 210 215 220

Phe Phe Glu Arg Leu His Thr Ser Arg Val Ala Arg Val Cys Lys Asn  
 225 230 235 240

Asp Val Gly Gly Glu Lys Leu Leu Gln Lys Lys Trp Thr Thr Phe Leu  
 245 250 255

Lys Ala Gln Leu Leu Cys Thr Gln Pro Gly Gln Leu Pro Phe Asn Val  
 260 265 270

Ile Arg His Ala Val Leu Leu Pro Ala Asp Ser Pro Thr Ala Pro His  
 275 280 285

Ile Tyr Ala Val Phe Thr Ser Gln Trp Gln Val Gly Gly Thr Arg Ser  
 290 295 300

Ser Ala Val Cys Ala Phe Ser Leu Leu Asp Ile Glu Arg Val Phe Lys  
 305 310 315 320

Gly Lys Tyr Lys Glu Leu Asn Lys Glu Thr Ser Arg Trp Thr Thr Tyr  
 325 330 335

Arg Gly Pro Glu Thr Asn Pro Arg Pro Gly Ser Cys Ser Val Gly Pro  
 340 345 350

Ser Ser Asp Lys Ala Leu Thr Phe Met Lys Asp His Phe Leu Met Asp  
 355 360 365

Glu Gln Val Val Gly Thr Pro Leu Leu Val Lys Ser Gly Val Glu Tyr  
 370 375 380

Thr Arg Leu Ala Val Glu Thr Ala Gln Gly Leu Asp Gly His Ser His  
 385 390 395 400

Leu Val Met Tyr Leu Gly Thr Thr Thr Gly Ser Leu His Lys Ala Val  
 405 410 415  
 Val Ser Gly Asp Ser Ser Ala His Leu Val Glu Glu Ile Gln Leu Phe  
 420 425 430  
 Pro Asp Pro Glu Pro Val Arg Asn Leu Gln Leu Ala Pro Thr Gln Gly  
 435 440 445  
 Ala Val Phe Val Gly Phe Ser Gly Gly Val Trp Arg Val Pro Arg Ala  
 450 455 460  
 Asn Cys Ser Val Tyr Glu Ser Cys Val Asp Cys Val Leu Ala Arg Asp  
 465 470 475 480  
 Pro His Cys Ala Trp Asp Pro Glu Ser Arg Thr Cys Cys Leu Leu Ser  
 485 490 495  
 Ala Pro Asn Leu Asn Ser Trp Lys Gln Asp Met Glu Arg Gly Asn Pro  
 500 505 510  
 Glu Trp Ala Cys Ala Ser Gly Pro Met Ser Arg Ser Leu Arg Pro Gln  
 515 520 525  
 Ser Arg Pro Gln Ile Ile Lys Glu Val Leu Ala Val Pro Asn Ser Ile  
 530 535 540  
 Leu Glu Leu Pro Cys Pro His Leu Ser Ala Leu Ala Ser Tyr Tyr Trp  
 545 550 555 560  
 Ser His Gly Pro Ala Ala Val Pro Glu Ala Ser Ser Thr Val Tyr Asn  
 565 570 575  
 Gly Ser Leu Leu Leu Ile Val Gln Asp Gly Val Gly Gly Leu Tyr Gln  
 580 585 590  
 Cys Trp Ala Thr Glu Asn Gly Phe Ser Tyr Pro Val Ile Ser Tyr Trp  
 595 600 605  
 Val Asp Ser Gln Asp Gln Thr Leu Ala Leu Asp Pro Glu Leu Ala Gly  
 610 615 620  
 Ile Pro Arg Glu His Val Lys Val Pro Leu Thr Arg Val Ser Gly Gly  
 625 630 635 640  
 Ala Ala Leu Ala Ala Gln Gln Ser Tyr Trp Pro His  
 645 650

<210> 23  
 <211> 21  
 <212> PRT  
 <213> Homo sapiens

<400> 23  
 Phe Val Thr Val Thr Val Leu Phe Ala Leu Val Leu Ser Gly Ala Leu  
 1 5 10 15  
 Ile Ile Leu Val Ala  
 20

<210> 24  
 <211> 57  
 <212> PRT  
 <213> Homo sapiens

<400> 24  
 Ser Pro Leu Arg Ala Leu Arg Ala Arg Gly Lys Val Gln Gly Cys Glu  
 1 5 10 15  
 Thr Leu Arg Pro Gly Glu Lys Ala Pro Leu Ser Arg Glu Gln His Leu  
 20 25 30  
 Gln Ser Pro Lys Glu Cys Arg Thr Ser Ala Ser Asp Val Asp Ala Asp  
 35 40 45  
 Asn Asn Cys Leu Gly Thr Glu Val Ala  
 50 55

<210> 25  
 <211> 2964  
 <212> DNA  
 <213> Homo sapiens

<400> 25  
 gtcgacccac gcgtccgcgg acgcgtgggg acggctcccg gctgcagtct gcccgcccgc 60  
 cccgcgcggg ggccgagtcg cgaagcgcgc ctgcgacccg gcgtccgggc gcgtggaga 120  
 ggacgcgagg agccatgagg cgcagcctg cgaaggtggc ggcgctgctg ctccggctgc 180  
 tcttgaggatg cacagaagcc aaaaagcatt gctggtattt cgaaggactc tatccaacct 240  
 attatatatg ccgctcctac gaggactgct gtggctccag gtgctgtgtg cgggccctct 300  
 ccatacagag gctgtggtac ttctggttcc ttctgatgat gggcgtgctt ttctgctgcg 360  
 gagccggctt cttcatccgg aggcgcattg acccccgcgc gctgatcgag gagccagcct 420  
 tcaatgtgtc ctacaccagg cagcccccaa atcccggccc aggagcccag cagccggggc 480



<400> 26

atgaggcgcc agcctgcgaa ggtggcgggc ctgctgctcg ggctgctctt ggagtgcaca 60  
gaagccaaaa agcattgctg gtatttcgaa ggactctatc caacctatta tatatgcgc 120  
tctacgagg actgctgtgg ctccagggtc tgtgtgcggg ccctctccat acagaggctg 180  
tggtacttct ggttccttct gatgatgggc gtgcttttct gctgcggagc cggcttcttc 240  
atccggaggc gcatgtaccc cccgccgctg atcgaggagc cagccttcaa tgtgtcctac 300  
accaggcagc ccccaaatcc cggcccagga gcccagcagc cggggccgcc ctattacact 360  
gacccaggag gaccggggat gaaccctgtc ggaattcca tggcaatggc tttccaggtc 420  
ccaccaact caccocaggg gagtgtggcc tgcccgcccc ctccagccta ctgcaacacg 480  
cctccgcccc cgtacgaaca ggtagtgaag gccaag 516

<210> 27

<211> 172

<212> PRT

<213> Homo sapiens

<400> 27

Met Arg Arg Gln Pro Ala Lys Val Ala Ala Leu Leu Leu Gly Leu Leu  
1 5 10 15

Leu Glu Cys Thr Glu Ala Lys Lys His Cys Trp Tyr Phe Glu Gly Leu  
20 25 30

Tyr Pro Thr Tyr Tyr Ile Cys Arg Ser Tyr Glu Asp Cys Cys Gly Ser  
35 40 45

Arg Cys Cys Val Arg Ala Leu Ser Ile Gln Arg Leu Trp Tyr Phe Trp  
50 55 60

Phe Leu Leu Met Met Gly Val Leu Phe Cys Cys Gly Ala Gly Phe Phe  
65 70 75 80

Ile Arg Arg Arg Met Tyr Pro Pro Pro Leu Ile Glu Glu Pro Ala Phe  
85 90 95

Asn Val Ser Tyr Thr Arg Gln Pro Pro Asn Pro Gly Pro Gly Ala Gln  
100 105 110

Gln Pro Gly Pro Pro Tyr Tyr Thr Asp Pro Gly Gly Pro Gly Met Asn  
115 120 125

Pro Val Gly Asn Ser Met Ala Met Ala Phe Gln Val Pro Pro Asn Ser  
130 135 140

Pro Gln Gly Ser Val Ala Cys Pro Pro Pro Pro Ala Tyr Cys Asn Thr  
145 150 155 160

Pro Pro Pro Pro Tyr Glu Gln Val Val Lys Ala Lys

165

170

<210> 28  
 <211> 22  
 <212> PRT  
 <213> Homo sapiens

<400> 28  
 Met Arg Arg Gln Pro Ala Lys Val Ala Ala Leu Leu Leu Gly Leu Leu  
           1                  5                  10                  15  
 Leu Glu Cys Thr Glu Ala  
                   20

<210> 29  
 <211> 150  
 <212> PRT  
 <213> Homo sapiens

<400> 29  
 Lys Lys His Cys Trp Tyr Phe Glu Gly Leu Tyr Pro Thr Tyr Tyr Ile  
           1                  5                  10                  15  
 Cys Arg Ser Tyr Glu Asp Cys Cys Gly Ser Arg Cys Cys Val Arg Ala  
                   20                  25                  30  
 Leu Ser Ile Gln Arg Leu Trp Tyr Phe Trp Phe Leu Leu Met Met Gly  
           35                  40                  45  
 Val Leu Phe Cys Cys Gly Ala Gly Phe Phe Ile Arg Arg Arg Met Tyr  
           50                  55                  60  
 Pro Pro Pro Leu Ile Glu Glu Pro Ala Phe Asn Val Ser Tyr Thr Arg  
           65                  70                  75                  80  
 Gln Pro Pro Asn Pro Gly Pro Gly Ala Gln Gln Pro Gly Pro Pro Tyr  
                   85                  90                  95  
 Tyr Thr Asp Pro Gly Gly Pro Gly Met Asn Pro Val Gly Asn Ser Met  
           100                  105                  110  
 Ala Met Ala Phe Gln Val Pro Pro Asn Ser Pro Gln Gly Ser Val Ala  
           115                  120                  125  
 Cys Pro Pro Pro Pro Ala Tyr Cys Asn Thr Pro Pro Pro Pro Tyr Glu  
           130                  135                  140

Gln Val Val Lys Ala Lys  
145 150

<210> 30  
<211> 38  
<212> PRT  
<213> Homo sapiens

<400> 30  
Lys Lys His Cys Trp Tyr Phe Glu Gly Leu Tyr Pro Thr Tyr Tyr Ile  
1 5 10 15

Cys Arg Ser Tyr Glu Asp Cys Cys Gly Ser Arg Cys Cys Val Arg Ala  
20 25 30

Leu Ser Ile Gln Arg Leu  
35

<210> 31  
<211> 21  
<212> PRT  
<213> Homo sapiens

<400> 31  
Trp Tyr Phe Trp Phe Leu Leu Met Met Gly Val Leu Phe Cys Cys Gly  
1 5 10 15

Ala Gly Phe Phe Ile  
20

<210> 32  
<211> 91  
<212> PRT  
<213> Homo sapiens

<400> 32  
Arg Arg Arg Met Tyr Pro Pro Pro Leu Ile Glu Glu Pro Ala Phe Asn  
1 5 10 15

Val Ser Tyr Thr Arg Gln Pro Pro Asn Pro Gly Pro Gly Ala Gln Gln  
20 25 30

Pro Gly Pro Pro Tyr Tyr Thr Asp Pro Gly Gly Pro Gly Met Asn Pro  
35 40 45

Val Gly Asn Ser Met Ala Met Ala Phe Gln Val Pro Pro Asn Ser Pro  
 50 55 60

Gln Gly Ser Val Ala Cys Pro Pro Pro Pro Ala Tyr Cys Asn Thr Pro  
 65 70 75 80

Pro Pro Pro Tyr Glu Gln Val Val Lys Ala Lys  
 85 90

<210> 33  
 <211> 1980  
 <212> DNA  
 <213> Homo sapiens

<400> 33  
 gtgcacccac gcgtccgcag ctttggacac ttcctctgct tgaggacacc ttgactaacc 60  
 tccaagggca actaaaggat caagaaaggc ccagcacagc agaagatcag ctggatctag 120  
 ctctgcagg agatgtgtac aaagacaatc ccagtcctct ggggatgttt cctcctgtgg 180  
 aatctctatg tctcatcctc tcagaccatt taccctggaa tcaaggcaag gattactcag 240  
 agggcacttg actatggtgt tcaagctgga atgaagatga ttgagcaaata gctaaaagaa 300  
 aagaaactcc cagattttaag cggttctgag tctcttgaat ttctaaaagt tgattatgta 360  
 aactacaatt tttaaataat aaaaatcagt gccttttcat ttccaaatac ctcatgggt 420  
 tttgtgcctg gagtgggaat caaagcgcta accaaccatg gcaactgcaa catcagcaca 480  
 gactgggggt tcgagtctcc actttttgtt ctgtataact cctttgctga gcccatggag 540  
 aaacccattt taaagaactt aaatgaaatg ctctgtccca ttattgcaag tgaagtcaaa 600  
 gcgctaaatg ccaacctcag cactactggag gttttaacca agattgacaa ctacactctg 660  
 ctggattact ccctaatacag ttctccagaa attactgaga actacottga cctgaacttg 720  
 aaggggtgat tctaccactt ggaaaacctc accgacccc ccttctcacc agttcctttt 780  
 gtgtccccag aacgcagcaa ctccatgctc tacattggaa tcgccgagta tttcttttaa 840  
 tctgcgtcct ttgtcattt cacagctggg gttttcaatc tactctctc caccgaagag 900  
 atttccaacc attttgttca aaactctcaa ggccttggca acgtgtctc ccggattgca 960  
 gagatctaca tcttgtccca gcccttcatg gtgaggatca tggccacaga gcctcccata 1020  
 atcaatctac aaccaggcaa ttccacctg gacatccctg cctccatcat gatgtcacc 1080  
 caacccaaga actccacagt tgaaaccatc gtttccatgg acttcgttgc tagtaccagt 1140  
 gttggcctgg ttattttggg acaaagactg gtctgtcctc tgtctctgaa cagattccgc 1200  
 cttgctttgc cagagtccaa tcgcagcaac attgaggtct tgaggtttga aaatattcta 1260  
 tcgtccattc ttcacttttg agtctccca ctggccaatg caaaattgca gcaaggattt 1320  
 cctctgcca atccacacaa attcttattc gtcaattcag atattgaagt tcttgagggg 1380  
 ttctttttga tttccaccga cctgaagtat gaaacatcct caaagcagca gccaaagttt 1440  
 cacgtatggg aaggtctgaa cctgataagc agacagtgga gggggaagtc agccccttga 1500  
 ttgccggttt gcaattcacc ccaggaagta aatggctcct aatcctacaa ctactgtaaa 1560  
 ccagaaagg aaagacagta cactactggaa ttgtaaagcc cttgtgaatt gcttaggcag 1620  
 aaagttttct ttcttaagcc ttcaggaacc cagaataagg cagactctgt taaagggata 1680  
 aatagaggtg tctgaatgtg agtgtatgca tgctgcgtgt gtctgtgttt atgtttgttt 1740  
 gtttgtttgg ggcaagaaag attctaggac aagagctagg catgtacttc tgaccaggtg 1800  
 ggtaagcaac tctaagtctg tatttgtatt ggtcattctc agtggaaatc ccttaggccc 1860

tctagtgggtt ttccccctacc tgcattattgg ttttcatggtt ttatattcac tggtactatc 1920  
 ttctgtgtttt aattaaaatt gttttctatc aaaaaaaaaa aaaaaaaaaa gggcggccgc 1980

<210> 34  
 <211> 1365  
 <212> DNA  
 <213> Homo sapiens

<400> 34  
 atgtgtacaa agacaatccc agtcctctgg ggatgtttcc tcctgtggaa tctctatgtc 60  
 tcacccctctc agaccattta ccctggaatc aaggcaagga ttactcagag ggcacttgac 120  
 tatggtgttc aagctggaat gaagatgatt gagcaaagtc taaaagaaaa gaaactccca 180  
 gatttaagcg gttctgagtc tcttgaattt ctaaaagtgt attatgtaaa ctacaatttt 240  
 tcaaatataa aaatcagtg cttttcattt ccaaatacct cattggcttt tgtgcctgga 300  
 gtgggaatca aagcgctaac caaccatggc actgccaaca tcagcacaga ctggggggttc 360  
 gagtctccac tttttgttct gtataactcc tttgctgagc ccatggagaa acccatttta 420  
 aagaacttaa atgaaatgct ctgtcccatt attgcaagtg aagtcaaagc gctaaatgcc 480  
 aacctcagca cactggaggt tttaaccaag attgacaact acactctgct ggattactcc 540  
 ctaatcagtt ctccagaaat tactgagaac taccttgacc tgaacttgaa ggggtgtattc 600  
 taccactggg aaaacctcac cgaccccccc ttctcaccag ttctttttgt gctcccagaa 660  
 cgcagcaact ccatgctcta cattggaatc gccgagtatt tctttaaatc tgcgtccttt 720  
 gctcatttca cagctgggggt tttcaatctc actctctcca ccgaagagat ttccaacccat 780  
 tttgttcaaa actctcaagg ccttggaac gtgctctccc ggattgcaga gatctacatc 840  
 ttgtcccagc ccttcatggt gaggatcatg gccacagagc ctcccataat caatctacaa 900  
 ccaggcaatt tcaccctgga catccctgcc tccatcatga tgctcaccca acccaagaac 960  
 tccacagttg aaaccatcgt ttccatggac ttctgtgcta gtaccagtgt tggcctgggt 1020  
 attttgggac aaagactggt ctgctccttg tctctgaaca gattccgcct tgctttgcca 1080  
 gagtccaatc gcagcaacat tgaggctctg aggtttgaaa atattctatc gtccattctt 1140  
 cacttttgag tcctccact ggccaatgca aaattgcagc aaggatttcc tctgccaat 1200  
 ccacacaaat tcttattcgt caattcagat attgaagttc ttgaggggtt ccttttgatt 1260  
 tccaccgacc tgaagtatga aacatcctca aagcagcagc caagtttcca cgtatgggaa 1320  
 ggtctgaacc tgataagcag acagtggagg gggaagtcag cccct 1365

<210> 35  
 <211> 455  
 <212> PRT  
 <213> Homo sapiens

<400> 35  
 Met Cys Thr Lys Thr Ile Pro Val Leu Trp Gly Cys Phe Leu Leu Trp  
 1 5 10 15  
 Asn Leu Tyr Val Ser Ser Ser Gln Thr Ile Tyr Pro Gly Ile Lys Ala  
 20 25 30  
 Arg Ile Thr Gln Arg Ala Leu Asp Tyr Gly Val Gln Ala Gly Met Lys  
 35 40 45

Met Ile Glu Gln Met Leu Lys Glu Lys Lys Leu Pro Asp Leu Ser Gly  
 50 55 60

Ser Glu Ser Leu Glu Phe Leu Lys Val Asp Tyr Val Asn Tyr Asn Phe  
 65 70 75 80

Ser Asn Ile Lys Ile Ser Ala Phe Ser Phe Pro Asn Thr Ser Leu Ala  
 85 90 95

Phe Val Pro Gly Val Gly Ile Lys Ala Leu Thr Asn His Gly Thr Ala  
 100 105 110

Asn Ile Ser Thr Asp Trp Gly Phe Glu Ser Pro Leu Phe Val Leu Tyr  
 115 120 125

Asn Ser Phe Ala Glu Pro Met Glu Lys Pro Ile Leu Lys Asn Leu Asn  
 130 135 140

Glu Met Leu Cys Pro Ile Ile Ala Ser Glu Val Lys Ala Leu Asn Ala  
 145 150 155 160

Asn Leu Ser Thr Leu Glu Val Leu Thr Lys Ile Asp Asn Tyr Thr Leu  
 165 170 175

Leu Asp Tyr Ser Leu Ile Ser Ser Pro Glu Ile Thr Glu Asn Tyr Leu  
 180 185 190

Asp Leu Asn Leu Lys Gly Val Phe Tyr Pro Leu Glu Asn Leu Thr Asp  
 195 200 205

Pro Pro Phe Ser Pro Val Pro Phe Val Leu Pro Glu Arg Ser Asn Ser  
 210 215 220

Met Leu Tyr Ile Gly Ile Ala Glu Tyr Phe Phe Lys Ser Ala Ser Phe  
 225 230 235 240

Ala His Phe Thr Ala Gly Val Phe Asn Leu Thr Leu Ser Thr Glu Glu  
 245 250 255

Ile Ser Asn His Phe Val Gln Asn Ser Gln Gly Leu Gly Asn Val Leu  
 260 265 270

Ser Arg Ile Ala Glu Ile Tyr Ile Leu Ser Gln Pro Phe Met Val Arg  
 275 280 285

Ile Met Ala Thr Glu Pro Pro Ile Ile Asn Leu Gln Pro Gly Asn Phe  
 290 295 300

Thr Leu Asp Ile Pro Ala Ser Ile Met Met Leu Thr Gln Pro Lys Asn  
 305 310 315 320

Ser Thr Val Glu Thr Ile Val Ser Met Asp Phe Val Ala Ser Thr Ser  
 325 330 335

Val Gly Leu Val Ile Leu Gly Gln Arg Leu Val Cys Ser Leu Ser Leu  
 340 345 350

Asn Arg Phe Arg Leu Ala Leu Pro Glu Ser Asn Arg Ser Asn Ile Glu  
 355 360 365

Val Leu Arg Phe Glu Asn Ile Leu Ser Ser Ile Leu His Phe Gly Val  
 370 375 380

Leu Pro Leu Ala Asn Ala Lys Leu Gln Gln Gly Phe Pro Leu Pro Asn  
 385 390 395 400

Pro His Lys Phe Leu Phe Val Asn Ser Asp Ile Glu Val Leu Glu Gly  
 405 410 415

Phe Leu Leu Ile Ser Thr Asp Leu Lys Tyr Glu Thr Ser Ser Lys Gln  
 420 425 430

Gln Pro Ser Phe His Val Trp Glu Gly Leu Asn Leu Ile Ser Arg Gln  
 435 440 445

Trp Arg Gly Lys Ser Ala Pro  
 450 455

<210> 36

<211> 23

<212> PRT

<213> Homo sapiens

<400> 36

Met Cys Thr Lys Thr Ile Pro Val Leu Trp Gly Cys Phe Leu Leu Trp  
 1 5 10 15

Asn Leu Tyr Val Ser Ser Ser  
 20

<210> 37

<211> 432

<212> PRT

<213> Homo sapiens

<400> 37

Gln Thr Ile Tyr Pro Gly Ile Lys Ala Arg Ile Thr Gln Arg Ala Leu  
1 5 10 15

Asp Tyr Gly Val Gln Ala Gly Met Lys Met Ile Glu Gln Met Leu Lys  
20 25 30

Glu Lys Lys Leu Pro Asp Leu Ser Gly Ser Glu Ser Leu Glu Phe Leu  
35 40 45

Lys Val Asp Tyr Val Asn Tyr Asn Phe Ser Asn Ile Lys Ile Ser Ala  
50 55 60

Phe Ser Phe Pro Asn Thr Ser Leu Ala Phe Val Pro Gly Val Gly Ile  
65 70 75 80

Lys Ala Leu Thr Asn His Gly Thr Ala Asn Ile Ser Thr Asp Trp Gly  
85 90 95

Phe Glu Ser Pro Leu Phe Val Leu Tyr Asn Ser Phe Ala Glu Pro Met  
100 105 110

Glu Lys Pro Ile Leu Lys Asn Leu Asn Glu Met Leu Cys Pro Ile Ile  
115 120 125

Ala Ser Glu Val Lys Ala Leu Asn Ala Asn Leu Ser Thr Leu Glu Val  
130 135 140

Leu Thr Lys Ile Asp Asn Tyr Thr Leu Leu Asp Tyr Ser Leu Ile Ser  
145 150 155 160

Ser Pro Glu Ile Thr Glu Asn Tyr Leu Asp Leu Asn Leu Lys Gly Val  
165 170 175

Phe Tyr Pro Leu Glu Asn Leu Thr Asp Pro Pro Phe Ser Pro Val Pro  
180 185 190

Phe Val Leu Pro Glu Arg Ser Asn Ser Met Leu Tyr Ile Gly Ile Ala  
195 200 205

Glu Tyr Phe Phe Lys Ser Ala Ser Phe Ala His Phe Thr Ala Gly Val  
210 215 220

Phe Asn Leu Thr Leu Ser Thr Glu Glu Ile Ser Asn His Phe Val Gln  
225 230 235 240

Asn Ser Gln Gly Leu Gly Asn Val Leu Ser Arg Ile Ala Glu Ile Tyr



Leu Val Ala Ile Gly Thr Ala Val Thr Ala Ala Val Asn Pro Gly Val  
 20 25 30

Val Val Arg Ile Ser Gln Lys Gly Leu Asp Tyr Ala Ser Gln Gln Gly  
 35 40 45

Thr Ala Ala Leu Gln Lys Glu Leu Lys Arg Ile Lys Ile Pro Asp Tyr  
 50 55 60

Ser Asp Ser Phe Lys Ile Lys His Leu Gly Lys Gly His Tyr Ser Phe  
 65 70 75 80

Tyr Ser Met Asp Ile Arg Glu Phe Gln Leu Pro Ser Ser Gln Ile Ser  
 85 90 95

Met Val Pro Asn Val Gly Leu Lys Phe Ser Ile Ser Asn Ala Asn Ile  
 100 105 110

Lys Ile Ser Gly Lys Trp Lys Ala Gln Lys Arg Phe Leu Lys Met Ser  
 115 120 125

Gly Asn Phe Asp Leu Ser Ile Glu Gly Met Ser Ile Ser Ala Asp Leu  
 130 135 140

Lys Leu Gly Ser Asn Pro Thr Ser Gly Lys Pro Thr Ile Thr Cys Ser  
 145 150 155 160

Ser Cys Ser Ser His Ile Asn Ser Val His Val His Ile Ser Lys Ser  
 165 170 175

Lys Val Gly Trp Leu Ile Gln Leu Phe His Lys Lys Ile Glu Ser Ala  
 180 185 190

Leu Arg Asn Lys Met Asn Ser Gln Val Cys Glu Lys Val Thr Asn Ser  
 195 200 205

Val Ser Ser Lys Leu Gln Pro Tyr Phe Gln Thr Leu Pro Val Met Thr  
 210 215 220

Lys Ile Asp Ser Val Ala Gly Ile Asn Tyr Gly Leu Val Ala Pro Pro  
 225 230 235 240

Ala Thr Thr Ala Glu Thr Leu Asp Val Gln Met Lys Gly Glu Phe Tyr  
 245 250 255

Ser Glu Asn His His Asn Pro Pro Pro Phe Ala Pro Pro Val Met Glu  
 260 265 270

Phe Pro Ala Ala His Asp Arg Met Val Tyr Leu Gly Leu Ser Asp Tyr  
 275 280 285

Phe Phe Asn Thr Ala Gly Leu Val Tyr Gln Glu Ala Gly Val Leu Lys  
 290 295 300

Met Thr Leu Arg Asp Asp Met Ile Pro Lys Glu Ser Lys Phe Arg Leu  
 305 310 315 320

Thr Thr Lys Phe Phe Gly Thr Phe Leu Pro Glu Val Ala Lys Lys Phe  
 325 330 335

Pro Asn Met Lys Ile Gln Ile His Val Ser Ala Ser Thr Pro Pro His  
 340 345 350

Leu Ser Val Gln Pro Thr Gly Leu Thr Phe Tyr Pro Ala Val Asp Val  
 355 360 365

Gln Ala Phe Ala Val Leu Pro Asn Ser Ser Leu Ala Ser Leu Phe Leu  
 370 375 380

Ile Gly Met His Thr Thr Gly Ser Met Glu Val Ser Ala Glu Ser Asn  
 385 390 395 400

Arg Leu Val Gly Glu Leu Lys Leu Asp Arg Leu Leu Leu Glu Leu Lys  
 405 410 415

His Ser Asn Ile Gly Pro Phe Pro Val Glu Leu Leu Gln Asp Ile Met  
 420 425 430

Asn Tyr Ile Val Pro Ile Leu Val Leu Pro Arg Val Asn Glu Lys Leu  
 435 440 445

Gln Lys Gly Phe Pro Leu Pro Thr Pro Ala Arg Val Gln Leu Tyr Asn  
 450 455 460

Val Val Leu Gln Pro His Gln Asn Phe Leu Leu Phe Gly Ala Asp Val  
 465 470 475 480

Val Tyr Lys

<210> 39

<211> 481

<212> PRT

<213> Homo sapiens

<400> 39

Met Gly Ala Leu Ala Arg Ala Leu Pro Ser Ile Leu Leu Ala Leu Leu  
1 5 10 15

Leu Thr Ser Thr Pro Glu Ala Leu Gly Ala Asn Pro Gly Leu Val Ala  
20 25 30

Arg Ile Thr Asp Lys Gly Leu Gln Tyr Ala Ala Gln Glu Gly Leu Leu  
35 40 45

Ala Leu Gln Ser Glu Leu Leu Arg Ile Thr Leu Pro Asp Phe Thr Gly  
50 55 60

Asp Leu Arg Ile Pro His Val Gly Arg Gly Arg Tyr Glu Phe His Ser  
65 70 75 80

Leu Asn Ile His Glu Phe Gln Leu Pro Ser Ser Gln Ile Ser Met Val  
85 90 95

Pro Asn Val Gly Leu Lys Phe Ser Ile Ser Asn Ala Asn Ile Lys Ile  
100 105 110

Ser Gly Lys Trp Lys Ala Gln Lys Arg Phe Leu Lys Met Ser Gly Asn  
115 120 125

Phe Asp Leu Ser Ile Glu Gly Met Ser Ile Ser Ala Asp Leu Lys Leu  
130 135 140

Gly Ser Asn Pro Thr Ser Gly Lys Pro Thr Ile Thr Cys Ser Ser Cys  
145 150 155 160

Ser Ser His Ile Asn Ser Val His Val His Ile Ser Lys Ser Lys Val  
165 170 175

Gly Trp Leu Ile Gln Leu Phe His Lys Lys Ile Glu Ser Ala Leu Arg  
180 185 190

Asn Lys Met Asn Ser Gln Val Cys Glu Lys Val Thr Asn Ser Val Ser  
195 200 205

Ser Lys Leu Gln Pro Tyr Phe Gln Thr Leu Pro Val Met Thr Lys Ile  
210 215 220

Asp Ser Val Ala Gly Ile Asn Tyr Gly Leu Val Ala Pro Pro Ala Thr  
225 230 235 240

Thr Ala Glu Thr Leu Asp Val Gln Met Lys Gly Glu Phe Tyr Ser Glu  
245 250 255

Asn His His Asn Pro Pro Pro Phe Ala Pro Pro Val Met Glu Phe Pro  
 260 265 270

Ala Ala His Asp Arg Met Val Tyr Leu Gly Leu Ser Asp Tyr Phe Phe  
 275 280 285

Asn Thr Ala Gly Leu Val Tyr Gln Glu Ala Gly Val Leu Lys Met Thr  
 290 295 300

Leu Arg Asp Asp Met Ile Pro Lys Glu Ser Lys Phe Arg Leu Thr Thr  
 305 310 315 320

Lys Phe Phe Gly Thr Phe Leu Pro Glu Val Ala Lys Lys Phe Pro Asn  
 325 330 335

Met Lys Ile Gln Ile His Val Ser Ala Ser Thr Pro Pro His Leu Ser  
 340 345 350

Val Gln Pro Thr Gly Leu Thr Phe Tyr Pro Ala Val Asp Val Gln Ala  
 355 360 365

Leu Ala Val Leu Pro Asn Ser Ser Leu Ala Ser Leu Phe Leu Ile Gly  
 370 375 380

Met His Thr Thr Gly Ser Met Glu Val Ser Ala Glu Ser Asn Arg Leu  
 385 390 395 400

Val Gly Glu Leu Lys Leu Asp Arg Leu Leu Leu Glu Leu Lys His Ser  
 405 410 415

Asn Ile Gly Pro Phe Pro Val Glu Leu Leu Gln Asp Ile Met Asn Tyr  
 420 425 430

Ile Val Pro Ile Leu Val Leu Pro Arg Val Asn Glu Lys Leu Gln Lys  
 435 440 445

Gly Phe Pro Leu Pro Thr Pro Ala Arg Val Gln Leu Tyr Asn Val Val  
 450 455 460

Leu Gln Pro His Gln Asn Phe Leu Leu Phe Gly Ala Asp Val Val Tyr  
 465 470 475 480

Lys

<210> 40

<211> 383

<212> PRT

<213> *Caenorhabditis elegans*

<400> 40

Met Arg Ile Ala His Ala Ser Ser Arg Gly Asn Ile Ser Ile Phe Ser  
1 5 10 15

Val Phe Leu Ile Pro Leu Ile Ala Tyr Ile Leu Ile Leu Pro Gly Val  
20 25 30

Arg Arg Lys Arg Val Val Thr Thr Val Thr Tyr Val Leu Met Leu Ala  
35 40 45

Val Gly Gly Ala Leu Ile Ala Ser Leu Ile Tyr Pro Cys Trp Ala Ser  
50 55 60

Gly Ser Gln Met Ile Tyr Thr Gln Phe Arg Gly His Ser Asn Glu Arg  
65 70 75 80

Ile Leu Ala Lys Ile Gly Val Glu Ile Gly Leu Gln Lys Val Asn Val  
85 90 95

Thr Leu Lys Phe Glu Arg Leu Leu Ser Ser Asn Asp Val Leu Pro Gly  
100 105 110

Ser Asp Met Thr Glu Leu Tyr Tyr Asn Glu Gly Phe Asp Ile Ser Gly  
115 120 125

Ile Ser Ser Met Ala Glu Ala Leu His His Gly Leu Glu Asn Gly Leu  
130 135 140

Pro Tyr Pro Met Leu Ser Val Leu Glu Tyr Phe Ser Leu Asn Gln Asp  
145 150 155 160

Ser Phe Asp Trp Gly Arg His Tyr Arg Val Ala Gly His Tyr Thr His  
165 170 175

Ala Ala Ile Trp Phe Ala Phe Ala Cys Trp Cys Leu Ser Val Val Leu  
180 185 190

Met Leu Phe Leu Pro His Asn Ala Tyr Lys Ser Ile Leu Ala Thr Gly  
195 200 205

Ile Ser Cys Leu Ile Ala Cys Leu Val Tyr Leu Leu Leu Ser Pro Cys  
210 215 220

Glu Leu Arg Ile Ala Phe Thr Gly Glu Asn Phe Glu Arg Val Asp Leu

225                      230                      235                      240  
 Thr Ala Thr Phe Ser Phe Cys Phe Tyr Leu Ile Phe Ala Ile Gly Ile  
                                  245                                   250                                   255  
 Leu Cys Val Leu Cys Gly Leu Gly Leu Gly Ile Cys Glu His Trp Arg  
                                  260                                   265                                   270  
 Ile Tyr Thr Leu Ser Thr Phe Leu Asp Ala Ser Leu Asp Glu His Val  
                                  275                                   280                                   285  
 Gly Pro Lys Trp Lys Lys Leu Pro Thr Gly Gly Pro Ala Leu Gln Gly  
                                  290                                   295                                   300  
 Val Gln Ile Gly Ala Tyr Gly Thr Asn Thr Thr Asn Ser Ser Arg Asp  
 305                                   310                                   315                                   320  
 Lys Asn Asp Ile Ser Ser Asp Lys Thr Ala Gly Ser Ser Gly Phe Gln  
                                  325                                   330                                   335  
 Ser Arg Thr Ser Thr Cys Gln Ser Ser Ala Ser Ser Ala Ser Leu Arg  
                                  340                                   345                                   350  
 Ser Gln Ser Ser Ile Glu Thr Val His Asp Glu Ala Glu Leu Glu Arg  
                                  355                                   360                                   365  
 Thr His Val His Phe Leu Gln Glu Pro Cys Ser Ser Ser Ser Thr  
                                  370                                   375                                   380

<210> 41  
 <211> 399  
 <212> PRT  
 <213> Homo sapiens

<400> 41  
 Met Lys Met Arg Phe Leu Gly Leu Val Val Cys Leu Val Leu Trp Pro  
   1                                   5                                   10                                   15  
 Leu His Ser Glu Gly Ser Gly Gly Lys Leu Thr Ala Val Asp Pro Glu  
                                  20                                   25                                   30  
 Thr Asn Met Asn Val Ser Glu Ile Ile Ser Tyr Trp Gly Phe Pro Ser  
                                  35                                   40                                   45  
 Glu Glu Tyr Leu Val Glu Thr Glu Asp Gly Tyr Ile Leu Cys Leu Asn  
                                  50                                   55                                   60



Asn Gln Ser Tyr Pro Pro Thr Tyr Asn Val Lys Asp Met Leu Val Pro  
 325 330 335

Thr Ala Val Trp Ser Gly Gly His Asp Trp Leu Ala Asp Val Tyr Asp  
 340 345 350

Val Asn Ile Leu Leu Thr Gln Ile Thr Asn Leu Val Phe His Glu Ser  
 355 360 365

Ile Pro Glu Trp Glu His Leu Asp Phe Ile Trp Gly Leu Asp Ala Pro  
 370 375 380

Trp Arg Leu Tyr Asn Lys Ile Ile Asn Leu Met Arg Lys Tyr Gln  
 385 390 395

<210> 42

<211> 19

<212> PRT

<213> Mus sp.

<400> 42

Met Ala Pro Pro Ala Ala Arg Leu Ala Leu Leu Ser Ala Ala Ala Leu  
 1 5 10 15

Thr Leu Ala

<210> 43

<211> 451

<212> PRT

<213> Mus sp.

<400> 43

Ala Arg Pro Ala Pro Gly Pro Arg Ser Gly Pro Glu Cys Phe Thr Ala  
 1 5 10 15

Asn Gly Ala Asp Tyr Arg Gly Thr Gln Ser Trp Thr Ala Leu Gln Gly  
 20 25 30

Gly Lys Pro Cys Leu Phe Trp Asn Glu Thr Phe Gln His Pro Tyr Asn  
 35 40 45

Thr Leu Lys Tyr Pro Asn Gly Glu Gly Gly Leu Gly Glu His Asn Tyr  
 50 55 60

Cys Arg Asn Pro Asp Gly Asp Val Ser Pro Trp Cys Tyr Val Ala Glu

65		70		75		80									
His	Glu	Asp	Gly	Val	Tyr	Trp	Lys	Tyr	Cys	Glu	Ile	Pro	Ala	Cys	Gln
			85						90					95	
Met	Pro	Gly	Asn	Leu	Gly	Cys	Tyr	Lys	Asp	His	Gly	Asn	Pro	Pro	Pro
			100					105					110		
Leu	Thr	Gly	Thr	Ser	Lys	Thr	Ser	Asn	Lys	Leu	Thr	Ile	Gln	Thr	Cys
		115					120					125			
Ile	Ser	Phe	Cys	Arg	Ser	Gln	Arg	Phe	Lys	Phe	Ala	Gly	Met	Glu	Ser
		130				135					140				
Gly	Tyr	Ala	Cys	Phe	Cys	Gly	Asn	Asn	Pro	Asp	Tyr	Trp	Lys	His	Gly
145					150					155					160
Glu	Ala	Ala	Ser	Thr	Glu	Cys	Asn	Ser	Val	Cys	Phe	Gly	Asp	His	Thr
				165					170					175	
Gln	Pro	Cys	Gly	Gly	Asp	Gly	Arg	Ile	Ile	Leu	Phe	Asp	Thr	Leu	Val
			180					185						190	
Gly	Ala	Cys	Gly	Gly	Asn	Tyr	Ser	Ala	Met	Ala	Ala	Val	Val	Tyr	Ser
		195					200					205			
Pro	Asp	Phe	Pro	Asp	Thr	Tyr	Ala	Thr	Gly	Arg	Val	Cys	Tyr	Trp	Thr
		210				215					220				
Ile	Arg	Val	Pro	Gly	Ala	Ser	Arg	Ile	His	Phe	Asn	Phe	Thr	Leu	Phe
225					230					235					240
Asp	Ile	Arg	Asp	Ser	Ala	Asp	Met	Val	Glu	Leu	Leu	Asp	Gly	Tyr	Thr
				245					250					255	
His	Arg	Val	Leu	Val	Arg	Leu	Ser	Gly	Arg	Ser	Arg	Pro	Pro	Leu	Ser
			260					265						270	
Phe	Asn	Val	Ser	Leu	Asp	Phe	Val	Ile	Leu	Tyr	Phe	Phe	Ser	Asp	Arg
		275					280					285			
Ile	Asn	Gln	Ala	Gln	Gly	Phe	Ala	Val	Leu	Tyr	Gln	Ala	Thr	Lys	Glu
		290				295						300			
Glu	Pro	Pro	Gln	Glu	Arg	Pro	Ala	Val	Asn	Gln	Thr	Leu	Ala	Glu	Val
305					310					315					320
Ile	Thr	Glu	Gln	Ala	Asn	Leu	Ser	Val	Ser	Ala	Ala	His	Ser	Ser	Lys



Gln Pro Pro Asn Pro Ala Pro Gly Ala Gln Gln Met Gly Pro Pro Tyr  
85 90 95

Tyr Thr Asp Pro Gly Gly Pro Gly Met Asn Pro Val Gly Asn Thr Met  
100 105 110

Ala Met Ala Phe Gln Val Gln Pro Asn Ser Pro His Gly Gly Thr Thr  
115 120 125

Tyr Pro Pro Pro Pro Ser Tyr Cys Asn Thr Pro Pro Pro Pro Tyr Glu  
130 135 140

Gln Val Val Lys Asp Lys  
145 150

<210> 45  
<211> 2044  
<212> DNA  
<213> Homo sapiens

<400> 45  
gtcgaccac gcgtccgggg aattgcagca ggaaaatatg tgaagagttt ttaaaccac 60  
aaattcttct tactttagaa ttagttgtta cattggcagg aaaaaataaa tgcagatgtt 120  
ggaccatgtt ggaaaccttg tcaagacagt ggattgtctc acacagaatg gaaatgtggc 180  
ttctgattct ggtggcgat atgttccaga gaaatgtgaa ttcagtacat atgccaacta 240  
aagctgtgga cccagaagca ttcatgaata ttagtgaaat catccaacat caaggctatc 300  
cctgtgagga atatgaagtc gcaactgaag atgggtatat ctttctgtt aacaggattc 360  
ctcgaggcct agtgcaacct aagaagacag gttccaggcc tgtggtgtta ctgcagcatg 420  
gcctagttag aggtgctagc aactggattt ccaacctgcc caacaatagc ctgggcttca 480  
ttctggcaga tgctggtttt gacgtgtgga tggggaacag caggggaaac gcctggtctc 540  
gaaaacacaa gacactctcc atagaccaag atgagttctg ggctttcagt tatgatgaga 600  
tggctaggtt tgaccttcc gacgtgataa actttatttt gcagaaaacg ggccaggaaa 660  
agatctatta tgcgggctat tcacagggca ccacctggg ctttattgca ttttccacca 720  
tgccagagct ggctcagaaa atcaaaatgt attttgcttt agcaccata gccactgtta 780  
agcatgcaaa aagccccggg accaaatttt tgttgctgcc agatatgatg atcaagggat 840  
tgtttgcaa aaaagaattt ctgtatcaga ccagatttct cagacaactt gttatttacc 900  
tttgtggcca ggtgattctt gafcagattt gtagtaatat catgttactt ctgggtggat 960  
tcaacaccaa caatatgaac atgagccgag caagtgtata tgctgcccac actcttgctg 1020  
gaacatctgt gcaaaatatt ctacactgga gccaggcagt gaattctggt gaactccggg 1080  
catttgactg ggggagttag accaaaaatc tggaaaaatg caatcagcca actcctgtaa 1140  
ggtacagagt cagagatatg acggtcccta cagcaatgtg gacaggaggt caggactggc 1200  
tttcaaatcc agaagacgtg aaaatgctgc tctctgaggt gaccaacctc atctaccata 1260  
agaatattcc tgaatgggct cacgtggatt tcatctggg tttggatgct cctcaccgta 1320  
tgtacaatga aatcatccat ctgatgcagc aggaggagac caacctttcc cagggacggg 1380  
gtgaggccgt attgtgaagc atctgacact gacgatctta ggacaacctc ctgagggatg 1440  
gggctaggac ccatgaaggc agaattacgg agagcagaga cctagtatac atttttcaga 1500  
ttccctgcac ttggcactaa atccgacact tacatttaca ttttttttct gtaaattaaa 1560

```

gtacttatta ggtaaataga ggttttgtat gctattatat attctacat cttgaagggt 1620
aggttttacc tgatagccag aaaatatcta gacattctct atatcattca ggtaaattctc 1680
tttaaaacac ctattgtttt ttctataagc catatTTTTg gagcactaaa gtaaaatggc 1740
aaattgggac agatattgag gtctggagtc tgtggattat tgttgacttt gacaaaataa 1800
gctagacatt ttcaccttgt tgccacagag acataacact acctcaggaa gctgagctgc 1860
tttaaggaca acaacaacaa aatcagtgtt acagtatgga tgaaatctat gttaagcatt 1920
ctcagaataa ggccaagttt tatagttgca tctcaggga gaaaatttta taggatgttt 1980
atgagttctc caataaatgc attctgcatt acataaaaaa aaaaaaaaaa aaaagggcgg 2040
ccgc 2044

```

<210> 46  
 <211> 1269  
 <212> DNA  
 <213> Homo sapiens

```

<400> 46
atgttggaac ccttgtcaag acagtggatt gtctcacaca gaatggaaat gtggcttctg 60
attctgggtg cgtatatgtt ccagagaaat gtgaattcag tacatatgcc aactaaagct 120
gtggaccag aagcattcat gaatattagt gaaatcatcc aacatcaagg ctatccctgt 180
gaggaatatg aagtcgcaac tgaagatggg tatatccttt ctgttaacag gattcctcga 240
ggcctagtgc aacctaagaa gacaggttcc aggcctgtgg tgttactgca gcatggccta 300
gttgagggtg ctagcaactg gatttccaac ctgcccacaa atagcctggg cttcattctg 360
gcagatgctg gttttgacgt gtggatgggg aacagcaggg gaaacgcctg gtctcgaaaa 420
cacaagacac tctccataga ccaagatgag ttctgggctt tcagttatga tgagatggct 480
aggtttgacc ttctctcagt gataaacttt attttgcaga aaacggggcca ggaaaagatc 540
tattatgtcg gctattcaca gggcaccacc atgggcttta ttgcattttc caccatgcc 600
gagctggctc agaaaatcaa aatgtatttt gctttagcac ccatagccac tgttaagcat 660
gcaaaaagcc ccgggaccaa atttttgttg ctgccagata tgatgatcaa gggattgttt 720
ggcaaaaaag aattttctgta tcagaccaga tttctcagac aacttggttat ttacctttgt 780
ggccaggtga ttcttgatca gattttagt aatatcatgt tacttctggg tggattcaac 840
accaacaata tgaacatgag ccgagcaagt gtatatgctg cccacactct tgctggaaca 900
tctgtgcaaa atattctaca ctggagccag gcagtgaatt ctggtgaact ccgggcattt 960
gactggggga gtgagaccaa aaatctggaa aaatgcaatc agccaactcc tgaagggtac 1020
agagtacagag atatgacggt ccctacagca atgtggacag gaggtcagga ctggctttca 1080
aatccagaag acgtgaaaat gctgctctct gaggtgacca acctcatcta coataagaat 1140
attcctgaat gggctcacgt ggatttcac tgggggttgg atgctcctca ccgtatgtac 1200
aatgaaatca tccatctgat gcagcaggag gagaccaacc tttcccaggg acggtgtgag 1260
gccgtattg 1269

```

<210> 47  
 <211> 423  
 <212> PRT  
 <213> Homo sapiens

```

<400> 47
Met Leu Glu Thr Leu Ser Arg Gln Trp Ile Val Ser His Arg Met Glu
1           5           10           15

```

Met Trp Leu Leu Ile Leu Val Ala Tyr Met Phe Gln Arg Asn Val Asn  
20 25 30

Ser Val His Met Pro Thr Lys Ala Val Asp Pro Glu Ala Phe Met Asn  
35 40 45

Ile Ser Glu Ile Ile Gln His Gln Gly Tyr Pro Cys Glu Glu Tyr Glu  
50 55 60

Val Ala Thr Glu Asp Gly Tyr Ile Leu Ser Val Asn Arg Ile Pro Arg  
65 70 75 80

Gly Leu Val Gln Pro Lys Lys Thr Gly Ser Arg Pro Val Val Leu Leu  
85 90 95

Gln His Gly Leu Val Gly Gly Ala Ser Asn Trp Ile Ser Asn Leu Pro  
100 105 110

Asn Asn Ser Leu Gly Phe Ile Leu Ala Asp Ala Gly Phe Asp Val Trp  
115 120 125

Met Gly Asn Ser Arg Gly Asn Ala Trp Ser Arg Lys His Lys Thr Leu  
130 135 140

Ser Ile Asp Gln Asp Glu Phe Trp Ala Phe Ser Tyr Asp Glu Met Ala  
145 150 155 160

Arg Phe Asp Leu Pro Ala Val Ile Asn Phe Ile Leu Gln Lys Thr Gly  
165 170 175

Gln Glu Lys Ile Tyr Tyr Val Gly Tyr Ser Gln Gly Thr Thr Met Gly  
180 185 190

Phe Ile Ala Phe Ser Thr Met Pro Glu Leu Ala Gln Lys Ile Lys Met  
195 200 205

Tyr Phe Ala Leu Ala Pro Ile Ala Thr Val Lys His Ala Lys Ser Pro  
210 215 220

Gly Thr Lys Phe Leu Leu Leu Pro Asp Met Met Ile Lys Gly Leu Phe  
225 230 235 240

Gly Lys Lys Glu Phe Leu Tyr Gln Thr Arg Phe Leu Arg Gln Leu Val  
245 250 255

Ile Tyr Leu Cys Gly Gln Val Ile Leu Asp Gln Ile Cys Ser Asn Ile  
260 265 270

Met Leu Leu Leu Gly Gly Phe Asn Thr Asn Asn Met Asn Met Ser Arg  
 275 280 285

Ala Ser Val Tyr Ala Ala His Thr Leu Ala Gly Thr Ser Val Gln Asn  
 290 295 300

Ile Leu His Trp Ser Gln Ala Val Asn Ser Gly Glu Leu Arg Ala Phe  
 305 310 315 320

Asp Trp Gly Ser Glu Thr Lys Asn Leu Glu Lys Cys Asn Gln Pro Thr  
 325 330 335

Pro Val Arg Tyr Arg Val Arg Asp Met Thr Val Pro Thr Ala Met Trp  
 340 345 350

Thr Gly Gly Gln Asp Trp Leu Ser Asn Pro Glu Asp Val Lys Met Leu  
 355 360 365

Leu Ser Glu Val Thr Asn Leu Ile Tyr His Lys Asn Ile Pro Glu Trp  
 370 375 380

Ala His Val Asp Phe Ile Trp Gly Leu Asp Ala Pro His Arg Met Tyr  
 385 390 395 400

Asn Glu Ile Ile His Leu Met Gln Gln Glu Glu Thr Asn Leu Ser Gln  
 405 410 415

Gly Arg Cys Glu Ala Val Leu  
 420

<210> 48

<211> 33

<212> PRT

<213> Homo sapiens

<400> 48

Met Leu Glu Thr Leu Ser Arg Gln Trp Ile Val Ser His Arg Met Glu  
 1 5 10 15

Met Trp Leu Leu Ile Leu Val Ala Tyr Met Phe Gln Arg Asn Val Asn  
 20 25 30

Ser

<210> 49

<211> 390  
 <212> PRT  
 <213> Homo sapiens

<400> 49

Val	His	Met	Pro	Thr	Lys	Ala	Val	Asp	Pro	Glu	Ala	Phe	Met	Asn	Ile
1				5				10						15	
Ser	Glu	Ile	Ile	Gln	His	Gln	Gly	Tyr	Pro	Cys	Glu	Glu	Tyr	Glu	Val
			20					25					30		
Ala	Thr	Glu	Asp	Gly	Tyr	Ile	Leu	Ser	Val	Asn	Arg	Ile	Pro	Arg	Gly
		35					40					45			
Leu	Val	Gln	Pro	Lys	Lys	Thr	Gly	Ser	Arg	Pro	Val	Val	Leu	Leu	Gln
	50					55					60				
His	Gly	Leu	Val	Gly	Gly	Ala	Ser	Asn	Trp	Ile	Ser	Asn	Leu	Pro	Asn
	65				70					75				80	
Asn	Ser	Leu	Gly	Phe	Ile	Leu	Ala	Asp	Ala	Gly	Phe	Asp	Val	Trp	Met
				85					90					95	
Gly	Asn	Ser	Arg	Gly	Asn	Ala	Trp	Ser	Arg	Lys	His	Lys	Thr	Leu	Ser
			100					105					110		
Ile	Asp	Gln	Asp	Glu	Phe	Trp	Ala	Phe	Ser	Tyr	Asp	Glu	Met	Ala	Arg
			115				120					125			
Phe	Asp	Leu	Pro	Ala	Val	Ile	Asn	Phe	Ile	Leu	Gln	Lys	Thr	Gly	Gln
		130				135					140				
Glu	Lys	Ile	Tyr	Tyr	Val	Gly	Tyr	Ser	Gln	Gly	Thr	Thr	Met	Gly	Phe
	145				150				155					160	
Ile	Ala	Phe	Ser	Thr	Met	Pro	Glu	Leu	Ala	Gln	Lys	Ile	Lys	Met	Tyr
				165					170					175	
Phe	Ala	Leu	Ala	Pro	Ile	Ala	Thr	Val	Lys	His	Ala	Lys	Ser	Pro	Gly
			180					185					190		
Thr	Lys	Phe	Leu	Leu	Leu	Pro	Asp	Met	Met	Ile	Lys	Gly	Leu	Phe	Gly
			195				200					205			
Lys	Lys	Glu	Phe	Leu	Tyr	Gln	Thr	Arg	Phe	Leu	Arg	Gln	Leu	Val	Ile
	210					215					220				
Tyr	Leu	Cys	Gly	Gln	Val	Ile	Leu	Asp	Gln	Ile	Cys	Ser	Asn	Ile	Met

225		230		235		240									
Leu	Leu	Leu	Gly	Gly	Phe	Asn	Thr	Asn	Asn	Met	Asn	Met	Ser	Arg	Ala
			245					250						255	
Ser	Val	Tyr	Ala	Ala	His	Thr	Leu	Ala	Gly	Thr	Ser	Val	Gln	Asn	Ile
			260					265					270		
Leu	His	Trp	Ser	Gln	Ala	Val	Asn	Ser	Gly	Glu	Leu	Arg	Ala	Phe	Asp
			275					280				285			
Trp	Gly	Ser	Glu	Thr	Lys	Asn	Leu	Glu	Lys	Cys	Asn	Gln	Pro	Thr	Pro
			290				295				300				
Val	Arg	Tyr	Arg	Val	Arg	Asp	Met	Thr	Val	Pro	Thr	Ala	Met	Trp	Thr
305					310					315					320
Gly	Gly	Gln	Asp	Trp	Leu	Ser	Asn	Pro	Glu	Asp	Val	Lys	Met	Leu	Leu
			325						330					335	
Ser	Glu	Val	Thr	Asn	Leu	Ile	Tyr	His	Lys	Asn	Ile	Pro	Glu	Trp	Ala
			340					345					350		
His	Val	Asp	Phe	Ile	Trp	Gly	Leu	Asp	Ala	Pro	His	Arg	Met	Tyr	Asn
		355					360					365			
Glu	Ile	Ile	His	Leu	Met	Gln	Gln	Glu	Glu	Thr	Asn	Leu	Ser	Gln	Gly
		370				375					380				
Arg	Cys	Glu	Ala	Val	Leu										
385					390										

<210> 50  
 <211> 221  
 <212> PRT  
 <213> Homo sapiens

<400> 50  
 Val His Met Pro Thr Lys Ala Val Asp Pro Glu Ala Phe Met Asn Ile  
 1 5 10 15  
 Ser Glu Ile Ile Gln His Gln Gly Tyr Pro Cys Glu Glu Tyr Glu Val  
 20 25 30  
 Ala Thr Glu Asp Gly Tyr Ile Leu Ser Val Asn Arg Ile Pro Arg Gly  
 35 40 45

Leu Val Gln Pro Lys Lys Thr Gly Ser Arg Pro Val Val Leu Leu Gln  
 50 55 60

His Gly Leu Val Gly Gly Ala Ser Asn Trp Ile Ser Asn Leu Pro Asn  
 65 70 75 80

Asn Ser Leu Gly Phe Ile Leu Ala Asp Ala Gly Phe Asp Val Trp Met  
 85 90 95

Gly Asn Ser Arg Gly Asn Ala Trp Ser Arg Lys His Lys Thr Leu Ser  
 100 105 110

Ile Asp Gln Asp Glu Phe Trp Ala Phe Ser Tyr Asp Glu Met Ala Arg  
 115 120 125

Phe Asp Leu Pro Ala Val Ile Asn Phe Ile Leu Gln Lys Thr Gly Gln  
 130 135 140

Glu Lys Ile Tyr Tyr Val Gly Tyr Ser Gln Gly Thr Thr Met Gly Phe  
 145 150 155 160

Ile Ala Phe Ser Thr Met Pro Glu Leu Ala Gln Lys Ile Lys Met Tyr  
 165 170 175

Phe Ala Leu Ala Pro Ile Ala Thr Val Lys His Ala Lys Ser Pro Gly  
 180 185 190

Thr Lys Phe Leu Leu Leu Pro Asp Met Met Ile Lys Gly Leu Phe Gly  
 195 200 205

Lys Lys Glu Phe Leu Tyr Gln Thr Arg Phe Leu Arg Gln  
 210 215 220

<210> 51

<211> 25

<212> PRT

<213> Homo sapiens

<400> 51

Leu Val Ile Tyr Leu Cys Gly Gln Val Ile Leu Asp Gln Ile Cys Ser  
 1 5 10 15

Asn Ile Met Leu Leu Leu Gly Gly Phe  
 20 25

<210> 52

<211> 144  
 <212> PRT  
 <213> Homo sapiens

<400> 52

Asn	Thr	Asn	Asn	Met	Asn	Met	Ser	Arg	Ala	Ser	Val	Tyr	Ala	Ala	His
1				5					10					15	
Thr	Leu	Ala	Gly	Thr	Ser	Val	Gln	Asn	Ile	Leu	His	Trp	Ser	Gln	Ala
		20					25					30			
Val	Asn	Ser	Gly	Glu	Leu	Arg	Ala	Phe	Asp	Trp	Gly	Ser	Glu	Thr	Lys
	35					40					45				
Asn	Leu	Glu	Lys	Cys	Asn	Gln	Pro	Thr	Pro	Val	Arg	Tyr	Arg	Val	Arg
50					55						60				
Asp	Met	Thr	Val	Pro	Thr	Ala	Met	Trp	Thr	Gly	Gly	Gln	Asp	Trp	Leu
65					70					75					80
Ser	Asn	Pro	Glu	Asp	Val	Lys	Met	Leu	Leu	Ser	Glu	Val	Thr	Asn	Leu
			85					90						95	
Ile	Tyr	His	Lys	Asn	Ile	Pro	Glu	Trp	Ala	His	Val	Asp	Phe	Ile	Trp
		100					105						110		
Gly	Leu	Asp	Ala	Pro	His	Arg	Met	Tyr	Asn	Glu	Ile	Ile	His	Leu	Met
	115					120					125				
Gln	Gln	Glu	Glu	Thr	Asn	Leu	Ser	Gln	Gly	Arg	Cys	Glu	Ala	Val	Leu
130					135						140				

<210> 53  
 <211> 2133  
 <212> DNA  
 <213> Homo sapiens

<400> 53

gtcgacccac gcgtccacgg cgagggctcc cggggcgcag cattgcccc cctgcaccac 60  
 ctcaccaaga tggctacttt gggacacaca ttcccttct atgctggccc caagccaacc 120  
 ttcccgatgg acaccacttt ggccagcatc atcatgatct ttctgactgc actggccacg 180  
 ttcacgtgca tcttgcttgg cattcgggga aagacgaggg tgttctggct gcttcgggtg 240  
 gtgaccagct tattcatcgg ggctgcaatc ctggctgtga atttcagttc tgagtgggtc 300  
 gtggggccagg tcagcaccaa cacatcatat aaggccttca gttctgagtg gatcagcgct 360

```

gatattgggc tgcaggctcg gctgggtgga gtcaacatca cactcacagg gacccccgtg 420
cagcagctga atgagaccat caattacaac gaggagttca cctggcgccct ggggtgagaac 480
tatgctgagg agtgtgcaaa ggctctggag aaggggctgc cagaccctgt gttgtacct 540
gctgagaagt tcaactccaag aagcccatgt ggcctatacc gccagtaccg cctggcgagg 600
cactacacct cagccatgct atgggtggca ttctctgctt ggctgctggc caatgtgatg 660
ctctccatgc ctgtgctggt atatggtggc tacatgctat tggccacggg catcttccag 720
ctgttggctc tgctcttctt ctccatggcc acatcactca cctcaccctg tcccctgcac 780
ctgggcgctt ctgtgctgca tactcaccat gggcctgcct tctggatcac attgaccaca 840
ggactgctgt gtgtgctgct gggcctggct atggcggtgg cccacaggat gcagcctcac 900
aggctgaagg ctttcttcaa ccagagtgtg gatgaagacc ccatgctgga gtggagtcct 960
gaggaaggtg gactcctgag ccccgctac cggctccatg ctgacagtc caagtcccag 1020
gacattcccc tgtcagaggc ttctccacc aaggcatact gtaaggaggc acacccaaa 1080
gatcctgatt gtgctttata acattcctcc ccgtggaggc cacctggact tccagtctgg 1140
ctccaaacct cattggcgcc ccataaaacc agcagaactg ccctcagggg ggctgttacc 1200
agacaccag caccaatcta cagacggagt agaaaaagga ggctctatat actgatgta 1260
aaaaacaaaa caaaacaaaa agccctaagg gactgaagag atgctggggc tgtccataaa 1320
gcctgttgcc atgataaggc caagcagggg ctagcttata tgcacagcaa cccagccttt 1380
ccgtgctgcc ttgcctcttc aagatgctat tcaactgaaac ctaacttcac ccccataaca 1440
ccagcagggg ggggggttaca tatgattctc ctatggtttc ctctcatccc tggcacctc 1500
ttgttttctt ttttctggtt ttcttttgt tcttcttcta cttctccagc ttgtgtggcc 1560
ttttggtaca atgaaagaca gcaactgaaa ggaggggaaa ccaaacttct catcctaggt 1620
ctaacattaa ccaactatgc cacattctct ttgagcttca gttccaaat ttgctacata 1680
agattgcaag acttgccaag aatcttggga tttatctttc tatgccttgc tgacacctac 1740
cttgccctc aaacaccacc tcacaagaag ccaggtggga agttaggga tcaactccaa 1800
aacgctatct cttccacccc cactcagctg ggctagctga gtggcatcca ggacggggga 1860
gtgggtgacc tgcctcatca ctgccacct acgtccccc ggggtggttc agaaagatgc 1920
tagctctggt agggctcctc cggcctcact agagggcgcc cctattactc tggagtcgac 1980
gcagagaatc aggtttcaca gcaactgcga gagtgacta ggctgtctcc agcccagcga 2040
agctcatgag gacgtgagc cccggcgcg agaaagccatg aaaattaatg ggaaaaacag 2100
tttttaaaaa aaaaaaaaaa aaagggcggc cgc 2133

```

<210> 54  
 <211> 1029  
 <212> DNA  
 <213> Homo sapiens

```

<400> 54
atggctactt tgggacacac attcccttc tatgctggcc ccaagccaac cttcccgatg 60
gacaccactt tggccagcat catcatgatc tttctgactg cactggccac gttcatcgtc 120
atcctgcctg gcattcgggg aaagacgagg ctgttctggc tgcttcgggt ggtgaccagc 180
ttattcatcg gggctgcaat cctggctgtg aatttcagtt ctgagtggtc tgtgggcccag 240
gtcagcacca acacatcata caaggccttc agttctgagt ggatcagcgc tgatattggg 300
ctgcaggctg ggctgggtgg agtcaacatc aactcacag ggacccccgt gcagcagctg 360
aatgagacca tcaattacaa cgaggagttc acctggcgcc tgggtgagaa ctatgctgag 420
gagtgtgcaa aggtctgga gaaggggctg ccagaccctg tgttgtaoct agctgagaag 480
ttcaactcaa gaagcccatg tggcctatac cgccagtacc gcctggcggg acactacacc 540
tcagccatgc tatgggtggc attcctctgc tggctgctgg ccaatgtgat gctctccatg 600
cctgtgctgg tatatggtg ctacatgcta ttggccacgg gcatcttcca gctgttggct 660

```

ctgtctttct tctccatggc cacatcactc acctcaccct gtccccctgca cctgggagct 720  
tctgtgtgtgc atactcacca tgggcctgcc ttctggatca cattgaccac aggactgctg 780  
tgtgtgtgtgc tgggcctggc tatggcggtg gccacagga tgcagcctca caggctgaag 840  
gcttttttca accagagtgt ggatgaagac cccatgctgg agtggagtcc tgaggaaggt 900  
ggactcctga gccccgcta ccggtccatg gctgacagtc ccaagtccca ggacattccc 960  
ctgtcagagg cttcctccac caaggcatac tgtaaggagg cacaccccaa agatcctgat 1020  
tgtgcttta 1029

<210> 55  
<211> 343  
<212> PRT  
<213> Homo sapiens

<400> 55

Met Ala Thr Leu Gly His Thr Phe Pro Phe Tyr Ala Gly Pro Lys Pro  
1 5 10 15

Thr Phe Pro Met Asp Thr Thr Leu Ala Ser Ile Ile Met Ile Phe Leu  
20 25 30

Thr Ala Leu Ala Thr Phe Ile Val Ile Leu Pro Gly Ile Arg Gly Lys  
35 40 45

Thr Arg Leu Phe Trp Leu Leu Arg Val Val Thr Ser Leu Phe Ile Gly  
50 55 60

Ala Ala Ile Leu Ala Val Asn Phe Ser Ser Glu Trp Ser Val Gly Gln  
65 70 75 80

Val Ser Thr Asn Thr Ser Tyr Lys Ala Phe Ser Ser Glu Trp Ile Ser  
85 90 95

Ala Asp Ile Gly Leu Gln Val Gly Leu Gly Gly Val Asn Ile Thr Leu  
100 105 110

Thr Gly Thr Pro Val Gln Gln Leu Asn Glu Thr Ile Asn Tyr Asn Glu  
115 120 125

Glu Phe Thr Trp Arg Leu Gly Glu Asn Tyr Ala Glu Glu Cys Ala Lys  
130 135 140

Ala Leu Glu Lys Gly Leu Pro Asp Pro Val Leu Tyr Leu Ala Glu Lys  
145 150 155 160

Phe Thr Pro Arg Ser Pro Cys Gly Leu Tyr Arg Gln Tyr Arg Leu Ala  
165 170 175

Gly His Tyr Thr Ser Ala Met Leu Trp Val Ala Phe Leu Cys Trp Leu

180	185	190
Leu Ala Asn Val Met Leu Ser Met Pro Val Leu Val Tyr Gly Gly Tyr		
195	200	205
Met Leu Leu Ala Thr Gly Ile Phe Gln Leu Leu Ala Leu Leu Phe Phe		
210	215	220
Ser Met Ala Thr Ser Leu Thr Ser Pro Cys Pro Leu His Leu Gly Ala		
225	230	235 240
Ser Val Leu His Thr His His Gly Pro Ala Phe Trp Ile Thr Leu Thr		
245	250	255
Thr Gly Leu Leu Cys Val Leu Leu Gly Leu Ala Met Ala Val Ala His		
260	265	270
Arg Met Gln Pro His Arg Leu Lys Ala Phe Phe Asn Gln Ser Val Asp		
275	280	285
Glu Asp Pro Met Leu Glu Trp Ser Pro Glu Glu Gly Gly Leu Leu Ser		
290	295	300
Pro Arg Tyr Arg Ser Met Ala Asp Ser Pro Lys Ser Gln Asp Ile Pro		
305	310	315 320
Leu Ser Glu Ala Ser Ser Thr Lys Ala Tyr Cys Lys Glu Ala His Pro		
325	330	335
Lys Asp Pro Asp Cys Ala Leu		
340		

<210> 56  
 <211> 23  
 <212> PRT  
 <213> Homo sapiens

<400> 56  
 Met Ala Thr Leu Gly His Thr Phe Pro Phe Tyr Ala Gly Pro Lys Pro  
 1 5 10 15  
 Thr Phe Pro Met Asp Thr Thr  
 20

<210> 57  
 <211> 112

<212> PRT  
<213> Homo sapiens

<400> 57  
Asn Phe Ser Ser Glu Trp Ser Val Gly Gln Val Ser Thr Asn Thr Ser  
1 5 10 15  
Tyr Lys Ala Phe Ser Ser Glu Trp Ile Ser Ala Asp Ile Gly Leu Gln  
20 25 30  
Val Gly Leu Gly Gly Val Asn Ile Thr Leu Thr Gly Thr Pro Val Gln  
35 40 45  
Gln Leu Asn Glu Thr Ile Asn Tyr Asn Glu Glu Phe Thr Trp Arg Leu  
50 55 60  
Gly Glu Asn Tyr Ala Glu Glu Cys Ala Lys Ala Leu Glu Lys Gly Leu  
65 70 75 80  
Pro Asp Pro Val Leu Tyr Leu Ala Glu Lys Phe Thr Pro Arg Ser Pro  
85 90 95  
Cys Gly Leu Tyr Arg Gln Tyr Arg Leu Ala Gly His Tyr Thr Ser Ala  
100 105 110

<210> 58  
<211> 22  
<212> PRT  
<213> Homo sapiens

<400> 58  
Thr Ser Leu Thr Ser Pro Cys Pro Leu His Leu Gly Ala Ser Val Leu  
1 5 10 15  
His Thr His His Gly Pro  
20

<210> 59  
<211> 19  
<212> PRT  
<213> Homo sapiens

<400> 59

Leu Ala Ser Ile Ile Met Ile Phe Leu Thr Ala Leu Ala Thr Phe Ile  
 1 5 10 15

Val Ile Leu

<210> 60  
 <211> 20  
 <212> PRT  
 <213> Homo sapiens

<400> 60  
 Leu Phe Trp Leu Leu Arg Val Val Thr Ser Leu Phe Ile Gly Ala Ala  
 1 5 10 15

Ile Leu Ala Val  
 20

<210> 61  
 <211> 22  
 <212> PRT  
 <213> Homo sapiens

<400> 61  
 Met Leu Trp Val Ala Phe Leu Cys Trp Leu Leu Ala Asn Val Met Leu  
 1 5 10 15

Ser Met Pro Val Leu Val  
 20

<210> 62  
 <211> 17  
 <212> PRT  
 <213> Homo sapiens

<400> 62  
 Leu Ala Thr Gly Ile Phe Gln Leu Leu Ala Leu Leu Phe Phe Ser Met  
 1 5 10 15

Ala

<210> 63  
 <211> 22

<212> PRT  
<213> Homo sapiens

<400> 63

Ala Phe Trp Ile Thr Leu Thr Thr Gly Leu Leu Cys Val Leu Leu Gly  
1 5 10 15

Leu Ala Met Ala Val Ala  
20

<210> 64  
<211> 8  
<212> PRT  
<213> Homo sapiens

<400> 64

Pro Gly Ile Arg Gly Lys Thr Arg  
1 5

<210> 65  
<211> 6  
<212> PRT  
<213> Homo sapiens

<400> 65

Tyr Gly Gly Tyr Met Leu  
1 5

<210> 66  
<211> 72  
<212> PRT  
<213> Homo sapiens

<400> 66

His Arg Met Gln Pro His Arg Leu Lys Ala Phe Phe Asn Gln Ser Val  
1 5 10 15

Asp Glu Asp Pro Met Leu Glu Trp Ser Pro Glu Glu Gly Gly Leu Leu  
20 25 30

Ser Pro Arg Tyr Arg Ser Met Ala Asp Ser Pro Lys Ser Gln Asp Ile  
35 40 45

Pro Leu Ser Glu Ala Ser Ser Thr Lys Ala Tyr Cys Lys Glu Ala His  
50 55 60

Pro Lys Asp Pro Asp Cys Ala Leu

65

70

<210> 67

<211> 4928

<212> DNA

<213> Mus sp.

<400> 67

gtcgacccac gcgtccgccc ggctcccggg gctgcccct ctgcccggg ccgcgcccgg 60  
gggtcccgcg ctgacggccc atggcgccgc ccgcgcccg tctcgcgctg ctctccgccc 120  
ctgcgctcac tctggcgggc cggcccgcgc ccggtcccgc ctccggcccc gagggttca 180  
cagccaacgg tgcagattac aggggaacac agagctggac agcgtgcaa ggtgggaagc 240  
catgtctgtt ctggaacgag actttccagc atccgtacaa cacgctgaag taccccaacg 300  
gggaaggagg actgggagag cacaattatt gcagaaatcc agatggagac gtgagccctt 360  
ggtgctacgt ggccgagcat gaggacggag tctactggaa gtactgtgaa attcctgcct 420  
gccagatgcc tggaaacctt ggctgctaca aggatcatgg aaaccacct cctctcacgg 480  
gcaccagtaa aacctctaac aagctcacca tacaacctg tatcagcttc tgcggagtc 540  
agagattcaa gtttgctggg atggagtcag gctatgcctg cttctgtggg aacaatcctg 600  
actactgaa gcacggggag gcggccagca ccgagtcaa tagtgtctgc ttcggggacc 660  
acacgcagcc ctgcggtggg gacggcagga ttatcctctt tgacactctc gtgggagcct 720  
gcggtgggaa ctactcagcc atggcagccg tgggtgactc ccctgacttc cctgacacct 780  
acgccactgg cagagtctgc tactggacca tccgggttcc aggagcctct cgcattccatt 840  
tcaacttcac cctgtttgat atcagggaact ctgcagacat ggtggagctg ctggacggct 900  
acaccaccg cgtcctggtc cggctcagtg ggaggagccg ccgcctctg tctttcaatg 960  
tctctctgga ttttgtcatt ttgtatttct tctctgatcg catcaatcag gccagggat 1020  
ttgctgtgtt gtaccaagcc accaaggagg aaccgccaca ggagagacct gctgtcaacc 1080  
agacctggc agaggtgatc accgagcaag ccaacctcag tgtcagcgt gccactcct 1140  
ccaaagtct ctatgtcatc acccccagcc ccagccacct tccgcagact gccaggtag 1200  
ccattcctgg gcaccgtcag ttggggccaa cagccacaga gtggaaggat ggactgtgta 1260  
cggcctggcg accctcctca tctcagagt cacagcagtt gtgcgaaaga ttcttctgca 1320  
tgtcacattt aaatctcatc gaggccctgc atcaggagac cttagggact gtcgtcagcc 1380  
tggggcttct ggagatatct ggaccatttt ctatgaacct tccactaaa tctccatctt 1440  
taagaagaag ctcaagggtc agagtcaaca agatgacgcg aatcccctcg tgagtgactg 1500  
aagcccacgc ctgcatgaga ggctccgctc caagctcgag tttgctcccc tgagttctcc 1560  
tctgatgagt tccctgcctt cctattcacc accatctctt ttgggagcac cctgcttttag 1620  
aggcagcca gcctgggatc ctccatcaca tgtaccagcc tggctgctct gctggggatg 1680  
gtaagacagg cccaggctga caggacacag ctggacctga ctccagaaga ctcttgggtg 1740  
gtggggagggt atagtgtagg atgagttttc ttgcttcttc tctgttttgt ccacatacag 1800  
atcggtttcc cctgtcttta cagtttgcaa tagagccaga ctgaaagaac tgtcaggttt 1860  
tctaggctgg cctggttccc cactaagagt ggcattggcg ccttagaggc ccagaggccc 1920  
agtgtaggct tggagctttc tctgtgcca actaccatgt gtcattctagt ccgaggggac 1980  
tgagagcagg gccacaccag atgtcatctt tctagagggt tcttttttagt acccactgac 2040  
caatggggca agcctgagga ttggtccatc tgtttgtcca tggacagac acagtgaact 2100  
tcctggatac tagacttaac tagcctagcc ctcaagtagt tgccaatcct gtggaatcag 2160  
aattcagcct gtcttctgtt cctcagccca agcctgtagc ctagagctgg ggctgtagcc 2220

tagagctggg	gctgtagcct	agagctgggg	ctgtagcaca	gagctggggc	tgtagcctag	2280
agctggggct	gtagcacaga	gctggggctg	tagcctagag	ctggggctgt	agcacagagc	2340
tggggctgta	gcacagagct	ggggctgtag	cctagagctg	gggctgtagc	acagagctgg	2400
ggctgtaact	cagcgatcaa	gagcttgctt	tgtatacatc	ggaccctagg	ttctatccca	2460
gcactatcag	aaggtgggag	agaaaaagac	tgaccatag	catgcgggca	gcactctgtg	2520
ttcctacgtg	aggtgtcatc	attttaaaag	cagatcaaaa	ctaccgagag	ttttgtcctt	2580
tgtcccttat	catgggagca	gagtaggagt	aagggtctctg	gtcttgctca	ttgtcccca	2640
gacagggagg	caggaaaagg	tcaggcttgg	gaactggaga	tcctcccagg	aaaagctgca	2700
agattgagag	accagctgc	agttgggaga	ggaaggcca	tcccogactg	agaagtcctg	2760
cagtctggaa	gtggcctttg	tcagcagcag	ctgtgccctg	aaggtagacc	ttggctcactc	2820
tcctgccagc	ccttgagcct	ctgctctcct	gggtaccctc	ctggaacacc	atgctaacct	2880
tccccgagtc	tctcagtcac	tgccattgag	gcctctcctc	tagctgctgc	tccccaggac	2940
tgtctggggc	catctgggga	tcagggagag	gcagcaggag	tactgacgag	gcagtgcact	3000
gagctgatga	gtcaaccaga	ggacaccaga	gtctacagtg	ggctggctgc	tggctcagct	3060
cctatgggag	gcctacaggg	gtactaagct	aggggtcat	catctcattt	gatctgggaa	3120
aggctacagg	ctcctggatg	tgaagacagg	cccactacat	aagaagacca	ctggaaatag	3180
actgacagga	gcaggttcca	ctctaggctg	tccatagcgt	ttgcaggact	cccctgagac	3240
caagtgttga	gtcacagagt	gccatgtgcg	tagtgcataa	aggatatggg	ttcttaacca	3300
gggaaggctc	atagcaggcc	aggacatttt	ttcagctcag	agcactggcc	ccaggcttcc	3360
tctaagccac	cactcacctg	tctcttccca	tctcggacac	aggaagcaag	cccagtggtg	3420
gtggcagctg	cggctcagca	ttggtgtccc	caggaagggc	ggtggatgtg	cccacgctcc	3480
ttttgctgtg	ggcctggcac	agcccaacac	tgaggggccc	accttctctc	ttggggggta	3540
gggacacata	aggaaaacta	accacactcc	aacaacagca	gaggacagtg	ggaaggaagg	3600
gctgtaaata	accaggcca	gacctccaga	aatgacaggc	acagtctgtt	agaacctgta	3660
ggcagccagt	cacagagggc	ctttgtgctg	gtaacaccct	gcctggagca	taggggtaag	3720
ccgagggaga	agagcagccc	tcagagacat	cagctaaaaa	cataggtgcc	ctatgtccct	3780
cccttctctg	cacactgctt	acaaagcaga	gacagagtag	gaaagaggtc	ttcatcctct	3840
cccacatcag	caaggatagg	gctgcggctg	cctaaagtga	gcaaggagaa	cagagctctg	3900
gacttctcta	aatgtgggct	ctggcttcag	actcctcagc	caaaagctct	tgaagatcaa	3960
agctctggcg	ggtacagctg	tcctggcctg	tgggccagcc	catgggatgt	gcctgggcca	4020
ggtgccaccc	cacggctcac	tgtcatccca	ggagggaccc	cacctgatgc	tcctcatcat	4080
ccgctggcct	gacactatca	gagctcgctc	cggctgttgc	cagggacaga	ctgactacac	4140
ttgaccttca	agagcactta	gaagtggatg	gcctccagac	tctgtcagcc	tctgcagggg	4200
ccacacaagt	ctcccagacc	aagtccacaa	gcctccatgg	ttccctggct	cctctcctgt	4260
ggagtgtcct	gtttgatgtc	tgaggtctgc	tttgggtacc	gccctgggaa	ctgctaacct	4320
ccgattggtc	cctttgtgtc	tctgtttact	gtcctcttct	acctccaggt	cacttagctc	4380
tggctgctct	ggctgggagt	gggggggtgg	gatgctggct	gcacccccac	cctgggtctgc	4440
caacagaacc	tgggggcctc	acacgggctc	ctgtcttgcc	aagctggagc	tgagcacact	4500
ggcccaggct	gagtggggca	gagcaaacaa	gtggaagggg	atctctctcc	ttagagggag	4560
gtggccgaag	gtgtagatcc	agcgaggggag	ctgccatccc	cgccaccttc	atagcagcaa	4620
gaccttccca	tttccaatct	caccctccag	cagggatatg	actttggaca	acaaggcttt	4680
atttgtaaat	atgctcttaa	tatgcaactt	tgagaataag	atagaaacat	catgtatttt	4740
aaaatataaa	atgaagtgtg	acacactgta	tacaatttaa	tatatatttt	taggattttg	4800
ttatttaaga	aaatggaatg	tgatggtact	taacttttac	aaaagagaga	aaatgttatt	4860
tttactgttt	gaagaaaata	aatattctca	ttgtttaga	aaaaaaaaa	aaaaaaagg	4920
gcggccgc						4928

<210> 68

<211> 1410  
 <212> DNA  
 <213> Mus sp.

<400> 68

```

atggcgccgc ccgcccgcgc tctcgcgctg ctctccgcgc ctgcgctcac tctggcggcc 60
cgccccgcgc ccggtccccg ctccggcccc gagtgcttca cagccaacgg tgcagattac 120
aggggaacac agagctggac agcgctgcaa ggtgggaagc catgtctgtt ctggaacgag 180
actttccagc atccgtacaa cacgctgaag taccocaaacg gggaaggagg actgggcgag 240
cacaattatt gcagaaatcc agatggagac gtgagccctt ggtgctacgt ggccgagcat 300
gaggacggag tctactggaa gtactgtgaa attcctgcct gccagatgcc tggaaacctt 360
ggctgctaca aggatcatgg aaacccacct cctctcacgg gcaccagtaa aacctctaac 420
aagctcacca taaaaacctg tatcagcttc tgtcggagtc agagattcaa gtttgcgggg 480
atggagtcag gctatgcctg cttctgtggg aacaatcctg actactggaa gcacggggag 540
gcgccagca ccgagtgcaa tagtgtctgc ttccggggacc acacgcagcc ctgcgggtggg 600
gacggcagga ttatcctctt tgacactctc gtgggcgcct gcggtgggaa ctactcagcc 660
atggcagccg tgggtgtactc ccctgacttc cctgacacct acgccactgg cagagtctgc 720
tactggacca tccgggttcc aggagcctct cgcattccatt tcaacttcac cctgtttgat 780
atcagggact ctgcagacat ggtggagctg ctggacggct acaaccaccg cgtcctggtc 840
cggctcagtg ggaggagccg cccgcctctg tctttcaatg tctctctgga ttttgtcatt 900
ttgtatttct tctctgatcg catcaatcag gccagggat ttgctgtgtt gtaccaagcc 960
accaaggagg aaccgccaca ggagagacct gctgtcaacc agaccctggc agaggtgatc 1020
accgagcaag ccaacctcag tgtcagcgct gccactcct ccaaagtcct ctatgtcatc 1080
acccccagcc ccagccaccc tccgcagact gccaggtag ccattcctgg gcaccgtcag 1140
ttggggccaa cagccacaga gtggaaggat ggactgtgta cggcctggcg accctcctca 1200
tcctcacagt cacagcagtt gtcgcaaaga ttcttctgca tgtcacattt aaatctcatc 1260
gagtcctctg atcaggagac cttagggact gtcgtcagcc tggggcttct ggagatatct 1320
ggaccatttt ctatgaacct tccactacaa tctccatctt taagaagaag ctcaagggtc 1380
agagtcaaca agatgaccgc aatcccctcg                                     1410
  
```

<210> 69  
 <211> 470  
 <212> PRT  
 <213> Mus sp.

<400> 69

```

Met Ala Pro Pro Ala Ala Arg Leu Ala Leu Leu Ser Ala Ala Ala Leu
  1             5             10            15

Thr Leu Ala Ala Arg Pro Ala Pro Gly Pro Arg Ser Gly Pro Glu Cys
          20            25            30

Phe Thr Ala Asn Gly Ala Asp Tyr Arg Gly Thr Gln Ser Trp Thr Ala
    35            40            45

Leu Gln Gly Gly Lys Pro Cys Leu Phe Trp Asn Glu Thr Phe Gln His
    50            55            60
  
```



Thr Lys Glu Glu Pro Pro Gln Glu Arg Pro Ala Val Asn Gln Thr Leu  
 325 330 335

Ala Glu Val Ile Thr Glu Gln Ala Asn Leu Ser Val Ser Ala Ala His  
 340 345 350

Ser Ser Lys Val Leu Tyr Val Ile Thr Pro Ser Pro Ser His Pro Pro  
 355 360 365

Gln Thr Ala Gln Val Ala Ile Pro Gly His Arg Gln Leu Gly Pro Thr  
 370 375 380

Ala Thr Glu Trp Lys Asp Gly Leu Cys Thr Ala Trp Arg Pro Ser Ser  
 385 390 395 400

Ser Ser Gln Ser Gln Gln Leu Ser Gln Arg Phe Phe Cys Met Ser His  
 405 410 415

Leu Asn Leu Ile Glu Ser Leu His Gln Glu Thr Leu Gly Thr Val Val  
 420 425 430

Ser Leu Gly Leu Leu Glu Ile Ser Gly Pro Phe Ser Met Asn Leu Pro  
 435 440 445

Leu Gln Ser Pro Ser Leu Arg Arg Ser Ser Arg Val Arg Val Asn Lys  
 450 455 460

Met Thr Ala Ile Pro Ser  
 465 470

<210> 70  
 <211> 760  
 <212> PRT  
 <213> Mus sp.

<400> 70  
 Met Ala Leu Pro Ser Leu Gly Gln Asp Ser Trp Ser Leu Leu Arg Val  
 1 5 10 15

Phe Phe Phe Gln Leu Phe Leu Leu Pro Ser Leu Pro Pro Ala Ser Gly  
 20 25 30

Thr Gly Gly Gln Gly Pro Met Pro Arg Val Lys Tyr His Ala Gly Asp  
 35 40 45

Gly His Arg Ala Leu Ser Phe Phe Gln Gln Lys Gly Leu Arg Asp Phe  
 50 55 60

Asp Thr Leu Leu Leu Ser Asp Asp Gly Asn Thr Leu Tyr Val Gly Ala  
65 70 75 80

Arg Glu Thr Val Leu Ala Leu Asn Ile Gln Asn Pro Gly Ile Pro Arg  
85 90 95

Leu Lys Asn Met Ile Pro Trp Pro Ala Ser Glu Arg Lys Lys Thr Glu  
100 105 110

Cys Ala Phe Lys Lys Lys Ser Asn Glu Thr Gln Cys Phe Asn Phe Ile  
115 120 125

Arg Val Leu Val Ser Tyr Asn Ala Thr His Leu Tyr Ala Cys Gly Thr  
130 135 140

Phe Ala Phe Ser Pro Ala Cys Thr Phe Ile Glu Leu Gln Asp Ser Leu  
145 150 155 160

Leu Leu Pro Ile Leu Ile Asp Lys Val Met Asp Gly Lys Gly Gln Ser  
165 170 175

Pro Leu Thr Leu Phe Thr Ser Thr Gln Ala Val Leu Val Asp Gly Met  
180 185 190

Leu Tyr Ser Gly Thr Met Asn Asn Phe Leu Gly Ser Glu Pro Ile Leu  
195 200 205

Met Arg Thr Leu Gly Ser His Pro Val Leu Lys Thr Asp Ile Phe Leu  
210 215 220

Arg Trp Leu His Ala Asp Ala Ser Phe Val Ala Ala Ile Pro Ser Thr  
225 230 235 240

Gln Val Val Tyr Phe Phe Phe Glu Glu Thr Ala Ser Glu Phe Asp Phe  
245 250 255

Phe Glu Glu Leu Tyr Ile Ser Arg Val Ala Gln Val Cys Lys Asn Asp  
260 265 270

Val Gly Gly Glu Lys Leu Leu Gln Lys Lys Trp Thr Thr Phe Leu Lys  
275 280 285

Ala Gln Leu Leu Cys Ala Gln Pro Gly Gln Leu Pro Phe Asn Ile Ile  
290 295 300

Arg His Ala Val Leu Leu Pro Ala Asp Ser Pro Ser Val Ser Arg Ile  
305 310 315 320

Tyr Ala Val Phe Thr Ser Gln Trp Gln Val Gly Gly Thr Arg Ser Ser  
 325 330 335

Ala Val Cys Ala Phe Ser Leu Thr Asp Ile Glu Arg Val Phe Lys Gly  
 340 345 350

Lys Tyr Lys Glu Leu Asn Lys Glu Thr Ser Arg Trp Thr Thr Tyr Arg  
 355 360 365

Gly Ser Glu Val Ser Pro Arg Pro Gly Ser Cys Ser Met Gly Pro Ser  
 370 375 380

Ser Asp Lys Ala Leu Thr Phe Met Lys Asp His Phe Leu Met Asp Glu  
 385 390 395 400

His Val Val Gly Thr Pro Leu Leu Val Lys Ser Gly Val Glu Tyr Thr  
 405 410 415

Arg Leu Ala Val Glu Ser Ala Arg Gly Leu Asp Gly Ser Ser His Val  
 420 425 430

Val Met Tyr Leu Gly Thr Ser Thr Gly Pro Leu His Lys Ala Val Val  
 435 440 445

Pro Gln Asp Ser Ser Ala Tyr Leu Val Glu Glu Ile Gln Leu Ser Pro  
 450 455 460

Asp Ser Glu Pro Val Arg Asn Leu Gln Leu Ala Pro Ala Gln Gly Ala  
 465 470 475 480

Val Phe Ala Gly Phe Ser Gly Gly Ile Trp Arg Val Pro Arg Ala Asn  
 485 490 495

Cys Ser Val Tyr Glu Ser Cys Val Asp Cys Val Leu Ala Arg Asp Pro  
 500 505 510

His Cys Ala Trp Asp Pro Glu Ser Arg Leu Cys Ser Leu Leu Ser Gly  
 515 520 525

Ser Thr Lys Pro Trp Lys Gln Asp Met Glu Arg Gly Asn Pro Glu Trp  
 530 535 540

Val Cys Thr Arg Gly Pro Met Ala Arg Ser Pro Arg Arg Gln Ser Pro  
 545 550 555 560

Pro Gln Leu Ile Lys Glu Val Leu Thr Val Pro Asn Ser Ile Leu Glu  
 565 570 575

Leu Arg Cys Pro His Leu Ser Ala Leu Ala Ser Tyr His Trp Ser His  
580 585 590

Gly Arg Ala Lys Ile Ser Glu Ala Ser Ala Thr Val Tyr Asn Gly Ser  
595 600 605

Leu Leu Leu Leu Pro Gln Asp Gly Val Gly Gly Leu Tyr Gln Cys Val  
610 615 620

Ala Thr Glu Asn Gly Tyr Ser Tyr Pro Val Val Ser Tyr Trp Val Asp  
625 630 635 640

Ser Gln Asp Gln Pro Leu Ala Leu Asp Pro Glu Leu Ala Gly Val Pro  
645 650 655

Arg Glu Arg Val Gln Val Pro Leu Thr Arg Val Gly Gly Gly Ala Ser  
660 665 670

Met Ala Ala Gln Arg Ser Tyr Trp Pro His Phe Leu Ile Val Thr Val  
675 680 685

Leu Leu Ala Ile Val Leu Leu Gly Val Leu Thr Leu Leu Leu Ala Ser  
690 695 700

Pro Leu Gly Ala Leu Arg Ala Arg Gly Lys Val Gln Gly Cys Gly Met  
705 710 715 720

Leu Pro Pro Arg Glu Lys Ala Pro Leu Ser Arg Asp Gln His Leu Gln  
725 730 735

Pro Ser Lys Asp His Arg Thr Ser Ala Ser Asp Val Asp Ala Asp Asn  
740 745 750

Asn His Leu Gly Ala Glu Val Ala  
755 760

<210> 71  
<211> 3046  
<212> DNA  
<213> Mus sp.

<400> 71  
ctcggacgcc tgggttaggg gtctgtactg ctggggaacc atctggtgac catctcaggc 60  
tgaccatggc cctaccatcc ctgggccagg actcatggag tctcctgcgt gtttttttct 120  
tccaactott cctgctgcc tcaactgccac ctgcttctgg gactggtggt caggggcca 180  
tgcccagagt caaataccat gctggagacg ggcacagggc cctcagcttc ttccaacaaa 240

```

aaggcctccg agactttgac acgctgctcc tgagtgcga tggcaacact ctctatgtgg 300
gggctcgaga gaccgtcctg gccttgaata tccagaaccc aggaatccca aggctaaaga 360
acatgatacc ctggccagcc agtgagagaa aaaagaccga atgtgccttt aagaagaaga 420
gcaatgagac acagtgtttc aacttcattc gagtctggt ctcttacaat gctactcacc 480
tctatgcctg tgggaccttt gccttcagcc ctgcctgtac cttcattgaa ctccaagatt 540
ccctcctgtt gcccatcttg atagacaagg tcatggacgg gaagggccaa agccctttga 600
ccctgttcac aagcacacaa gctgtcttg tcatgggat gctttattcc ggcaccatga 660
acaacttcct gggcagcgag cccatcctga tgcggacact gggatcccat cctgttctca 720
agactgacat cttcttacgc tggctgcacg cggatgcctc cttcgtggca gccattccat 780
ccaccaggt cgtctatttc ttctttgagg agacagccag cgagtttgac ttctttgaag 840
agctgtatat atccagggtg gctcaagtct gcaagaacga cgtgggagggt gaaaagctgc 900
tgcagaagaa gtggaccacc ttctcaaaag cccagttgct ctgcgctcag ccagggcagc 960
tgccattcaa catcatccgc cacgcggtcc tgctgcccgc cgattctccc tctgtttccc 1020
gcatctacgc agtctttacc tcccagtggc aggttgccgc gaccaggagc tcagcagctc 1080
gtgccttctc tctcacggac attgagcgag tctttaaagg gaagtacaag gagctgaaca 1140
aggagacctc ccgctggacc acttacggg gctcagaggt cagcccgagg ccaggcagtt 1200
gctccatggg cccctcctct gacaaagcct tgaccttcac gaaggacct tttctgatgg 1260
atgagcacgt ggtaggaaca cccctgctgg tgaagtctgg tgtggagtac acacggcttg 1320
ctgtggagtc agctcggggc cttgatggga gcagccatgt ggtcatgtat ctgggtacct 1380
ccacgggtcc cctgcacaag gctgtggtgc ctcaggacag cagtgccttat ctcgtggagg 1440
agattcagct gagccctgac tctgagcctg ttcgaaacct gcagctggcc cccgcccagg 1500
gtgcagtggt tgcaggcttc tctggaggca tctggagagt tcccagggcc aattgcagtg 1560
totacgagag ctgtgtggac tgtgtgcttg ccagggaccc tcaactgtgc tgggaccttg 1620
aatcaagact ctgcagcctt ctgtctggct ctaccaagcc ttggaagcag gacatggaac 1680
gcggaacccc ggagtgggta tgcacccgtg gcccctatgg caggagcccc cggcgctcaga 1740
gccccctca actaattaaa gaagtcctga cagtcccaa ctccatcctg gagctgcgct 1800
gccccacct gtcagcactg gcctcttacc actggagtca tggccgagcc aaaatctcag 1860
aagcctctgc taccgtctac aatggctccc tcttgtctgt gccgcaggat ggtgtcgggg 1920
gcctctacca gtgtgtggcg actgagaacg gctactcata ccctgtggtc tccatttggg 1980
tagacagcca ggaccagccc ctggcgctgg accctgagct ggcggcggtt ccccgctgagc 2040
gtgtgcaggt cccgctgacc agggctggag gcggagcttc catggtgtgc cagcggtcct 2100
actggcccca ttttctcctc gttacogtcc tccctggccat cgtgctcctg ggagtgtctc 2160
ctctcctcct cgtctcccca ctggggggcg tgccggctcg gggtaagggt cagggtgtgt 2220
ggatgtgtgc cccaggggaa aaggctccac tgagcaggga ccagcacctc cagccctcca 2280
aggaccacag gacctctgcc agtgacgtag atgccgacaa caaccatctg ggcgccgaag 2340
tggtctaaac agggacacag atccgcagct gagcagagca agccactggc cttgttggt 2400
atgccaggca cagtgccact ctgaccaggg taggaggctc tcctgctaac gtgtgtcacc 2460
tacagacccc agtaggtcct cccctgtggg actctcttct gcaagcacat tgggtgtgt 2520
ccatacctgt acttgtgtg tgacaggaag agccagacag gtttctttga ttttgattga 2580
cccaagagcc ctgcctgtaa caaacgtgct ccaggagacc atgaaagggt tggctgtctg 2640
ggattctgtg gtgacaaacc taagcatccg agcaagctgg ggctattcct gcaaactcca 2700
tcctgaacgc tgtcactcta gaagcagctg ctgctttgaa caccagccca cccctcctcc 2760
caagagtctc tatggagttg gccccttggt tttcctttac cagtccgggc atactgtttg 2820
ggaagtcatc tctgaagtct aaccaccttc cttcttggtt cagtttggtg agattgttat 2880
tattgtctct gccctggcta gaatgggggc ataactctgag ccttgttccc ttgtccagt 2940
tggctgaccc ttgacctct ccttctcct cccttggtt tgggattcag aaaactgctt 3000
gtcacagaca atttatTTTT tattaAAAAA gatataagct ttaaaag 3046

```

<210> 72  
 <211> 2915  
 <212> DNA  
 <213> Mus sp.

<400> 72

```

gtcgaaccac gcgtccggcc gcgcgtcctt ctgccggcgt cagctcgtat ccccgagatc 60
caccgcgccg tcccgggggtg cggactggcc ctgagctggc cgtacagccc ggcttcggac 120
ggtcctcgct ggagccatgg gccgcgggct cggcaggggt gcggcgctgc tgctcgggct 180
gctagtggag tgcactgagg ccaaaaaaca ttgctggtat tttgaaggac tctatccac 240
atactatata tgccgttcct atgaagactg ctgtggtctc aggtgctgtg tgagggccct 300
ttccatacag aggctgtggt attttttggt cctgctgatg atgggtgtgc tgttctgctg 360
tggtgccggg ttcttcattc gccggcgcat gtatccgcca ccactcattg aggagccac 420
attcaatgtg tcctatacca ggcagccacc aaatcctgt ccaggagcac agcaaattgg 480
accgccatat tacaccgacc ctggaggacc cgggatgaat cctggttgga ataccatggc 540
tatggctttc caggtccagc ccaattcacc tcacggaggc acaacttacc caccctctcc 600
ttcctactgc aacacgcctc caccctccta tgaacagggt gtgaaggaca agtagcaaga 660
tgctacatca aaggcaaaga ggatggacag gcccttttgt ttaccttccc atcctcaccg 720
atacttgctg ataggggtggt ccaagggaaa acttggtatg tctcaaagca agcccagctc 780
tctttcaagt cttttgtgga ggacatttga atccacactg tctcctctgt tgcttctgtt 840
tctgatgtag tctgtgctct ctgagagagt gtggcaacag tccctgaggg ttgatattcc 900
taggggtgtc agggtagatc ctcgaggag aggctaagg gaaaggaagg catagcctgt 960
gtgttagggg gcagataaag tggtcaggct gagataagac tcacatgatg cagtagttgg 1020
cagtgaactt cgaagagaca ctatccacca tcccagccca ttctcctaag agaagctgtg 1080
gggctgtgtt gttgatgtct tttggtctcc actcacattt tgaaaatagg ctttctctctg 1140
caggaatagg aaagacccaa gtacatattt gcttccactt aaaaatgagg gtcagaacca 1200
ggcctcagtt ggacatctat agttaaataa aggccattag agaggggaaa tctttaagtt 1260
aggggaaatt ctctaaatgg agacattgcg ttttatgaat catcgtctgg cttttctttt 1320
agtgcattga ttgaagttag ggtgtccttt gagatcagat ggggagagtg aactctgcgg 1380
ggggtggggg gtctctactc agagggctcc aacacccttt tcttaggtag ttctggtgat 1440
gggttttatg ggcactatag agctgagggg cacattaggc cgggtagtta cattgaccct 1500
tgagagaggaa gaggacagcc aaagaaactc agcaaagcaa gaccagcatt gctgagttag 1560
agctaggggt gtatgtgatc ccaacagaga tgtgctggcc tcagaagagg ggacgtttgt 1620
ggatagagcc gtgaaaacct acttagttgc acagatgaca taatcaaaag tagagaaaga 1680
agtgtagtta gagatgccat ttcccagggt agaatcagag ctcatccata gatttacaag 1740
tagtggctgg agttaacagt atggagttct tttcccttgc gtagttagtc acgttgatgt 1800
gtattttaa acaggttgag acctgtgtga ctaagagcaa ggaagtatag ctaagatgtc 1860
tagattattt atatgtagta tgggtggggg tggggctgca aggaaggggg ctgacattgt 1920
aatgagaaa atcagagcca tttgataaac tgttacttgt tggatcaggc atccaaaagt 1980
gtctcttgag tggacattga gtattcttta ccacctacaa gaccaggagg catggtgtca 2040
ttctocattg gggatattat atgaggtaga gggtcaggaa tcgacagtag ctgtgtgggc 2100
ttagtttaag gactgaaagc atagggactg gtagacagtt tcataggaaa ctgcggggaa 2160
ggaatggata cttttaaaga cagtttgttg atgcagatgc tgccacccat cattgagcac 2220
ccttgtgtct ctggcttcct gtcactggat ccagtacccc tccatgcttg ggtccttgtt 2280
ttacataaga caacaaagca caatgtctgc tgtttacaat caagacgact acatggtcca 2340
aacatttctt ctctcttcta tcaattgttg ctttaacttc catttctctc gttccttttt 2400
aaaatcaaga agcacagtca gagctgcccc tgggattgca tcagggaaag gctgatcaag 2460
gcattcagtg tccatgacta aatcttatct ttttgatagc aaatcctttt aagaaactga 2520

```

acaattgcta aggctcagca attttatact ccaatgtctg tgtaaggtaa attttgtttg 2580  
ccattgagcc cacattggaa ttccttctga cgtcaacact gacaatgcct atggaaattg 2640  
cacttctggg tatatgtccc agcatccttg ttttcttatg tttggtagt aaggctcacc 2700  
ccttccagca gctctacttc tgtgtgctga ggctcgttag agccggggct tgggcacaga 2760  
catgaggcag acttgtgcat gctctttctt ggcaacactt ggctcatatt tcttgttctc 2820  
ttttgataga gtcctgtttc ctatgtattt aaaaaataat aaaagtgaat ttagtcaaaa 2880  
aaaaaaaaaa aaaaaaaaaa aaaaagggcg gccgc 2915

<210> 73  
<211> 516  
<212> DNA  
<213> Mus sp.

<400> 73  
atgggccgcc ggctcggcag ggtggcggcg ctgctgctcg ggctgctagt ggagtgcact 60  
gaggccaaaa aacattgctg gtattttgaa ggactctatc ccacatacta tatatgccgt 120  
tcctatgaag actgctgtgg ctccagggtc tgtgtgaggg ccctttccat acagaggctg 180  
tggtattttt ggttcctgct gatgatgggt gtgctgttct gctgtggtgc cggtttcttc 240  
attcgccggc gcatgtatcc gccaccactc attgaggagc ccacattcaa tgtgtcctat 300  
accaggcagc caccaaatcc tgctccagga gcacagcaaa tgggaccgcc atattacacc 360  
gaccctggag gaccgggat gaatcctgtt ggcaatacca tggctatggc tttccaggtc 420  
cagcccaatt cacctcacgg aggcacaact taccaccccc ctcttctcta ctgcaacacg 480  
cctccacccc cctatgaaca ggtggtgaag gacaag 516

<210> 74  
<211> 172  
<212> PRT  
<213> Mus sp.

<400> 74  
Met Gly Arg Arg Leu Gly Arg Val Ala Ala Leu Leu Leu Gly Leu Leu  
1 5 10 15  
Val Glu Cys Thr Glu Ala Lys Lys His Cys Trp Tyr Phe Glu Gly Leu  
20 25 30  
Tyr Pro Thr Tyr Tyr Ile Cys Arg Ser Tyr Glu Asp Cys Cys Gly Ser  
35 40 45  
Arg Cys Cys Val Arg Ala Leu Ser Ile Gln Arg Leu Trp Tyr Phe Trp  
50 55 60  
Phe Leu Leu Met Met Gly Val Leu Phe Cys Cys Gly Ala Gly Phe Phe  
65 70 75 80  
Ile Arg Arg Arg Met Tyr Pro Pro Pro Leu Ile Glu Glu Pro Thr Phe  
85 90 95

Asn Val Ser Tyr Thr Arg Gln Pro Pro Asn Pro Ala Pro Gly Ala Gln  
100 105 110

Gln Met Gly Pro Pro Tyr Tyr Thr Asp Pro Gly Gly Pro Gly Met Asn  
115 120 125

Pro Val Gly Asn Thr Met Ala Met Ala Phe Gln Val Gln Pro Asn Ser  
130 135 140

Pro His Gly Gly Thr Thr Tyr Pro Pro Pro Pro Ser Tyr Cys Asn Thr  
145 150 155 160

Pro Pro Pro Pro Tyr Glu Gln Val Val Lys Asp Lys  
165 170

<210> 75  
<211> 398  
<212> PRT  
<213> Homo sapiens

<400> 75  
Met Trp Leu Leu Leu Thr Met Ala Ser Leu Ile Ser Val Leu Gly Thr  
1 5 10 15

Thr His Gly Leu Phe Gly Lys Leu His Pro Gly Ser Pro Glu Val Thr  
20 25 30

Met Asn Ile Ser Gln Met Ile Thr Tyr Trp Gly Tyr Pro Asn Glu Glu  
35 40 45

Tyr Glu Val Val Thr Glu Asp Gly Tyr Ile Leu Glu Val Asn Arg Ile  
50 55 60

Pro Tyr Gly Lys Lys Asn Ser Gly Asn Thr Gly Gln Arg Pro Val Val  
65 70 75 80

Phe Leu Gln His Gly Leu Leu Ala Ser Ala Thr Asn Trp Ile Ser Asn  
85 90 95

Leu Pro Asn Asn Ser Leu Ala Phe Ile Leu Ala Asp Ala Gly Tyr Asp  
100 105 110

Val Trp Leu Gly Asn Ser Arg Gly Asn Thr Trp Ala Arg Arg Asn Leu  
115 120 125

Tyr Tyr Ser Pro Asp Ser Val Glu Phe Trp Ala Phe Ser Phe Asp Glu  
130 135 140

Met Ala Lys Tyr Asp Leu Pro Ala Thr Ile Asp Phe Ile Val Lys Lys  
145 150 155 160

Thr Gly Gln Lys Gln Leu His Tyr Val Gly His Ser Gln Gly Thr Thr  
165 170 175

Ile Gly Phe Ile Ala Phe Ser Thr Asn Pro Ser Leu Ala Lys Arg Ile  
180 185 190

Lys Thr Phe Tyr Ala Leu Ala Pro Val Ala Thr Val Lys Tyr Thr Lys  
195 200 205

Ser Leu Ile Asn Lys Leu Arg Phe Val Pro Gln Ser Leu Phe Lys Phe  
210 215 220

Ile Phe Gly Asp Lys Ile Phe Tyr Pro His Asn Phe Phe Asp Gln Phe  
225 230 235 240

Leu Ala Thr Glu Val Cys Ser Arg Glu Met Leu Asn Leu Leu Cys Ser  
245 250 255

Asn Ala Leu Phe Ile Ile Cys Gly Phe Asp Ser Lys Asn Phe Asn Thr  
260 265 270

Ser Arg Leu Asp Val Tyr Leu Ser His Asn Pro Ala Gly Thr Ser Val  
275 280 285

Gln Asn Met Phe His Trp Thr Gln Ala Val Lys Ser Gly Lys Phe Gln  
290 295 300

Ala Tyr Asp Trp Gly Ser Pro Val Gln Asn Arg Met His Tyr Asp Gln  
305 310 315 320

Ser Gln Pro Pro Tyr Tyr Asn Val Thr Ala Met Asn Val Pro Ile Ala  
325 330 335

Val Trp Asn Gly Gly Lys Asp Leu Leu Ala Asp Pro Gln Asp Val Gly  
340 345 350

Leu Leu Leu Pro Lys Leu Pro Asn Leu Ile Tyr His Lys Glu Ile Pro  
355 360 365

Phe Tyr Asn His Leu Asp Phe Ile Trp Ala Met Asp Ala Pro Gln Glu  
370 375 380

Val Tyr Asn Asp Ile Val Ser Met Ile Ser Glu Asp Lys Lys  
385 390 395

<210> 76  
 <211> 760  
 <212> PRT  
 <213> Mus sp.

<400> 76

Met Ala Leu Pro Ser Leu Gly Gln Asp Ser Trp Ser Leu Leu Arg Val  
 1 5 10 15

Phe Phe Phe Gln Leu Phe Leu Leu Pro Ser Leu Pro Pro Ala Ser Gly  
 20 25 30

Thr Gly Gly Gln Gly Pro Met Pro Arg Val Lys Tyr His Ala Gly Asp  
 35 40 45

Gly His Arg Ala Leu Ser Phe Phe Gln Gln Lys Gly Leu Arg Asp Phe  
 50 55 60

Asp Thr Leu Leu Leu Ser Asp Asp Gly Asn Thr Leu Tyr Val Gly Ala  
 65 70 75 80

Arg Glu Thr Val Leu Ala Leu Asn Ile Gln Asn Pro Gly Ile Pro Arg  
 85 90 95

Leu Lys Asn Met Ile Pro Trp Pro Ala Ser Glu Arg Lys Lys Thr Glu  
 100 105 110

Cys Ala Phe Lys Lys Lys Ser Asn Glu Thr Gln Cys Phe Asn Phe Ile  
 115 120 125

Arg Val Leu Val Ser Tyr Asn Ala Thr His Leu Tyr Ala Cys Gly Thr  
 130 135 140

Phe Ala Phe Ser Pro Ala Cys Thr Phe Ile Glu Leu Gln Asp Ser Leu  
 145 150 155 160

Leu Leu Pro Ile Leu Ile Asp Lys Val Met Asp Gly Lys Gly Gln Ser  
 165 170 175

Pro Leu Thr Leu Phe Thr Ser Thr Gln Ala Val Leu Val Asp Gly Met  
 180 185 190

Leu Tyr Ser Gly Thr Met Asn Asn Phe Leu Gly Ser Glu Pro Ile Leu  
 195 200 205

Met Arg Thr Leu Gly Ser His Pro Val Leu Lys Thr Asp Ile Phe Leu

210	215	220
Arg Trp Leu His Ala Asp Ala Ser Phe Val Ala Ala Ile Pro Ser Thr		
225	230	235 240
Gln Val Val Tyr Phe Phe Phe Glu Glu Thr Ala Ser Glu Phe Asp Phe		
	245	250 255
Phe Glu Glu Leu Tyr Ile Ser Arg Val Ala Gln Val Cys Lys Asn Asp		
	260	265 270
Val Gly Gly Glu Lys Leu Leu Gln Lys Lys Trp Thr Thr Phe Leu Lys		
	275	280 285
Ala Gln Leu Leu Cys Ala Gln Pro Gly Gln Leu Pro Phe Asn Ile Ile		
	290	295 300
Arg His Ala Val Leu Leu Pro Ala Asp Ser Pro Ser Val Ser Arg Ile		
305	310	315 320
Tyr Ala Val Phe Thr Ser Gln Trp Gln Val Gly Gly Thr Arg Ser Ser		
	325	330 335
Ala Val Cys Ala Phe Ser Leu Thr Asp Ile Glu Arg Val Phe Lys Gly		
	340	345 350
Lys Tyr Lys Glu Leu Asn Lys Glu Thr Ser Arg Trp Thr Thr Tyr Arg		
	355	360 365
Gly Ser Glu Val Ser Pro Arg Pro Gly Ser Cys Ser Met Gly Pro Ser		
	370	375 380
Ser Asp Lys Ala Leu Thr Phe Met Lys Asp His Phe Leu Met Asp Glu		
385	390	395 400
His Val Val Gly Thr Pro Leu Leu Val Lys Ser Gly Val Glu Tyr Thr		
	405	410 415
Arg Leu Ala Val Glu Ser Ala Arg Gly Leu Asp Gly Ser Ser His Val		
	420	425 430
Val Met Tyr Leu Gly Thr Ser Thr Gly Pro Leu His Lys Ala Val Val		
	435	440 445
Pro Gln Asp Ser Ser Ala Tyr Leu Val Glu Glu Ile Gln Leu Ser Pro		
	450	455 460
Asp Ser Glu Pro Val Arg Asn Leu Gln Leu Ala Pro Ala Gln Gly Ala		

465		470		475		480
Val Phe Ala Gly Phe Ser Gly Gly Ile Trp Arg Val Pro Arg Ala Asn						
	485			490		495
Cys Ser Val Tyr Glu Ser Cys Val Asp Cys Val Leu Ala Arg Asp Pro						
	500			505		510
His Cys Ala Trp Asp Pro Glu Ser Arg Leu Cys Ser Leu Leu Ser Gly						
	515			520		525
Ser Thr Lys Pro Trp Lys Gln Asp Met Glu Arg Gly Asn Pro Glu Trp						
	530			535		540
Val Cys Thr Arg Gly Pro Met Ala Arg Ser Pro Arg Arg Gln Ser Pro						
	545			550		555
						560
Pro Gln Leu Ile Lys Glu Val Leu Thr Val Pro Asn Ser Ile Leu Glu						
				565		570
						575
Leu Arg Cys Pro His Leu Ser Ala Leu Ala Ser Tyr His Trp Ser His						
	580			585		590
Gly Arg Ala Lys Ile Ser Glu Ala Ser Ala Thr Val Tyr Asn Gly Ser						
	595			600		605
Leu Leu Leu Leu Pro Gln Asp Gly Val Gly Gly Leu Tyr Gln Cys Val						
	610			615		620
Ala Thr Glu Asn Gly Tyr Ser Tyr Pro Val Val Ser Tyr Trp Val Asp						
	625			630		635
						640
Ser Gln Asp Gln Pro Leu Ala Leu Asp Pro Glu Leu Ala Gly Val Pro						
	645			650		655
Arg Glu Arg Val Gln Val Pro Leu Thr Arg Val Gly Gly Gly Ala Ser						
	660			665		670
Met Ala Ala Gln Arg Ser Tyr Trp Pro His Phe Leu Ile Val Thr Val						
	675			680		685
Leu Leu Ala Ile Val Leu Leu Gly Val Leu Thr Leu Leu Leu Ala Ser						
	690			695		700
Pro Leu Gly Ala Leu Arg Ala Arg Gly Lys Val Gln Gly Cys Gly Met						
	705			710		715
						720
Leu Pro Pro Arg Glu Lys Ala Pro Leu Ser Arg Asp Gln His Leu Gln						

725

730

735

Pro Ser Lys Asp His Arg Thr Ser Ala Ser Asp Val Asp Ala Asp Asn  
 740 745 750

Asn His Leu Gly Ala Glu Val Ala  
 755 760

&lt;210&gt; 77

&lt;211&gt; 3046

&lt;212&gt; DNA

&lt;213&gt; Mus sp.

&lt;400&gt; 77

ctcggacgcc tgggttaggg gtctgtactg ctggggaacc atctggtgac catctcaggc 60  
 tgaccatggc cctaccatcc ctgggccagg actcatggag tctcctgcgt gtttttttct 120  
 tccaactcct cctgctgcc aactgccac ctgcttctgg gactggtggt caggggcca 180  
 tgcccagagt caaataccat gctggagacg ggcacagggc cctcagcttc ttccaacaaa 240  
 aaggcctccg agactttgac acgctgctcc tgagtgcga tggcaacact ctctatgtgg 300  
 gggctcgaga gaccgtcctg gccttgaata tccagaaccc aggaatccca aggctaaaga 360  
 acatgatacc ctggccagcc agtgagagaa aaaagaccga atgtgccttt aagaagaaga 420  
 gcaatgagac acagtgtttc aacttcattc gagtctggt ctcttacaat gctactcacc 480  
 tctatgcctg tgggaccttt gccttcagcc ctgcctgtac cttcattgaa ctccaagatt 540  
 ccctcctgtt gccatctttg atagacaagg tcatggacgg gaagggccaa agccctttga 600  
 ccctgttcac aagcacacaa gctgtcttgg tcatgggat gctttattcc ggcaccatga 660  
 acaacttcct gggcagcgag cccatctga tgcggacact gggatcccat cctgtttctca 720  
 agactgacat cttcttacgc tggtgcacg cggatgcctc ctctgtggca gccattccat 780  
 ccaccaggt cgtctatttc ttctttgagg agacagccag cgagtttgac ttctttgaag 840  
 agctgtatat atccagggtg gctcaagtct gcaagaacga cgtgggcggt gaaaagctgc 900  
 tgcaagaaga gtggaccacc ttctcaaaag ccagttgct ctgcgctcag ccagggcagc 960  
 tgccattcaa catcatccgc cagcgggtcc tgcctcccgc cgattctccc tctgtttccc 1020  
 gcatctacgc agtctttacc tcccagtggt aggttggcgg gaccaggagc tcagcagtct 1080  
 gtgccttctc tctcacggac attgagcgag tctttaaagg gaagtacaag gagctgaaca 1140  
 aggagacctc ccgctggacc acttacggg gctcagaggt cagcccagg ccaggcagtt 1200  
 gctccatggg cccctcctct gacaaagcct tgaccttcat gaaggacct tttctgatgg 1260  
 atgagcacgt ggtaggaaca cccctgctgg tgaagtctgg tgtggagtac acacggcttg 1320  
 ctgtggagtc agctcggggc ctfgatggga gcagccatgt ggtcatgtat ctgggtacct 1380  
 ccacgggtcc cctgcacaag gctgtggtgc ctccaggacag cagtgttat ctctgtggag 1440  
 agattcagct gagccctgac tctgagcctg ttccgaaacct gcagctggcc cccgcccagg 1500  
 gtgcagtggt tgcaggcttc tctggaggca tctggagagt tcccagggcc aattgcagtg 1560  
 tctacgagag ctgtgtggac tgtgtgcttg ccagggaccc tcaactgtgc tgggacctg 1620  
 aatcaagact ctgcagcctt ctgtctggct ctaccaagcc ttggaagcag gacatggaac 1680  
 gcggcaaccc ggagtgggta tgcacccgtg gcccattggc caggagcccc cggcgtcaga 1740  
 gccccctca actaattaaa gaagtcctga cagtcccaa ctccatcctg gagctgcgct 1800  
 gccccacct gtcagcactg gcctcttacc actggagtca tggccgagcc aaaatctcag 1860  
 aagcctctgc taccgtctac aatggctccc tcttctgctg gccgcaggat ggtgtcgggg 1920  
 gcctctacca gtgtgtggcg actgagaacg gctactcata ccctgtggtc tcctattggg 1980

tagacagcca ggaccagccc ctggcgctgg accctgagct ggcgggcggt ccccgtaggc 2040  
 gtgtgcaggt cccgctgacc agggctcgag gcgagagcttc catggctgcc cagcgggtcct 2100  
 actggcccca ttttctcatc gttaccgtcc tcctggccat cgtgctcctg ggagtgtcct 2160  
 ctctcctcct cgcttcccca ctggggggcg tgcgggctcg gggttaagggt cagggctgtg 2220  
 ggatgctgcc ccccagggaa aaggctccac tgagcagggg ccagcacctc cagccctcca 2280  
 aggaccacag gacctctgcc agtgacgtag atgccgacaa caaccatctg ggcgccgaag 2340  
 tggcttaaag agggacacag atccgcagct gagcagagca agccactggc cttgttggt 2400  
 atgccaggca cagtgcact ctgaccaggg taggaggctc tcctgctaac gtgtgtcacc 2460  
 tacagcacc cagtaggtcct cccctgtggg actctcttct gcaagcacat tgggtgtct 2520  
 ccatacctgt acttgtgtgt tgacaggaag agccagacag gtttctttga ttttgattga 2580  
 cccaagagcc ctgcctgtaa caaacgtgct ccaggagacc atgaaagggt tggctgtctg 2640  
 ggattctgtg gtgacaaaacc taagcatccg agcaagctgg ggctattcct gcaaactcca 2700  
 tcctgaacgc tgtcactcta gaagcagctg ctgctttgaa caccagccca cctccttcc 2760  
 caagagtctc tatggagttg gcccttgtg tttcctttac cagtcggggc atactgtttg 2820  
 ggaagtcac tctgaagtct aaccaccttc cttcttggtt cagtttggac agattgttat 2880  
 tattgtctct gccctggcta gaatgggggc ataactctgag ccttggtccc ttgtccagt 2940  
 tggctgacc ttgacctct ccttcctcct ccctttgttt tgggattcag aaaactgctt 3000  
 gtcacagaca atttattttt tattaaaaaa gatataagct ttaaag 3046

<210> 78

<211> 1436

<212> PRT

<213> Bos sp.

<400> 78

Met Ala Leu Gly Arg His Leu Ser Leu Arg Gly Leu Cys Val Leu Leu  
 1 5 10 15

Leu Gly Thr Met Val Gly Gly Gln Ala Leu Glu Leu Arg Leu Lys Asp  
 20 25 30

Gly Val His Arg Cys Glu Gly Arg Val Glu Val Lys His Gln Gly Glu  
 35 40 45

Trp Gly Thr Val Asp Gly Tyr Arg Trp Thr Leu Lys Asp Ala Ser Val  
 50 55 60

Val Cys Arg Gln Leu Gly Cys Gly Ala Ala Ile Gly Phe Pro Gly Gly  
 65 70 75 80

Ala Tyr Phe Gly Pro Gly Leu Gly Pro Ile Trp Leu Leu Tyr Thr Ser  
 85 90 95

Cys Glu Gly Thr Glu Ser Thr Val Ser Asp Cys Glu His Ser Asn Ile  
 100 105 110

Lys Asp Tyr Arg Asn Asp Gly Tyr Asn His Gly Arg Asp Ala Gly Val  
 115 120 125

Val Cys Ser Gly Phe Val Arg Leu Ala Gly Gly Asp Gly Pro Cys Ser  
130 135 140

Gly Arg Val Glu Val His Ser Gly Glu Ala Trp Ile Pro Val Ser Asp  
145 150 155 160

Gly Asn Phe Thr Leu Ala Thr Ala Gln Ile Ile Cys Ala Glu Leu Gly  
165 170 175

Cys Gly Lys Ala Val Ser Val Leu Gly His Glu Leu Phe Arg Glu Ser  
180 185 190

Ser Ala Gln Val Trp Ala Glu Glu Phe Arg Cys Glu Gly Glu Glu Pro  
195 200 205

Glu Leu Trp Val Cys Pro Arg Val Pro Cys Pro Gly Gly Thr Cys His  
210 215 220

His Ser Gly Ser Ala Gln Val Val Cys Ser Ala Tyr Ser Glu Val Arg  
225 230 235 240

Leu Met Thr Asn Gly Ser Ser Gln Cys Glu Gly Gln Val Glu Met Asn  
245 250 255

Ile Ser Gly Gln Trp Arg Ala Leu Cys Ala Ser His Trp Ser Leu Ala  
260 265 270

Asn Ala Asn Val Ile Cys Arg Gln Leu Gly Cys Gly Val Ala Ile Ser  
275 280 285

Thr Pro Gly Gly Pro His Leu Val Glu Glu Gly Asp Gln Ile Leu Thr  
290 295 300

Ala Arg Phe His Cys Ser Gly Ala Glu Ser Phe Leu Trp Ser Cys Pro  
305 310 315 320

Val Thr Ala Leu Gly Gly Pro Asp Cys Ser His Gly Asn Thr Ala Ser  
325 330 335

Val Ile Cys Ser Gly Asn Gln Ile Gln Val Leu Pro Gln Cys Asn Asp  
340 345 350

Ser Val Ser Gln Pro Thr Gly Ser Ala Ala Ser Glu Asp Ser Ala Pro  
355 360 365

Tyr Cys Ser Asp Ser Arg Gln Leu Arg Leu Val Asp Gly Gly Gly Pro  
370 375 380

Cys Ala Gly Arg Val Glu Ile Leu Asp Gln Gly Ser Trp Gly Thr Ile  
385 390 395 400

Cys Asp Asp Gly Trp Asp Leu Asp Asp Ala Arg Val Val Cys Arg Gln  
405 410 415

Leu Gly Cys Gly Glu Ala Leu Asn Ala Thr Gly Ser Ala His Phe Gly  
420 425 430

Ala Gly Ser Gly Pro Ile Trp Leu Asp Asn Leu Asn Cys Thr Gly Lys  
435 440 445

Glu Ser His Val Trp Arg Cys Pro Ser Arg Gly Trp Gly Gln His Asn  
450 455 460

Cys Arg His Lys Gln Asp Ala Gly Val Ile Cys Ser Glu Phe Leu Ala  
465 470 475 480

Leu Arg Met Val Ser Glu Asp Gln Gln Cys Ala Gly Trp Leu Glu Val  
485 490 495

Phe Tyr Asn Gly Thr Trp Gly Ser Val Cys Arg Asn Pro Met Glu Asp  
500 505 510

Ile Thr Val Ser Thr Ile Cys Arg Gln Leu Gly Cys Gly Asp Ser Gly  
515 520 525

Thr Leu Asn Ser Ser Val Ala Leu Arg Glu Gly Phe Arg Pro Gln Trp  
530 535 540

Val Asp Arg Ile Gln Cys Arg Lys Thr Asp Thr Ser Leu Trp Gln Cys  
545 550 555 560

Pro Ser Asp Pro Trp Asn Tyr Asn Ser Cys Ser Pro Lys Glu Glu Ala  
565 570 575

Tyr Ile Trp Cys Ala Asp Ser Arg Gln Ile Arg Leu Val Asp Gly Gly  
580 585 590

Gly Arg Cys Ser Gly Arg Val Glu Ile Leu Asp Gln Gly Ser Trp Gly  
595 600 605

Thr Ile Cys Asp Asp Arg Trp Asp Leu Asp Asp Ala Arg Val Val Cys  
610 615 620

Lys Gln Leu Gly Cys Gly Glu Ala Leu Asp Ala Thr Val Ser Ser Phe  
625 630 635 640



Asn	His	Thr	Gln	Val	Leu	Pro	Gln	Cys	Asn	Asp	Phe	Leu	Ser	Gln	Pro	900	905	910
Ala	Gly	Ser	Ala	Ala	Ser	Glu	Glu	Ser	Ser	Pro	Tyr	Cys	Ser	Asp	Ser	915	920	925
Arg	Gln	Leu	Arg	Leu	Val	Asp	Gly	Gly	Gly	Pro	Cys	Gly	Gly	Arg	Val	930	935	940
Glu	Ile	Leu	Asp	Gln	Gly	Ser	Trp	Gly	Thr	Ile	Cys	Asp	Asp	Asp	Trp	945	950	955
Asp	Leu	Asp	Asp	Ala	Arg	Val	Val	Cys	Arg	Gln	Leu	Gly	Cys	Gly	Glu	965	970	975
Ala	Leu	Asn	Ala	Thr	Gly	Ser	Ala	His	Phe	Gly	Ala	Gly	Ser	Gly	Pro	980	985	990
Ile	Trp	Leu	Asp	Asp	Leu	Asn	Cys	Thr	Gly	Lys	Glu	Ser	His	Val	Trp	995	1000	1005
Arg	Cys	Pro	Ser	Arg	Gly	Trp	Gly	Arg	His	Asp	Cys	Arg	His	Lys	Glu	1010	1015	1020
Asp	Ala	Gly	Val	Ile	Cys	Ser	Glu	Phe	Leu	Ala	Leu	Arg	Met	Val	Ser	1025	1030	1035
Glu	Asp	Gln	Gln	Cys	Ala	Gly	Trp	Leu	Glu	Val	Phe	Tyr	Asn	Gly	Thr	1045	1050	1055
Trp	Gly	Ser	Val	Cys	Arg	Ser	Pro	Met	Glu	Asp	Ile	Thr	Val	Ser	Val	1060	1065	1070
Ile	Cys	Arg	Gln	Leu	Gly	Cys	Gly	Asp	Ser	Gly	Ser	Leu	Asn	Thr	Ser	1075	1080	1085
Val	Gly	Leu	Arg	Glu	Gly	Ser	Arg	Pro	Arg	Trp	Val	Asp	Leu	Ile	Gln	1090	1095	1100
Cys	Arg	Lys	Met	Asp	Thr	Ser	Leu	Trp	Gln	Cys	Pro	Ser	Gly	Pro	Trp	1105	1110	1115
Lys	Tyr	Ser	Ser	Cys	Ser	Pro	Lys	Glu	Glu	Ala	Tyr	Ile	Ser	Cys	Glu	1125	1130	1135
Gly	Arg	Arg	Pro	Lys	Ser	Cys	Pro	Thr	Ala	Ala	Ala	Cys	Thr	Asp	Arg	1140	1145	1150

Glu Lys Leu Arg Leu Arg Gly Gly Asp Ser Glu Cys Ser Gly Arg Val  
 1155 1160 1165

Glu Val Trp His Asn Gly Ser Trp Gly Thr Val Cys Asp Asp Ser Trp  
 1170 1175 1180

Ser Leu Ala Glu Ala Glu Val Val Cys Gln Gln Leu Gly Cys Gly Gln  
 1185 1190 1195 1200

Ala Leu Glu Ala Val Arg Ser Ala Ala Phe Gly Pro Gly Asn Gly Ser  
 1205 1210 1215

Ile Trp Leu Asp Glu Val Gln Cys Gly Gly Arg Glu Ser Ser Leu Trp  
 1220 1225 1230

Asp Cys Val Ala Glu Pro Trp Gly Gln Ser Asp Cys Lys His Glu Glu  
 1235 1240 1245

Asp Ala Gly Val Arg Cys Ser Gly Val Arg Thr Thr Leu Pro Thr Thr  
 1250 1255 1260

Thr Ala Gly Thr Arg Thr Thr Ser Asn Ser Leu Pro Gly Ile Phe Ser  
 1265 1270 1275 1280

Leu Pro Gly Val Leu Cys Leu Ile Leu Gly Ser Leu Leu Phe Leu Val  
 1285 1290 1295

Leu Val Ile Leu Val Thr Gln Leu Leu Arg Trp Arg Ala Glu Arg Arg  
 1300 1305 1310

Ala Leu Ser Ser Tyr Glu Asp Ala Leu Ala Glu Ala Val Tyr Glu Glu  
 1315 1320 1325

Leu Asp Tyr Leu Leu Thr Gln Lys Glu Gly Leu Gly Ser Pro Asp Gln  
 1330 1335 1340

Met Thr Asp Val Pro Asp Glu Asn Tyr Asp Asp Ala Glu Glu Val Pro  
 1345 1350 1355 1360

Val Pro Gly Thr Pro Ser Pro Ser Gln Gly Asn Glu Glu Glu Val Pro  
 1365 1370 1375

Pro Glu Lys Glu Asp Gly Val Arg Ser Ser Gln Thr Gly Ser Phe Leu  
 1380 1385 1390

Asn Phe Ser Arg Glu Ala Ala Asn Pro Gly Glu Gly Glu Glu Ser Phe  
 1395 1400 1405

Trp Leu Leu Gln Gly Lys Lys Gly Asp Ala Gly Tyr Asp Asp Val Glu  
 1410 1415 1420

Leu Ser Ala Leu Gly Thr Ser Pro Val Thr Phe Ser  
 1425 1430 1435

<210> 79  
 <211> 4308  
 <212> DNA  
 <213> Bos sp.

<400> 79

```

atggctctgg gcagacacct ctccctgcgg ggactctgtg tctcctcct cggcaccatg 60
gtgggtgggc aagctctgga gctgaggttg aaggatggag tccatcgctg tgaggggaga 120
gtggaagtga agcaccaagg agaatggggc acagtggatg gttacagggtg gacattgaag 180
gatgcatctg tagtgtgcag acagctgggg tgtggagctg ccattggttt tcctggaggg 240
gcttattttg ggccaggact tggcccatt tggcttttgt atacttcctg tgaagggaca 300
gagtcaactg tcagtgactg tgagcattct aatattaaag actatcgtaa tgatggctat 360
aatcatgggc gggatgctgg agtagtctgc tcaggatttg tgcgtctggc tggaggggat 420
ggaccctgct cagggcgagt agaagtgcac tctggagaag cttggatccc agtgtctgat 480
gggaacttca cacttgccac tgcccagatc atctgtgcag agttgggttg tggcaaggct 540
gtgtctgtcc tgggacatga gctcttcaga gaggccagtg ccaggtctg ggctgaagag 600
ttcagggtgtg agggggagga gcctgagctc tgggtctgcc ccagagtgcc ctgtccaggg 660
ggcacgtgtc accacagtgg atctgtctag gttgtttgtt cagcatactc agaagtcagg 720
ctcatgacaa acggctcctc tcagtgtgaa gggcaggtgg agatgaacat ttctggacaa 780
tggagagcgc tctgtgctcc ccaactggag ctggccaatg ccaatgttat ctgtcgtcag 840
ctcggctgtg gaggttgcat ctccaccccc ggaggaccac acttggtgga agaaggtgat 900
cagatcctaa cagcccgatt tcaactgctc tgggctgagt ccttcctgtg gaggttgtct 960
gtgactgcc tgggtggtcc tgactgttcc catggcaaca cagcctctgt gatctgtc 1020
ggaaaccaga tccaggtgct tccccagtg aacgactccg tgtctcaacc tacaggctct 1080
gcggcctcag aggacagcgc cccctactgc tcagacagca ggcagctccg cctggtggac 1140
gggggcggtc cctgcgccgg gagagtggag atccttgacc agggctcctg gggcaccatc 1200
tgtgatgacg gctgggacct ggacgatgcc cgcgtggtgt gcaggcagct gggctgtgga 1260
gaagccctca atgccacggg gtctgtctac ttcggggcag gatcagggcc catctggttg 1320
gacaacttga actgcacagg aaaggagtcc cacgtgtgga ggtgcccttc ccggggctgg 1380
gggcagcaca actgcagaca caagcaggac gcgggggtca tctgtctaga gttcctggcc 1440
ctcaggatgg tgagtgagga ccagcagtgt gctgggtggc tggagtttt ctacaatggg 1500
acctggggca gtgtctgcc taaccccatg gaagacatca ctgtgtccac gatctgcaga 1560
cagcttggct gtggggacag tggaaacctc aactctcttg ttgctcttag agaaggtttt 1620
aggccacagt ggggtgtag aatccagtgt cggaaaactg acacctctct ctggcagtgt 1680
ccttctgacc cttggaatta caactcatgc tctccaaagg aggaagccta tatctggtgt 1740
gcagacagca gacagatccg cctggtggat ggaggtggc gctgtctctg gagagtggag 1800
atccttgacc agggctcctg gggcaccatc tgtgatgacc gctgggacct ggacgatgcc 1860
cgtgtggtgt gcaagcagct gggctgtgga gaagccctgg acgccactgt ctcttccttc 1920
ttcgggacgg gatcagggcc catctggctg gatgaagtga actgcagagg agaggagtcc 1980
caagtatgga ggtgcccttc ctggggatgg cggaacaca actgcaatca tcaagaagat 2040

```

```

gcaggagtca tctgctcagg atttgtgcgt ctggctggag gagatggacc ctgctcaggg 2100
cgagtagaag tgcattctgg agaagcctgg accccagtgt ctgatggaaa cttcacactc 2160
cccactgccc aggtcatctg tgcagagctg ggatgtggca aggctgtgtc tgtcctggga 2220
cacatgccat tcagagagtc cgatggccag gtctgggctg aagagttcag gtgtgatggg 2280
ggggagcctg agctctggtc ctgccccaga gtgccctgtc caggaggcac atgtctccac 2340
agtggagctg ctcaggttgt ctgttcagtg tacacagaag tccagcttat gaaaaacggc 2400
acctctcaat gtgaggggca ggtggagatg aagatctctg gacgatggag agcgctctgt 2460
gcctcccact ggagtctggc caatgccaat gttgtctgtc gtcagctcgg ctgtggagtc 2520
gccatctcca ccccagagg accacacttg gtggaaggag gtgatcagat ctcaacagcc 2580
caatttcact gctcaggggc tgagtccttc ctgtggagtt gtcctgtgac tgccttgggt 2640
gggcctgact gttcccatgg caacacagcc tctgtgatct gtcaggaaa ccacaccag 2700
gtgctgcccc agtgcaacga cttcctgtct caacctgcag gctctgcggc ctcagaggag 2760
agttctccct actgctcaga cagcaggcag ctccgcctgg tggacggggg cggctccctgc 2820
ggcgggagag tggagatcct tgaccagggc tcctggggca ccatctgtga tgatgactgg 2880
gacctggacg atgcccggtg ggtgtgcagg cagctgggct gtggagaagc cctcaatgcc 2940
acggggtctg ctcacttcgg ggcaggatca gggcccatct ggctggacga cctgaactgc 3000
acaggaaagg agtcccacgt gtggaggtgc ccttcccggg gctgggggcg gcacgactgc 3060
agacacaagg aggacgccgg ggtcatctgc tcagagttcc tggccctcag gatggtgagc 3120
gaggaccagc agtgtgctgg gtggctggag gttttctaca acgggacctg gggcagtgtc 3180
tgccgcagcc ccatggaaga tatcactgtg tccgtgatct gcagacagct tggatgtggg 3240
gacagtggaa gtctcaacac ctctgttggg ctccagggaag gttctagacc ccggtgggta 3300
gatttaattc agtgtcggaa aatggatacc tctctctggc agtgtccttc tggcccatgg 3360
aaatacagtt catgctctcc aaaggaggaa gcctacatct catgtgaagg aagaagacc 3420
aagagctgtc caactgctgc cgcctgcaca gacagagaga agctccgcct caggggagga 3480
gacagcgagt gctcagggcg ggtggaggtg tggcacaacg gtcctgggg caccgtgtgc 3540
gatgactcct ggagcctggc agaggctgag gtggtgtgtc agcagctggg ctgtggccag 3600
gccctggaag ccgtgcggtc tgcagcattt ggccctggaa atgggagcat ctggctggac 3660
gaggtgcagt gcgggggccc ggagtcctcc ctgtgggact gtgttgcgga gccctggggg 3720
cagagcgact gcaagcacga ggaggatgct ggtgtgaggt gctctggtgt aaggacaaca 3780
ttgcccacga ccacagcagg gaccagaaca acctcaaatt ctctccctgg catcttctcc 3840
ctgcctgggg ttctctgcct tatcctgggg togtttctct tcctggctct cgtcatcctg 3900
gtgactcagc tactcagatg gagagcagag cgcagagcct tatccagcta tgaagatgct 3960
cttgctgaag ctgtgtatga ggagctcgat taccttctga cacagaagga aggtctgggc 4020
agcccagatc agatgactga tgtccctgat gaaaattatg atgatgctga agaagtacca 4080
gtgcctggaa ctcttctcc ctctcagggg aatgaggagg aagtgcccc agagaaggag 4140
gacggggtga ggtcctctca gacaggctct ttctgaact tctccagaga ggcagcta 4200
cctggggaag gagaagagag cttctggctg ctccagggga agaaagggga tgctgggtat 4260
gatgatgttg aactcagtgc cctgggaaca tcccagtgat ctttctcg 4308

```